

ISSN 2518-170X (Online)

ISSN 2224-5278 (Print)



«ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫ» РҚБ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

РОО «НАЦИОНАЛЬНОЙ
АКАДЕМИИ НАУК РЕСПУБЛИКИ
КАЗАХСТАН»

N E W S

OF THE NATIONAL ACADEMY
OF SCIENCES OF THE REPUBLIC
OF KAZAKHSTAN

SERIES

OF GEOLOGY AND TECHNICAL SCIENCES

3 (471)

MAY – JUNE 2025

THE JOURNAL WAS FOUNDED IN 1940

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of geology and technical sciences scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of geology and technical sciences in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of geology and engineering sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы «ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы» ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді геология және техникалық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия геологии и технических наук» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК. Серия геологии и технических наук в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по геологии и техническим наукам для нашего сообщества.

EDITOR-IN-CHIEF

ZHURINOV Murat Zhurinovich, Doctor of Chemical Sciences, Professor, Academician of NAS RK, President of National Academy of Sciences of the Republic of Kazakhstan, RPA, General Director of JSC "D.V. Sokolsky Institute of Fuel, Catalysis and Electrochemistry" (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>, <https://www.webofscience.com/wos/author/record/2017489>

DEPUTY EDITOR-IN-CHIEF

ABSADYKOV Bakhyt Narikbayevich, Doctor of Technical Sciences, Professor, Academician of NAS RK, Satbayev University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=6504694468>, <https://www.webofscience.com/wos/author/record/2411827>

EDITORIAL BOARD:

ABSAMETOV Malis Kudysovich, (Deputy Editor-in-Chief), Doctor of Geological and Mineralogical Sciences, Professor, Academician of NAS RK, Director of the Akhmedsafin Institute of Hydrogeology and Geoecology (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=56955769200>, <https://www.webofscience.com/wos/author/record/1937883>

ZHOLTAEV Geroy Zholtaevich, Doctor of Geological and Mineralogical Sciences, Professor, Honorary Academician of NAS RK (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=57112610200>, <https://www.webofscience.com/wos/author/record/1939201>

SNOW Daniel, PhD, Associate Professor, Director, Aquatic Sciences Laboratory, University of Nebraska (Nebraska, USA), <https://www.scopus.com/authid/detail.uri?authorId=7103259215>, <https://www.webofscience.com/wos/author/record/1429613>

SELTMAN Reimar, PhD, Head of Petrology and Mineral Deposits Research in the Earth Sciences Department, Natural History Museum (London, England), <https://www.scopus.com/authid/detail.uri?authorId=55883084800>, <https://www.webofscience.com/wos/author/record/1048681>

PANFILOV Mikhail Borisovich, Doctor of Technical Sciences, Professor at the University of Nancy (Nancy, France), <https://www.scopus.com/authid/detail.uri?authorId=7003436752>, <https://www.webofscience.com/wos/author/record/1230499>

SHEN Ping, PhD, Deputy Director of the Mining Geology Committee of the Chinese Geological Society, Member of the American Association of Economic Geologists (Beijing, China), <https://www.scopus.com/authid/detail.uri?authorId=57202873965>, <https://www.webofscience.com/wos/author/record/1753209>

FISCHER Axel, PhD, Associate Professor, Technical University of Dresden (Dresden, Berlin), <https://www.scopus.com/authid/detail.uri?authorId=35738572100>, <https://www.webofscience.com/wos/author/record/2085986>

AGABEKOV Vladimir Enokovich, Doctor of Chemical Sciences, Academician of NAS of Belarus, Honorary Director of the Institute of Chemistry of New Materials (Minsk, Belarus), <https://www.scopus.com/authid/detail.uri?authorId=7004624845>

CATALIN Stefan, PhD, Associate Professor, Technical University of Dresden, Germany, <https://www.scopus.com/authid/detail.uri?authorId=35203904500>, <https://www.webofscience.com/wos/author/record/1309251>

Jay Sagin, PhD, Associate Professor, Nazarbayev University (Astana, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=57204467637>, <https://www.webofscience.com/wos/author/record/907886>

FRATTINI Paolo, PhD, Associate Professor, University of Milano - Bicocca (Milan, Italy), <https://www.scopus.com/authid/detail.uri?authorId=56538922400>

NURPEISOVA Marzhan Baysanovna – Doctor of Technical Sciences, Professor of Satbayev University, (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=57202218883>, <https://www.webofscience.com/wos/author/record/AAD-1173-2019>

RATOV Boranbay Tovbasarovich, Doctor of Technical Sciences, Professor, Head of the Department of Geophysics and Seismology, Satbayev University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=55927684100>, <https://www.webofscience.com/wos/author/record/1993614>

RONNY Berndtsson, Professor at the Center of Promising Middle Eastern Research, Lund University (Sweden), <https://www.scopus.com/authid/detail.uri?authorId=7005388716>, <https://www.webofscience.com/wos/author/record/1324908>

MIRLAS Vladimir, Faculty chemical engineering and Oriental research center, Ariel University, (Israel), <https://www.scopus.com/authid/detail.uri?authorId=8610969300>, <https://www.webofscience.com/wos/author/record/53680261>

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Owner: RPA «National Academy of Sciences of the Republic of Kazakhstan» (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Social Development of the Republic of Kazakhstan **No. KZ39VPY00025420**, issued 29.07.2020.

Thematic scope: *geology, hydrogeology, geography, mining and chemical technologies of oil, gas and metals*

Periodicity: 6 times a year.

<http://www.geology-technical.kz/index.php/en/>

© National Academy of Sciences of the Republic of Kazakhstan, 2025

БАС РЕДАКТОР

ЖҰРЫНОВ Мұрат Жұрыңұлы, химия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, РКБ «Қазақстан Республикасы Ұлттық Ғылым академиясының» президенті, АҚ «Д.В. Сокольский атындағы отын, катализ және электрохимия институтының» бас директоры (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>, <https://www.webofscience.com/wos/author/record/2017489>

БАС РЕДАКТОРДЫҢ ОРЫНБАСАРЫ:

АБСАДЫҚОВ Бақыт Нәрікбайұлы, техника ғылымдарының докторы, профессор, ҚР ҰҒА академигі, Қ.И. Сәтбаев атындағы Қазақ ұлттық техникалық зерттеу университеті (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=6504694468>, <https://www.webofscience.com/wos/author/record/2411827>

РЕДАКЦИЯ АЛҚАСЫ:

ӘБСӘМЕТОВ Мәліс Құдысулы (бас редактордың орынбасары), геология-минералогия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, У.М. Ахмедсафин атындағы Гидрогеология және геоэкология институтының директоры, (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=56955769200>, <https://www.webofscience.com/wos/author/record/1937883>

ЖОЛТАЕВ Герой Жолтайұлы, геология-минералогия ғылымдарының докторы, профессор, ҚР ҰҒА құрметті академигі, (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=57112610200>, <https://www.webofscience.com/wos/author/record/1939201>

СНОУ Дэниел, PhD, қауымдастырылған профессор, Небраска университетінің Су ғылымдары зертханасының директоры, (Небраска штаты, АҚШ), <https://www.scopus.com/authid/detail.uri?authorId=7103259215>, <https://www.webofscience.com/wos/author/record/1429613>

ЗЕЛЪГМАН Раймар, PhD, Жер туралы ғылымдар бөлімінің петрология және пайдалы қазбалар кен орындары саласындағы зерттеулерінің жетекшісі, Табиғи тарих мұражайы, (Лондон, Ұлыбритания), <https://www.scopus.com/authid/detail.uri?authorId=55883084800>, <https://www.webofscience.com/wos/author/record/1048681>

ПАНФИЛОВ Михаил Борисович, техника ғылымдарының докторы, Нанси университетінің профессоры, (Нанси, Франция), <https://www.scopus.com/authid/detail.uri?authorId=7003436752>, <https://www.webofscience.com/wos/author/record/1230499>

ШЕН Пин, PhD, Қытай геологиялық қоғамының Тау-кен геологиясы комитеті директорының орынбасары, Американдық экономикалық геологтар қауымдастығының мүшесі, (Бейжін, Қытай), <https://www.scopus.com/authid/detail.uri?authorId=57202873965>, <https://www.webofscience.com/wos/author/record/1753209>

ФИШЕР Аксель, қауымдастырылған профессор, PhD, Дрезден техникалық университеті, (Дрезден, Берлин), <https://www.scopus.com/authid/detail.uri?authorId=35738572100>, <https://www.webofscience.com/wos/author/record/2085986>

АГАБЕКОВ Владимир Енокович, химия ғылымдарының докторы, Беларусь ҰҒА академигі, Жаңа материалдар химиясы институтының құрметті директоры, (Минск, Беларусь), <https://www.scopus.com/authid/detail.uri?authorId=7004624845>

КАТАЛИН Стефан, PhD, қауымдастырылған профессор, Техникалық университеті (Дрезден, Германия), <https://www.scopus.com/authid/detail.uri?authorId=35203904500>, <https://www.webofscience.com/wos/author/record/1309251>

САҒЫНТАЕВ Жанай, PhD, қауымдастырылған профессор, Назарбаев университеті (Астана, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=57204467637>, <https://www.webofscience.com/wos/author/record/907886>

ФРАТТИНИ Паоло, PhD, қауымдастырылған профессор, Бикокк Милан университеті, (Милан, Италия), <https://www.scopus.com/authid/detail.uri?authorId=56538922400>

НҮРПЕЙСОВА Маржан Байсанқызы – Техника ғылымдарының докторы, Қ.И. Сәтбаев атындағы Қазақ ұлттық зерттеу техникалық университетінің профессоры, (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=57202218883>, <https://www.webofscience.com/wos/author/record/AAD-1173-2019>

Ратов Боранбай Товбасарович, техника ғылымдарының докторы, профессор, «Геофизика және сейсмология» кафедрасының меңгерушісі, Қ.И. Сәтбаев атындағы Қазақ ұлттық зерттеу техникалық университеті, (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=55927684100>, <https://www.webofscience.com/wos/author/record/1993614>

РОННИ Беритссон, Лунд университетінің Таяу Шығысты перспективалы зерттеу орталығының профессоры, Лунд университетінің толық курсты профессоры, (Швеция), <https://www.scopus.com/authid/detail.uri?authorId=7005388716>, <https://www.webofscience.com/wos/author/record/1324908>

МИРЛАС Владимир, Ариэль университетінің Химиялық инженерия факультеті және Шығыс ғылыми-зерттеу орталығы, (Израиль), <https://www.scopus.com/authid/detail.uri?authorId=8610969300>, <https://www.webofscience.com/wos/author/record/53680261>

«ҚР ҰҒА» РКБ Хабарлары. Геология және техникалық ғылымдар сериясы».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктеуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РКБ (Алматы қ.).

Қазақстан Республикасының Ақпарат және қоғамдық даму министрлігінің Ақпарат комитетінде 29.07.2020 ж. берілген № **KZ39VPY00025420** мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Тақырыптық бағыты: *Геология, гидрогеология, география, тау-кен ісі, мұнай, газ және металдардың*

химиялық технологиялары

Мерзімділігі: жылына 6 рет.

<http://www.geolog-technical.kz/index.php/en/>

© «Қазақстан Республикасының Ұлттық ғылым академиясы» РКБ, 2025

ГЛАВНЫЙ РЕДАКТОР

ЖУРИНОВ Мурат Журинович, доктор химических наук, профессор, академик НАН РК, президент РОО Национальной академии наук Республики Казахстан, генеральный директор АО «Институт топлива, катализа и электрохимии им. Д.В. Сокольского» (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>, <https://www.webofscience.com/wos/author/record/2017489>

ЗАМЕСТИТЕЛЬ ГЛАВНОГО РЕДАКТОРА

АБСАДЫКОВ Бахыт Нарикбаевич, доктор технических наук, профессор, академик НАН РК, Казахский национальный исследовательский технический университет им. К.И. Сатпаева (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6504694468>, <https://www.webofscience.com/wos/author/record/2411827>

РЕДАКЦИОННАЯ КОЛЛЕГИЯ:

АБСАМЕТОВ Малис Кудысович, (заместитель главного редактора), доктор геолого-минералогических наук, профессор, академик НАН РК, директор Института гидрогеологии и геоэкологии им. У.М. Ахмедсафина (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=56955769200>, <https://www.webofscience.com/wos/author/record/1937883>

ЖОЛТАЕВ Герой Жолтаевич, доктор геологоминералогических наук, профессор, почетный академик НАН РК (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=57112610200>, <https://www.webofscience.com/wos/author/record/1939201>

СНОУ Дэниел, PhD, ассоциированный профессор, директор Лаборатории водных наук Университета Небраски (штат Небраска, США), <https://www.scopus.com/authid/detail.uri?authorId=7103259215>, <https://www.webofscience.com/wos/author/record/1429613>

ЗЕЛЪТМАНН Раймар, PhD, руководитель исследований в области петрологии и месторождений полезных ископаемых в Отделе наук о Земле Музея естественной истории (Лондон, Англия), <https://www.scopus.com/authid/detail.uri?authorId=55883084800>, <https://www.webofscience.com/wos/author/record/1048681>

ПАНФИЛОВ Михаил Борисович, доктор технических наук, профессор Университета Нанси (Нанси, Франция), <https://www.scopus.com/authid/detail.uri?authorId=7003436752>, <https://www.webofscience.com/wos/author/record/1230499>

ШЕН Пин, PhD, заместитель директора Комитета по горной геологии Китайского геологического общества, член Американской ассоциации экономических геологов (Пекин, Китай), <https://www.scopus.com/authid/detail.uri?authorId=57202873965>, <https://www.webofscience.com/wos/author/record/1753209>

ФИШЕР Аксель, ассоциированный профессор, PhD, технический университет Дрезден (Дрезден, Берлин), <https://www.scopus.com/authid/detail.uri?authorId=35738572100>, <https://www.webofscience.com/wos/author/record/2085986>

АГАБЕКОВ Владимир Енокович, доктор химических наук, академик НАН Беларуси, почетный директор Института химии новых материалов (Минск, Беларусь), <https://www.scopus.com/authid/detail.uri?authorId=7004624845>

КАТАЛИН Стефан, PhD, ассоциированный профессор, Технический университет (Дрезден, Германия), <https://www.scopus.com/authid/detail.uri?authorId=35203904500>, <https://www.webofscience.com/wos/author/record/1309251>

САГИНТАЕВ Жанай, PhD, ассоциированный профессор, Назарбаев университет (Астана, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=57204467637>, <https://www.webofscience.com/wos/author/record/907886>

ФРАТТИНИ Паоло, PhD, ассоциированный профессор, Миланский университет Биокк (Милан, Италия), <https://www.scopus.com/authid/detail.uri?authorId=56538922400>

НУРПЕЙСОВА Маржан Байсановна – доктор технических наук, профессор Казахского Национального исследовательского технического университета им. К.И. Сатпаева, (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=57202218883>, <https://www.webofscience.com/wos/author/record/AAD-1173-2019>

РАТОВ Боранбай Товбасарович, доктор технических наук, профессор, заведующий кафедрой «Геофизика и сейсмология», Казахский Национальный исследовательский технический университет им. К.И. Сатпаева, (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=55927684100>, <https://www.webofscience.com/wos/author/record/1993614>

РОННИ Берндтссон, Профессор Центра перспективных ближневосточных исследований Лундского университета, профессор (полный курс) Лундского университета, (Швеция), <https://www.scopus.com/authid/detail.uri?authorId=7005388716>, <https://www.webofscience.com/wos/author/record/1324908>

МИРЛАС Владимир, Факультет химической инженерии и Восточный научно-исследовательский центр, Университет Ариэля, (Израиль), <https://www.scopus.com/authid/detail.uri?authorId=8610969300>, <https://www.webofscience.com/wos/author/record/53680261>

«Известия РОО «НАН РК». Серия геологии и технических наук».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и общественного развития Республики Казахстан № KZ39VPY00025420, выданное 29.07.2020 г.

Тематическая направленность: *геология, гидрогеология, география, горное дело и химические технологии нефти, газа и металлов*

Периодичность: 6 раз в год.

<http://www.geolog-technical.kz/index.php/en/>

© РОО «Национальная академия наук Республики Казахстан», 2025

CONTENTS

Geology

A. Abetov, Zh. Katrenov, S. Kudaibergenova, Sh. Kisseyeva INTEGRATED GEODYNAMIC MONITORING AND RISK ASSESSMENT OF DEFORMATION PROCESSES AT THE BOZASHY NORTH OIL AND GAS FIELD.....	9
Ye. Bukayev, F. Nurbayeva, A. Bukayeva STUDY OF CHEMICAL-MINERALOGICAL COMPOSITION OF LIMESTONE-SHELL FROM THE ZHETIBAI FIELD.....	27
K.S. Dosaliyev, M.I. Karabaev, F.Kh. Aubakirova, A.M. Karabaeva, Ya.B. Kunanbayeva STRESS-STRAIN STATE CALCULATIONS FOR THE SOIL BASE OF THE SLAB FOUNDATION OF A HIGH-RISE BUILDING.....	39
A.S. Ibraim, B.N. Absadykov, S.A. Kalmaganbetov, D.B. Absadykov STUDY OF THE PROSPECTS OF USING 3D PRINTED METAL-CERAMIC ALLOYS IN ELECTRIC MOTORS.....	55
V. Ismailov, J. Bozorov, A. Khusomiddinov, E. Yadigarov, A. Mansurov DETERMINATION OF CHANGES IN SOIL PARAMETERS USING THE PLAXIS 3D PROGRAM USING REINFORCEMENT OF BORED PILES.....	69
Yu.I. Karlina, V.Y. Konyukhov, T.A. Oparina ANALYSIS OF THE INTERACTION OF TRADITIONAL AND NEW TECHNOLOGIES FOR THE EXTRACTION OF METALS FROM SUBSTANDARD RAW MATERIALS.....	83
D.M. Kirgizbaeva, T.B. Nurpeissova, A.Zh. Beisenova, T.A. Kuandykov, S.E. Tirzhanova METHOD OF RECULTIVATION OF POLLUTED SOILS WITH OIL PRODUCTS.....	96
Zh. Markabayeva, K. Koshimbayev, L. Abzhanova, Y. Orakbaev, S. Sagyndykova ANALYSIS OF MODERN METHODS FOR CONTROL AND MANAGEMENT OF THE FLOTATION PROCESS.....	109

N.A. Medeshova, D.A. Novikov, E.S. Auelkhan, A.R. Tasbolat, Sh.D. Miniskul HYDROGEOCHEMICAL FEATURES OF THE NORTH-WESTERN REGIONS OF THE TORGAY DEPRESSION IN RESPECT OF THE SEARCH FOR DEPOSITS OF STRATEGIC METALS.....	120
I.E. Nekrasova, R.V. Kononenko, M.A. Popov, M.I. Chazhaev, S.S. Khudoyorov OPTIMISATION OF DUST REGIME AND EXPLOSION SAFETY OF COAL MINES.....	139
S.H. Novruzova, I.N. Aliyev. E.V. Gadashova CONTROL OF THE FACTORS AFFECTING WELL PRODUCTIVITY.....	151
M.B. Nurpeissova, G. Meirambek, N.S. Donenbayeva, Ye.Zh. Ormambekov, R.Sh. Bek DEVELOPMENT OF METHOD FOR ASSESSING QUARRY SLOPE STABILITY USING SIDE MASSIF MAPPING.....	166
B. Orazbayev, B. Assanova, Zh. Shangitova, Zh. Moldasheva HEURISTIC APPROACH TO MULTI-CRITERIA OPTIMISATION OF A MODEL BASED DELAYED COKING PROCESS IN FUZZY ENVIRONMENT.....	179
B. Orymbetov, E. Orymbetov, G. Orymbetova, A. Khusanov, T. Orymbetov HYDRAULIC RESISTANCE OF THE ADSORBER WITH REGULAR NOZZLE.....	197
A.P. Permana, D.W.K. Baderan, R. Hutagalung, F.A. Ahmad TECTONIC GEOHISTORY OF THE GORONTALO REGION BASED ON FORAMINIFERA FOSSIL.....	207
V. Solonenko, N. Makhmetova, N. Ivanovtseva, M. Kvashnin, V. Nikolaev STABILITY OF WORKINGS OF THE CROSSHAIRS AND DRIFTS TYPE IN THE INCLINED-LAYERED ROCK MASSIF.....	220
V. Stanevich, O. Vyshar, G. Rakhimova, M. Rakhimov, S. Kovtareva TECHNOGENIC WASTE FROM COAL MINING - A PROMISING RAW MATERIAL FOR THE PRODUCTION OF BUILDING CERAMICS.....	233
Zh.K. Tukhfatov, M.K. Jexenov, Y.K. Bektay, G.S. Turysbekova, B.N. Shiderin EXPLORATION STUDIES FOR RAW CHEMICAL MINERAL RESOURCES IN THE CASPIAN BASIN SALT DOMES.....	252

**Y.A. Tynchenko, E.V. Khudyakova, V.V. Kukartsev, M.N. Stepancevich,
A.A. Stupina**

FORECASTING THE CONTENT OF RARE EARTH ELEMENTS BASED
ON GEOCHEMICAL DATA USING ENSEMBLE LEARNING

METHODS.....268

**B. Khusain, N.E. Zhumakhanova, A.Zh. Kenessary, D.N. Delikesheva,
T.D. Darzhokov**

OPTIMIZATION OF CO₂ HUFF-N-PUFF PARAMETERS FOR ENHANCED
GAS RECOVERY IN SHALE RESERVOIRS: A COMPOSITIONAL

SIMULATION STUDY.....281

NEWS of the National Academy of Sciences of the Republic of Kazakhstan
SERIES OF GEOLOGY AND TECHNICAL SCIENCES
ISSN 2224-5278
Volume 3. Number 471 (2025), 207–219

<https://doi.org/10.32014/2025.2518-170X.521>

UDC 622.276.54:622.276.72:519.876.5

© A.P. Permana*, D.W.K. Baderan, R. Hutagalung, F.A. Ahmad, 2025.

Universitas Negeri Gorontalo, Gorontalo, Indonesia.

E-mail: aang@ung.ac.id*

TECTONIC GEOHISTORY OF THE GORONTALO REGION BASED ON FORAMINIFERA FOSSIL

Aang Panji Permana — PhD, Researcher and Associate Professor in Geological Engineering Study Program, Earth Science and Technology Department, Faculty of Mathematics and Natural Sciences, Universitas Negeri Gorontalo, Indonesia, E-mail: aang@ung.ac.id, <https://orcid.org/0000-0002-6865-3564>;

Dewi Wahyuni K. Baderan — Researcher and Professor in Environmental Science Doctoral Study Program, Postgraduate, Universitas Negeri Gorontalo, Indonesia, <https://orcid.org/0000-0003-3014-0832>;

Ronal Hutagalung — Researcher and Lecturer in Geological Engineering Study Program, Earth Science and Technology Department, Faculty of Mathematics and Natural Sciences, Universitas Negeri Gorontalo, Indonesia, <https://orcid.org/0000-0001-8142-8903>;

Fadhil Abdillah Ahmad — Researcher in Geological Engineering Study Program, Earth Science and Technology Department, Faculty of Mathematics and Natural Sciences, Universitas Negeri Gorontalo, Indonesia, <https://orcid.org/0009-0009-6728-3192>.

Abstract. The tectonic geohistory of the Gorontalo region, situated at the convergent boundary of the Pacific and Eurasian plates and influenced by microplate interactions, remains inadequately understood despite its high seismic vulnerability. This study reconstructs the Cenozoic tectonic evolution of Gorontalo using benthonic foraminifera fossils to address critical gaps in regional geological knowledge and support disaster mitigation efforts. Field stratigraphy was conducted in Ombulo Village, where limestone bed thicknesses (up to 25 m) and facies distributions were measured. Sixteen benthonic foraminifera species were identified using Olympus SZ61 microscopy, enabling detailed paleobathymetric reconstructions. Bathymetric corrections were applied using (van Hinte's) midpoint-range method, and local depth changes were compared to global sea-level data using (Miller et al) to distinguish between eustatic and tectonic influences. The results indicate a progressive deepening from middle shelf (178.4 m) to upper slope (500 m) environments between 6.4–5.8 Ma, corresponding to a ~321.5 m increase in depth that is not explained by global sea-level changes. The study concludes that tectonic subsidence is the dominant process shaping the Gorontalo basin, with foraminifera

distributions serving as reliable proxies for reconstructing paleodepths. These findings have practical applications in disaster mitigation (by providing baseline data for subsidence modeling), conservation planning (through the identification of geological protected areas), and energy exploration (by characterizing carbonate reservoirs relevant to hydrocarbon prospecting).

Keywords: Foraminifera Fossil, Geohistory, Gorontalo, Tectonics.

Acknowledgments.

This research was funded by LPPM UNG via PNBП-Universitas Negeri Gorontalo according to DIPA-UNG No. 653/UN47/HK.02/2024 through Riset Akselerasi Lektor Kepala under contract No. 822/UN47.D1/PT.01.03/2024.

© А.П. Пермана*, Д.В.К. Бадеран, Р. Хутагалунг, Ф.А. Ахмад, 2025.

Негери Горонтало университети, Горонтало, Индонезия.

E-mail: aang@ung.ac.id*

ФОРАМИНИФЕРА ҚАЗЫҒЫНЫҢ НЕГІЗІНДЕГІ ГОРОНТАЛО АЙМАҒЫНЫҢ ТЕКТНИКАЛЫҚ ГЕОТАРИХЫ

Аанг Панджи Пермана — PhD, зерттеуші және Геологиялық инженерияны зерттеу бағдарламасының доценті, Жер Туралы ғылымдар және технологиялар кафедрасы, Математика және жаратылыстану ғылымдары факультеті, Негери Горонтало Университеті, Индонезия, E-mail: aang@ung.ac.id, <https://orcid.org/0000-0002-6865-3564>;

Дьюи Вахюни К. Бадеран — зерттеуші және қоршаған ортаны қорғау ғылымдарының докторантурасын зерттеу бағдарламасының профессоры, аспирантура, Негери Горонтало университеттері, Индонезия, <https://orcid.org/0000-0003-3014-0832>;

Ронал Хутагалунг — Индонезиядағы Негери Горонтало университетінің Математика және жаратылыстану ғылымдары факультеті, Жер туралы ғылымдар және технологиялар кафедрасының Геологиялық инженерияны зерттеу бағдарламасының зерттеушісі және оқытушысы, <https://orcid.org/0000-0001-8142-8903>;

Фадхил Абдилла Ахмад — Геологиялық инженерияны зерттеу бағдарламасының ғылыми қызметкері, Жер туралы ғылымдар және технологиялар кафедрасы, Математика және жаратылыстану факультеті, Негери Горонтало университети, Индонезия, <https://orcid.org/0009-0009-6728-3192>.

Аннотация. Тынық мұхиты мен Еуразиялық плиталардың конвергентті шекарасында орналасқан және микропластиналық өзара әрекеттесулердің әсерінен Горонтало аймағының тектоникалық геотарихы оның жоғары сейсмикалық осалдығына қарамастан жеткілікті түрде зерттелмеген. Бұл зерттеу аймақтық геологиялық білімдегі маңызды олқылықтарды жою және апаттардың салдарын азайту жөніндегі күш-жігерді қолдау үшін бентондық фораминифера қазбаларын пайдалана отырып, Горонталоның кайнозойлық тектоникалық эволюциясын қалпына келтіреді. Омбуло ауылында дала стратиграфиясы жүргізілді, онда әктас қабатының қалыңдығы (25 м дейін) және фациялық таралу өлшенді. Толық палеобатиметриялық реконструкцияларды

жүргізуге мүмкіндік беретін Olympus SZ61 микроскопиясының көмегімен он алты бентондық фораминифера түрі анықталды. Батиметриялық түзетулер (ван Хинтенін) ортаңғы диапазон әдісі арқылы қолданылды, ал жергілікті тереңдік өзгерістері эвстатикалық және тектоникалық әсерлерді ажырату үшін (Миллер және т.б.) теңіз деңгейіндегі жаһандық деректермен салыстырылды. Нәтижелер теңіз деңгейінің жаһандық өзгерістерімен түсіндірілмейтін тереңдіктің ~321,5 м ұлғаюына сәйкес келетін орта қайраңнан (178,4 м) жоғарғы еңіске (500 м) дейін 6,4–5,8 млн аралығында біртіндеп тереңдеуді көрсетеді. Бұл аймақтағы ерте плиоцендегі теңіз деңгейінің көтерілуі эвстатикалық факторларға емес, тектоникалық шөгуге байланысты болды деген гипотезаны қолдайды. Зерттеу нәтижесінде тектоникалық шөгу Горонтало бассейнін қалыптастыратын басым процесс болып табылады, фораминифера таралулары палеотереңдіктерді қалпына келтіру үшін сенімді прокси ретінде қызмет етеді. Бұл нәтижелер апаттардың салдарын азайтуда (шөгуді модельдеу үшін бастапқы деректерді беру арқылы), консервацияны жоспарлауда (геологиялық қорғалатын аумақтарды анықтау арқылы) және энергетикалық барлауда (көмірсутектерді іздеуге қатысты карбонатты қабаттарды сипаттау арқылы) практикалық қолданбаларға ие.

Түйін сөздер: фораминифераның қазба қалдықтары, геотарих, Горонтало, тектоника

© А.П. Пермана*, Д.В.К. Бадеран, Р. Хутагалунг, Ф.А. Ахмад, 2025.

Университет Негери Горонтало, Горонтало, Индонезия.

E-mail: aang@ung.ac.id*

ТЕКТОНИЧЕСКАЯ ГЕОИСТОРИЯ РЕГИОНА ГОРОНТАЛЛО НА ОСНОВАНИИ ИСКОПАЕМЫХ ФОРАМИНИФЕР

Аанг Панджи Пермана — PhD, исследователь и доцент кафедры инженерно-геологических исследований факультета Наук о Земле факультета Математики и естественных наук университета Негери Горонтало, Индонезия, E-mail: aang@ung.ac.id, <https://orcid.org/0000-0002-6865-3564>;

Дьюни Вахьюни К. Бадеран — исследователь и профессор докторантуры по экологическим наукам, аспирант Университета Негери Горонтало, Индонезия, <https://orcid.org/0000-0003-3014-0832>;

Ронал Хутагалунг — исследователь и преподаватель программы инженерно-геологических исследований, факультет Наук о Земле и технологий, факультет Математики и естественных наук, Университет Негери Горонтало, Индонезия, <https://orcid.org/0000-0001-8142-8903>;

Фадхил Абдиллах Ахмад — исследователь программы инженерно-геологических исследований, факультет Наук о Земле и технологий, факультет математики и естественных наук, Университет Негери Горонтало, Индонезия, <https://orcid.org/0009-0009-6728-3192>.

Аннотация. Тектоническая геоистория региона Горонтало, расположенного на конвергентной границе Тихоокеанской и Евразийской плит и находящегося под влиянием микроплитных взаимодействий,

остается недостаточно изученной, несмотря на его высокую сейсмическую уязвимость. Это исследование реконструирует кайнозойскую тектоническую эволюцию Горонтало с использованием ископаемых бентосных фораминифер для устранения критических пробелов в региональных геологических знаниях и поддержки усилий по смягчению последствий стихийных бедствий. Полевая стратиграфия была проведена в деревне Омбуло, где были измерены толщина известняковых слоев (до 25 м) и распределение фаций. Шестнадцать видов бентосных фораминифер были идентифицированы с помощью микроскопии Olympus SZ61, что позволило провести подробные палеобатиметрические реконструкции. Батиметрические поправки были применены с использованием метода средней точки диапазона (van Hinte), а локальные изменения глубины были сопоставлены с глобальными данными об уровне моря с использованием (Miller et al) для различения эвстатических и тектонических влияний. Результаты указывают на постепенное углубление от среднего шельфа (178,4 м) до верхней части склона (500 м) в период 6,4–5,8 млн лет назад, что соответствует увеличению глубины примерно на 321,5 м, что не объясняется глобальными изменениями уровня моря. Исследование приходит к выводу, что тектоническое опускание является доминирующим процессом, формирующим бассейн Горонтало, а распределение фораминифер служит надежными прокси для реконструкции палеоглубин. Эти результаты имеют практическое применение в смягчении последствий стихийных бедствий (путем предоставления исходных данных для моделирования опускания), планировании охраны природы (путем определения геологических охраняемых территорий) и разведке энергоносителей (путем характеристики карбонатных резервуаров, имеющих отношение к разведке углеводородов).

Ключевые слова: ископаемые фораминиферы, геоистория, Горонтало, тектоника.

Introduction. Gorontalo is one of the areas prone to earthquake disasters; this condition is due to Gorontalo's tectonic position, which is influenced by the activities of the meeting of two large plates, namely the Pacific plate and the Eurasian plate. In addition, three microplates also contribute to the seismicity in Gorontalo, namely the Sulawesi, Banggai-Sula, and Sangihe sea plates. Evidence in the field is the uplifted limestone of the Gorontalo region, originally under the sea but now exposed above the land into hills. The presence of limestone at high elevations is evidence of a powerful tectonic influence (Jablonski, 2007; Pasari, et al., 2021; Efendi, et al., 2021; Handyarso, et al., 2023; Meidji, et al., 2023). The tectonic evolution of the Gorontalo area is a challenge in this study in providing new geological information. This new information on tectonic geohistory is related to the importance and originality of this research, which lies in the research methods and objectives to be achieved. The existence of rocks and foraminifera fossil species

as objects of geological study is beneficial in reconstructing tectonic geohistory. Unfortunately, research on tectonic geohistory analysis of the Gorontalo region has never been done. For this reason, research is needed on the tectonic geohistory that occurred in Gorontalo not only thousands of years ago but millions of years ago until now. One way to do this is to utilize the content of foraminifera fossil species in Gorontalo rocks. This reconstruction is very important to provide geological information for the government, researchers, academics, and the public. The main target is to develop a model of geological conservation area and geological natural disaster mitigation based on local wisdom in the future.

Foraminifera is a unicellular organism from the phylum Protozoa, class Sarcodina, and sub-class Rhizopod. They are characterized by the ability to form an outer shell. Some species may have a length of more than 100 μm or more but generally have a size of less than 1mm (Jones, 1969; Hanagata & Nobuhara, 2015).

Applying benthonic foraminifera fossils for paleo bathymetry is closely related to the distribution of characteristic species based on bathymetry or depth. This distribution is associated with the ability of benthonic foraminifera to live at specific depths so that environments with different depths will have different characteristic species (Jones, 1994; Herkat & Ladjal, 2013; Milker et al., 2014). The distribution of benthonic foraminifera fossils will be used as a determinant of paleo bathymetry using two classifications, namely the classification of Jones (1994) and Tipsword et al. (1966), which divide paleo bathymetry based on the depth of habitat of each benthonic foraminifera species. In fact, Tipsword et al. (1996) divided bathymetric zones for marine environments based on microfossil association data and planktonic/benthonic (P/B) ratio from the Gulf of Mexico. This classification combines the data associated with lithology, sedimentology, and tectonics. Based on this background, this research aims to analyze the Tectonic Geohistory of Gorontalo Region based on Benthonic Foraminifera Fossil.

Materials and Methods.

The research location is in Ombulo Village at coordinates $00^{\circ} 39' 7.08''$ East Longitude and $00^{\circ} 39' 7.08''$ North Latitude, precisely in the highland morphology. The research material is limestone outcrops scattered along the Gorontalo Outer Ring Road. The stratigraphic position of the study area can be seen on the geological map of the Ombulo Region, Gorontalo Regency (Figure 1).




Figure 1. Distribution of limestone outcrops (TQl) with other rocks on the geological map of the Ombulo area, Gorontalo Regency.

The material or object of research is limestone outcrops with a thickness of up to 25 meters. The types of research used in this study are qualitative and quantitative. Qualitative analysis in the form of field surveys in the form of stratigraphic measurements is known as the measured section (MS) method and sampling. Sampling for biostratigraphic analysis to be taken to the laboratory is a fine-sized limestone sample, namely mud. Quantitative analysis is based on microscopic observations from the samples taken in the field for biostratigraphic analysis (Kadar, 1986; Armstrong & Brasier, 2005). The MS method measures the details of stratigraphic sediment layers using Jacob’s stick, which has a length of 1.5 meters with sequential and systematic sampling. The 1.5-meter Jacob’s stick has an interval of 10 cm marked by a striking color difference between black and white. This MS method aims to obtain the stratigraphy of the study area based on the different characteristics of each facies (Compton, 1985; Permana, et al., 2019; 2021; 2024).

Biostratigraphic analysis was conducted at the Paleontology and Micropaleontology Laboratory of Universitas Negeri Gorontalo using an Olympus SZ61 binocular microscope connected to a computer to facilitate the observation and documentation of photos or images of very small (micro) foraminifera fossils.

This biostratigraphic analysis helps identify the type of benthonic foraminifera fossils so that the main purpose of ancient environments and ancient salinity content can be known (Jones, 1994; Nichlos, 2009; Permana, et al., 2020; 2021; Refalta, et al., 2022).

Paleo bathymetry data obtained from the analysis of benthonic small foraminifera must first be corrected to create a bathymetry change curve from the oldest to the youngest rock, where the connected point is generally the midpoint of the paleo bathymetry range (van Hinte, 1978) (Figure 2). The final analysis step is to make eustasy corrections by correlating regional eustasy data with age. After that, the main cause of the change in paleo bathymetry will be known, whether it is tectonic or eustasy.




Figure 2. Paleo bathymetry correction curve (Van Hinte., 1978)

Results.

Geological survey in the study area based on measured section (MS) results shows that the main constituent is limestone. Based on the facies analysis in the field, there is only one facies, namely coralline rudstone interbedded with mudstone. Full petrological description: the limestone is white, poorly sorted, open-fabric, matrix-supported grains with dominant grain abundance >2 mm. The bedding structure is thinning upward. The composition is mollusk fragments, coral fragments, quartz, and opaque minerals as fragments with a matrix of micrites. Based on the petrological description, the rock’s name is coralline rudstone, which refers to the classification (Embry & Klován, 1971; Permana et al., 2019a). The insertion

based on petrological analysis is fine brown limestone (grain size $<1/256$ mm), mud-supported, well-sorted, closed fabric with micrite and quartz composition. According to the results of the petrological analysis, the rock's name is mudstone, referring to the classification (Embry & Klován, 1971; Permana, et al., 2019a). The age of the rock based on the biozonation of planktonic foraminifera in the study area with LO *Globoquadrina dehiscens* bio datum (5.80 MA) is equivalent to Early Pliocene (Permana, et al., 2019b).

Micropaleontological observations in the study area using the measuring section (MS) method on ten samples sorted stratigraphically from oldest (bottom) to youngest (top), there are 16 species of benthonic foraminifera fossils. Sixteen species of benthonic foraminifera fossils, namely *Bolivinita quadrilatera* (Schwager, 1866), *Cornuspira foliacea* (Phillippi, 1844), *Fijiniopsis fijianensis* (Cushman & Edwards, 1937), *Gyrodinoides broeckhiana* (Karrer, 1878), *Gyrodinoides soldanii* (d'Orbigny, 1825), *Lenticulina denticulifera* (Cushman, 1913), *Mellonis Affinis* (Reuss, 1851), *Nautilus craticulatus* (Fichtel & Moll, 1798), *Nonion fabum* (Fichtel & Moll, 1798), *Pararotalia venusta* (Brady, 1884), *Praeglobobulimina ovata* (d'Orbigny, 1839), *Pyrgoella sphaera* (d'Orbigny, 1839), *Rhabdammina discreta* (Brady, 1881), *Saccorhiza ramosa* (Brady, 1879), *Spiroplectinella wrightii* (Silvestri, 1903), and *Uvigerina hollicki* (Thalmann, 1950).

Paleo bathymetry analysis on the MS Daenaa line on samples 3A, 3B, 3C, 3D, 3E, 3F, 3G, 3H, 3I and 3J. The paleo bathymetry zone on the MS Danaa line is divided into several zones. The paleo bathymetry zone for sample 3A is the middle shelf-upper slope (72.5-283.65 meters). The paleo bathymetry zone for sample 3B is the outer shelf-upper slope (183-283.65 meters). The paleo bathymetric zone for sample 3C is the upper slope (283.65-366 meters). The paleo bathymetry zone for sample 3D is the upper slope-lower slope (283.65-1830 meters). The paleo bathymetry zone for sample 3E is the upper slope-lower slope (283.65-1830 meters). The paleo bathymetry zone for sample 3F is the upper slope (283.65-466.65 meters). The paleo bathymetry zone for sample 3G is the upper slope (283.65-366 meters). The paleo bathymetry zone for sample 3G is the upper slope (283.65-366 meters). The paleo bathymetry zone for sample 3H is the upper slope (283.65-466.65 meters). The paleo bathymetric zone for sample 3I is middle shelf-abyssal (54.9-1830 meters). The paleo bathymetric zone for sample 3J is the upper slope (283.65-549 meters). The benthonic fossil content association can be seen in Table 1.

Table 1. Benthonic foraminifera distribution chart and paleo bathymetry of the study area

SAMPLE CODE	Benthonic Foraminifera											NUMBER OF SPECIMENS	Paleobathymetry							Paleobathymetry (Jones., 1994)				
	<i>Bolivinita quadrilobata</i> (Schwager,1866)	<i>Cornuspira foliacea</i> (Philippi,1844)	<i>Fijinionion fjiense</i> (Cushman & Edwards,1937)	<i>Gyrodonoides broeckiana</i> (Karver)	<i>Gyrodonoides soldanii</i> (d'Orbigny),1825)	<i>Lenticulina denticulifera</i> (Cushman,1913)	<i>Mellonis Affinis</i> (Reuss)	<i>Nautilus craticulatus</i> (Fichtel & Moll)	<i>Nonion fabum</i> (Fichtel & Moll,1798)	<i>Paratotalia venusta</i> (Brady)	<i>Praeglobobulimina ovata</i> (d'Orbigny,1839)		<i>Pyrgoella sphaera</i> (d'Orbigny,1839)	<i>Rhabdammina discreta</i> (Brady,1881)	<i>Saccorhiza ramosa</i> (Brady,1879)	<i>Spiroplectinella wrightii</i> (Silvestri,1903)	<i>Uvigerina hollicki</i> (Thalmann,1950)							
																		Shelf			Slope			
																		Inner	Middle		Outer	Upper		Middle
3J		12	2				1					1					15							283,65-549 m
3I												4					5							54.9-1830 m
3H		7				1						4			1		13							283,65-466.65 m
3G		2						1				12					15							283.65-366 m
3F		1										1					2							283,65-466,65 m
3E												2	1				3							283.65-1830 m
3D												1	1				2							283.65-1830 m
3C		2						2				2					6							283,65-366 m
3B				1			1	1		2		3					7							183-283.65 m
3A	2	1				1	2	2				2					8							72.5-283.65 m

Discussion.

Paleobathymetry data obtained from the analysis of benthonic small foraminifera must be corrected first because the paleo bathymetry results obtained are in the form of a range, while what is needed in the study of sea level change is paleo bathymetry in the form of absolute numbers. Therefore, correcting by making a paleo bathymetry change curve from the oldest to the youngest stratigraphic layer is necessary. The step taken is that the connected point is generally the midpoint of the paleo bathymetry range. The resulting curve is made as smooth as possible so that it will be corrected for some depths by not plotting at the center of the depth range.

After completing the benthonic foraminifera distribution chart, the paleo bathymetry correction curve should be made. The paleo bathymetry correction curve is created by smoothly connecting the depth points from each sample's paleo bathymetry range, then sorted stratigraphically from bottom to top (van Hinte., 1978). The paleo bathymetry correction curve aims to determine the pattern of sea level change to be compared to the global sea level change curve. The sea level will be known whether it drops or rises so that if it is related to global sea level changes, it is clear that the tectonic influence is very strong in the study area.

Paleobathymetry correction curve analysis based on plotting ten samples from

the oldest sample, 3A (bottom), to the youngest sample, 3J (top), shows a striking or significant sea level change in the study area during the Early Pliocene (6.4 MA - 5.8 MA). Changes in sea level tend to increase (getting deeper) because initially, the depth of the sea level was 178.425 meters and then changed drastically to be deeper to 500 meters. The difference in sea level change from the oldest to the youngest stratigraphic layer is $500 - 178.425 \text{ meters} = 321.575 \text{ meters}$ (Figure 3).




Figure 3. The paleo bathymetry correction curve of the study area shows a deepening sea level change from the oldest to the youngest stratigraphic layer.

The results of the paleo bathymetry curve analysis in the study area are the key to answering the mystery of the Early Pliocene period (6.4-5.8 MA) whether the sea level rise of 321.575 meters was due to global sea level rise or tectonic influences in the form of subsidence. The step that needs to be done is to make an eustasy correction. Eustasy correction is done by correlating regional eustasy data with age. After that, the leading cause of paleo bathymetry changes will be known, whether due to tectonic influence or eustasy.




Figure 4. The position of the sea level of the study area (Early Pliocene 6.4-5.8 MA) in the global sea level curve tends to be stable, with no changes in either rising or falling sea level, referring to the model Miller et al. (2005).

Analysis of the geohistory of the study area refers to changes in sea level that rose drastically due to tectonic subsidence because, according to the model Miller et al. (2005) (Figure 4) in the Early Pliocene (6.4-5.8 MA), no global water level changes occurred. Based on (Figure 4), it shows that in the Early Pliocene, sea level tends to be stable. For this reason, if there is an increase in water level in the research area based on the analysis of the paleo bathymetry curve, the tectonic influence in the form of subsidence is dominant because this subsidence causes a change in the depth of the rock deposition basin which was initially at a depth of about 178.425 meters to 500 meters.

Conclusion.

Referring to the results of research on the Tectonic Geohistory of the Gorontalo Region Based on Benthonic Foraminifera Fossils, the primary lithology of the research area is limestone, which contains 16 benthonic foraminifera fossil species. Those fossils indicate that paleo bathymetry ranges from the middle shelf to the abyssal, which facilitates making paleo bathymetry curves after correction. This curve becomes data to develop the eustasy curve, which shows that the decrease in sea level in the study area is due to tectonic influences, namely subsidence.

References

- Armstrong H.A., & Brasier M.D. (2005). Microfossils. 2nd edition, Blackwell Publishing, Oxford, United Kingdom, 304 pp.
- Compton R.R. (1985). Geology in the field. Wiley Press-New York. 416 p. (in Eng.)
- Efendi HA, Marliyani GI, Pramumijoyo S. (2021). Recent faulting along the Gorontalo fault based on seismicity data analysis and lineament mapping. E3S Web Conf. 2021;325(01013):1–4. <https://doi.org/10.1051/e3sconf/202132501013>. (in Eng.)
- Embry A.F., & Klován J. E. (1971). A late devonian reef tract on northeastern Banks Island, NWT. Bull. Can. Petroleum Geol. Vol. 19. – P. 730–781. (in Eng.)
- Hanagata S., & Nobuhara T. (2015). Illustrated guide to Pliocene foraminifera from Miyakojima, Ryukyu Island Arc, with comments on biostratigraphy. *Palaeontologia Electronica* 18.1.3A: 1-140. <https://doi.org/10.26879/44> (in Eng.)
- Handyarso A, Permána H, Hanafi M, Sendjaja P, Mukti MM. (2023). Extensional tectonic investigation based on gravity and magnetic data analysis in the Gulf of Tomini, Indonesia. Rud Geol Naft Zb. Vol.38(4):1–18. <https://doi.org/10.17794/rgn.2023.4.1:1-18> 5. (in Eng.)
- Herkat M., & Ladjal A. (2013). Paleo bathymetry of foraminiferal assemblages from the Pliocene of the Western Sahel (North-Algeria). *Palaeogeography, Palaeoclimatology, Palaeoecology*. 374. 144–163. [10.1016/j.palaeo.2013.01.013](https://doi.org/10.1016/j.palaeo.2013.01.013). (in Eng.)
- Jablonski D. (2007). Geology and Exploration Potential of The Gorontalo Basin, Central Indonesia– Eastern Extension of The North Makassar Basin. 10.29118/IPA.1482.07.G.083. (in Eng.)
- Jones D.K., (1969). Introduction to Microfossils; Hafner Publishing Company. New York, 406 p. (in Eng.)
- Jones R. W. (1994). The challenger foraminifera. Oxford, Oxford University Press, 149 p.(in Eng.)
- Kadar D., (1986). Neogene Planktonic Foraminiferal Biostratigraphy of The South Central Java Area, Indonesia. GRDC Special Publication, 5: 83 p. (in Eng.)
- Meidji I.U., Jatnika J., Samatowa L., Ramadani A. I. W. S., Ramdhan M., Efendi H. A., & Jayadi H. (2023). Analyzing Seismicity Pattern of the Gorontalo Region and Its Surroundings in the Form of Hypocenter Relocation Using Double Difference Method. *Jurnal Penelitian Pendidikan IPA*, 9(10), 8581–8587. <https://doi.org/10.29303/jppipa.v9i10.5400> (in Eng.)

Milker Y, Weinkauff MFG, Titschack J, Freiwald A, Krüger S, Jorissen FJ, & Schmiedl G. (2017). Testing the applicability of a benthic foraminiferal-based transfer function for the reconstruction of paleowater depth changes in Rhodes (Greece) during the early Pleistocene. *PLoS One*. Vol.12(11):e0188447. doi: 10.1371/journal.pone.0188447. PMID: 29166653; PMCID: PMC5699812. (in Eng.)

Miller K, G., Kominz A., Browning J V., Wright J D., Mountain G S., Katz M.E., Sugarman P.J., Cramer B.S., Christie-Blick N., & Pekar S.,F., (2005). The phanerozoic record of global sea-level change. *Review. Science* vol. 310. (in Eng.)

Nichols G. 2009. *Sedimentology and stratigraphy*. Blackwell Science Ltd., London, 335 p. (in Eng.)

Pasari, S., Simanjuntak AVH, Neha, Sharma Y. (2021). Nowcasting earthquakes in Sulawesi Island, Indonesia. *Geosci Lett*. Vol. 8(1):1–13. <https://doi.org/10.1186/s40562-021-00197-5> 6. (in Eng.)

Permana A.P., Pramumijoyo S., & Akmaluddin. (2019). Uplift Rate of Gorontalo Limestone (Indonesia) Based on Biostratigraphy Analysis. *News of the National Academy of Sciences of the Republic of Kazakhstan, Series of Geology and Technical Sciences*. Vol.6. No. 438. 6-11. <https://doi.org/10.32014/2019.2518-170X.150>. (in Eng.)

Permana A.P., Pramumijoyo S., and Akmaluddin. (2020). Paleo bathymetry Analysis of Limestone in Bongomeme Region Based on Content of Benthic Foraminifera Fossil, Gorontalo District, Indonesia. *Bulletin of the Iraq Natural History Museum*. Vol. 16. No. 1. 1-14. <https://doi.org/10.26842/binhm.7.2020.16.1.0001>. (in Eng.)

Permana A.P., Pramumijoyo S., Akmaluddin & Barianto, D.H. (2021). Planktonic Foraminiferal Biostratigraphy of the Limboto Limestone, Gorontalo Province, Indonesia. *Kuwait Journal of Science*. Vol. 48. No. 1. –P. 116-126. <https://doi.org/10.48129/kjs.v48i1.6916>. (in Eng.)

Permana A.P., Suaib A., Hutagalung R., & Eraku S.S. (2024). Analysis of the Relative Age of Limestone at Tanjung Kramat Region, Gorontalo City, Indonesia. *News of the National Academy of Sciences of the Republic of Kazakhstan, Series of Geology and Technical Sciences*. Vol.1. No. 463. – P. 190-200. <https://doi.org/10.32014/2024.2518-170X.375>. (in Eng.)

Refalta C. F., Yudha D.S., & Barianto D. H. (2022). Biostratigraphy and Paleobathymetry Microfossil Foraminera in the Sentolo Formation on the Jambon Line, Bantul Regency, Special Region of Yogyakarta Province. *Journal of Tropical Biodiversity and Biotechnology*. 7. 62239. 10.22146/jtbb.62239. (in Eng.)

Tippsword H.L., Setzer F.M., & Smith L.F., (1966). Interpretation of depositional environment in gulf coast petroleum exploration from paleoecology and related stratigraphy. *Gulf Coast Association of Geological Societies*, vol. 16: pp. 119-129, 1966(in Eng.)

Van Hinte J.E., (1978), *Geohistory Analysis – Application of Micropaleontology in Exploration Geology*. The American Association of Petroleum Geologist Bulletin, vol.62, No.2, - P.201-222. (in Eng.)

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

**[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)
<http://www.geolog-technical.kz/index.php/en/>
ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)**

Директор отдела издания научных журналов НАН РК *А. Ботанқызы*

Редакторы: *Д.С. Аленов, Ж.Ш. Әден*

Верстка на компьютере *Г.Д. Жадыранова*

Подписано в печать 15.06.2025.

Формат 70х90^{1/16}. Бумага офсетная. Печать – ризограф.
14,5 п.л. Заказ 3.