

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
әл-Фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
им. аль-Фараби

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-Farabi
Kazakh National University

SERIES PHYSICO-MATHEMATICAL

2 (336)

MARCH – APRIL 2021

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series physico-mathematical journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of chemistry and technologies in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of chemical sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Физикалық-математикалық сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Химия және технология сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді химиялық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия физико-математическая» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по химическим наукам для нашего сообщества.

Б а с р е д а к т о р
ф.-м.ғ.д., проф., ҚР ҰҒА академигі
Ғ.М. Мұтанов

Р е д а к ц и я а л қ а с ы:

Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Байгунчеков Ж.Ж. проф., академик (Қазақстан)
Quevedo Hernando проф. (Мексика)
Жүсіпов М.А. проф. (Қазақстан)
Ковалев А.М. проф., академик (Украина)
Калимолдаев М.Н. проф., академик (Қазақстан)
Михалевич А.А. проф., академик (Белорусь)
Мырзакулов Р. проф., академик (Қазақстан)
Рамазанов Т.С. проф., академик (Қазақстан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)
Уалиев З.Г. проф., чл.-корр. (Қазақстан)
Харин С.Н. проф., академик (Қазақстан)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан Республикасының Ақпарат және коммуникациялар министрлігінің Ақпарат комитетінде
14.02.2018 ж. берілген № **16906-Ж** мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Тақырыптық бағыты: *физика-математика ғылымдары және ақпараттық
технологиялар саласындағы басым ғылыми зерттеулерді
жариялау.*

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекен-жайы: 050010, Алматы қ., Шевченко көш., 28; 219 бөл.;
тел.: 272-13-19; 272-13-18

<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2021

Типографияның мекен-жайы: «Аруна» ЖК, Алматы қ., Мұратбаева көш., 75.

Главный редактор
д.ф.-м.н., проф. академик НАН РК
Г.М. Мутанов

Редакционная коллегия:

Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Байгунчечков Ж.Ж. проф., академик (Казахстан)
Quevedo Hernando проф. (Мексика)
Жусупов М.А. проф. (Казахстан)
Ковалев А.М. проф., академик (Украина)
Калимолдаев М.Н. проф., академик (Казахстан)
Михалевич А.А. проф., академик (Беларусь)
Мырзакулов Р. проф., академик (Казахстан)
Рамазанов Т.С. проф., академик (Казахстан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)
Уалиев З.Г. проф., чл.-корр. (Казахстан)
Харин С.Н. проф., академик (Казахстан)

«Известия НАН РК. Серия физика-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и коммуникаций Республики Казахстан № 16906-Ж, выданное 14.02.2018 г.

Тематическая направленность: *публикация приоритетных научных исследований в области физико-математических наук и информационных технологий.*

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219; тел.: 272-13-19; 272-13-18

<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2021

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

Editor in chief

doctor of physics and mathematics, professor, academician of NAS RK

G.M. Mutanov

Editorial board:

Asanova A.T. prof. (Kazakhstan)

Boshkayev K.A. PhD (Kazakhstan)

Baigunchekov Zh.Zh. prof., akademik (Kazakhstan)

Quevedo Hemando prof. (Mexico)

Zhusupov M.A. prof. (Kazakhstan)

Kovalev A.M. prof., academician (Ukraine)

Kalimoldaev M.N. prof., akademik (Kazakhstan)

Mikhalevich A.A. prof., academician (Belarus)

Myrzakulov R. prof., akademik (Kazakhstan)

Ramazanov T.S. prof., akademik (Kazakhstan)

Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.

Tiginyanu I. prof., academician (Moldova)

Ualiev Z.G. prof., chl.-korr. (Kazakhstan)

Kharin S.N. prof., academician (Kazakhstan)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Communications of the Republic of Kazakhstan No. 16906-Ж, issued on 14.02.2018.

Thematic scope: *publication of priority research in the field of physical and mathematical sciences and information technology.*

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, Almaty, 050010, tel. 272-13-19; 272-13-18

<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2021

Address of printing house: ST "Aruna", 75, Muratbayev str., Almaty.

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

Volume 2, Number 336 (2021), 6 – 14

<https://doi.org/10.32014/2021.2518-1726.15>

UDC 517.958; 517.968.7; 519.62/.64

MRNTI 27.29.17; 27.29.25; 27.33.19

Zh. M. Kadirbayeva^{1,2}, E. A. Bakirova^{1,3}, A. Sh. Dauletbayeva³, A. A. Kassymgali³

¹Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan;

²International Information Technology University, Almaty, Kazakhstan;

³Kazakh National Women's Teacher Training University, Almaty, Kazakhstan

E-mail: bakirova1974@mail.ru

AN ALGORITHM FOR SOLVING A BOUNDARY VALUE PROBLEM FOR ESSENTIALLY LOADED DIFFERENTIAL EQUATIONS

Abstract. A linear boundary value problem for essentially loaded differential equations is considered. Using the properties of essentially loaded differential we reduce the considering problem to a two-point boundary value problem for loaded differential equations. This problem is investigated by parameterization method. We offer algorithm for solving to boundary value problem for the system of loaded differential equations. This algorithm includes of the numerical solving of the Cauchy problems for system of the ordinary differential equations and solving of the linear system of algebraic equations. For numerical solving of the Cauchy problem we apply the Runge–Kutta method of 4th order. The proposed numerical implementation is illustrated by example.

Key words: essentially loaded differential equation, numerically approximate method, algorithm.

Funding: This research is funded by the Science Committee of the Ministry of Education and Science of the Republic of Kazakhstan (Grant No. AP08955489).

Introduction. The mathematical description of various dynamic control processes in which the future flow of processes depends not only on the present, but also is substantially determined by the history of the process, is carried out using ordinary differential equations with different types of memory, also called equations with aftereffect or loaded differential equations [1]. Loaded differential equations are used to solve problems of long-term prediction and control of the groundwater level and soil moisture [2-4]. Various problems for loaded differential equations and methods of finding their solutions considered in [1, 6-16].

In the present paper, a linear boundary value problem for essentially loaded differential equations is investigated. The significance is that the loading member of the equation appear in the form of derivative of solution at loaded point of the interval, i.e. the order of the loaded term is equal to the order of the differential part of the equation. Presence of derivative of solution in loaded point has a strong influence on the properties of equations.

Statement of problem.

We consider a linear boundary value problem for essentially loaded differential equations

$$\frac{dx}{dt} = A(t)x + \sum_{j=0}^m M_j(t)x(\theta_j) + K(t)\dot{x}(\theta_0) + f(t), t \in (0, T), \quad (1)$$

$$Bx(0) + Cx(T) = d, d \in R^n, x \in R^n, \quad (2)$$

where the $(n \times n)$ -matrices $A(t)$, $K(t)$, $M_j(t)$ ($j = 0, 1, \dots, m$), and n -vector-function $f(t)$ are continuous on $[0, T]$, B and C are constant $(n \times n)$ -matrices, d is constant n -vector, and $0 = \theta_0 < \theta_1 < \dots < \theta_m < \theta_{m+1} = T$, $\|x\| = \max_i |x_i|$.

Let $C([0, T], R^n)$ denote the space of continuous on $[0, T]$ functions $x(t)$ with norm $\|x\|_1 = \max_{t \in [0, T]} \|x(t)\|$.

A solution to problem (1), (2) is a continuously differentiable on $(0, T)$ function $x(t) \in C([0, T], R^n)$ satisfying the essentially loaded differential equations (1) and boundary condition (2).

By setting $t = \theta_0$ in equation (1), we get

$$[I - K(\theta_0)]\dot{x}(\theta_0) = A(\theta_0)x(\theta_0) + \sum_{j=0}^m M_j(\theta_0)x(\theta_j) + f(\theta_0). \quad (3)$$

Assume that the matrix $[I - K(\theta_0)]$ is invertible. We obtain

$$\dot{x}(\theta_0) = [I - K(\theta_0)]^{-1}\{A(\theta_0)x(\theta_0) + \sum_{j=0}^m M_j(\theta_0)x(\theta_j) + f(\theta_0)\}. \quad (4)$$

We consider the following linear boundary value problem for loaded differential equations

$$\frac{dx}{dt} = A(t)x + \sum_{i=0}^m D_i(t)x(\theta_i) + F(t), t \in (0, T), \quad (5)$$

$$Bx(0) + Cx(T) = d, d \in R^n, x \in R^n, \quad (6)$$

where

$$D_0(t) = M_0(t) + K(t)[I - K(\theta_0)]^{-1}\{A(\theta_0) + M_0(\theta_0)\}$$

$$D_j(t) = M_j(t) + K(t)[I - K(\theta_0)]^{-1}M_j(\theta_0), j = \overline{1, m},$$

$$F(t) = K(t)[I - K(\theta_0)]^{-1}f(\theta_0) + f(t).$$

Let us consider an example showing that loads influences significantly to the property of boundary value problem. Consider the following Cauchy problem for the loaded differential equation:

$$\frac{dx}{dt} = -\alpha x(0.4) + f(t), t \in [0, 1], \quad (7)$$

$$x(0) = 1. \quad (8)$$

Solving the problem (7), (8) we get

$$x(t) - 1 = -\alpha x(0.4)t + \int_0^t f(\tau) d\tau.$$

The value of $x(0.4)$ satisfies the following equation:

$$\left(1 + \frac{2}{5}\alpha\right)x(0.4) = 1 + \int_0^{0.4} f(\tau) d\tau. \quad (9)$$

But if we take $\alpha = -\frac{5}{2}$, $f(t) = 1$ then the equation (9) does not hold and the Cauchy problem (7), (8) is not solved. At the same time, the Cauchy problem for a linear system of ordinary differential equations (without loading) always has a unique solution.

On $[0, 1]$ we consider a periodic boundary value problem for an ordinary differential equation

$$\frac{dx}{dt} = t, x(0) = x(1), t \in [0, 1].$$

The General solution of the differential equation has the form: $x(t) = \frac{t^2}{2} + C$. Substituting the General solution in the boundary conditions for determining C , we obtain the relation: $C = C + \frac{1}{2}$. Since there is no such number C , the problem has no solution.

Now, adding the load at the point $t = 0.5$ to the right side of the differential equation we obtain the following periodic boundary value problem for a loaded differential equation

$$\frac{dx}{dt} = t + x(0.5), x(0) = x(1), t \in [0, 1],$$

and the solution of this problem has the form $x(t) = -\frac{3}{8} + \frac{t^2}{2} - \frac{t}{2}$.

Scheme of parametrization method.

We use the approach offered in [16-21] to solve the boundary value problem (5), (6). This approach based on the algorithms of the parameterization method and numerical methods for solving Cauchy problems.

Let us now investigate boundary value problem (5), (6) by the parametrization method. The interval $[0, T]$ is divided into subintervals by loading points:

$$[0, T] = \bigcup_{r=1}^{m+1} [\theta_{r-1}, \theta_r].$$

Introduce $C([0, T], \theta_m, R^{n(m+1)})$ as a space of systems of functions $x[t] = (x_1(t), x_2(t), \dots, x_{m+1}(t))$, where $x_r: [\theta_{r-1}, \theta_r] \rightarrow R^n$ are continuous on $[\theta_{r-1}, \theta_r]$ and have finite left-sided limits $\lim_{t \rightarrow \theta_r-0} x_r(t)$ for all $r = 1: (m+1)$, with norm $\|x[\cdot]\|_2 = \max_{r=1, m+1} \sup_{t \in [\theta_{r-1}, \theta_r]} \|x_r(t)\|$.

Let $x_r(t)$ be the restriction of the function $x(t)$ to the r -th interval $[\theta_{r-1}, \theta_r]$, i.e. $x_r(t) = x(t)$ for $t \in [\theta_{r-1}, \theta_r]$, $r = 1: (m+1)$. Then we reduce problem (5), (6) to the equivalent multipoint boundary value problem

$$\frac{dx_r}{dt} = A(t)x_r + \sum_{i=0}^m D_i(t)x_{i+1}(\theta_i) + F(t), t \in [\theta_{r-1}, \theta_r], r = 1: (m+1), \quad (10)$$

$$Bx_1(0) + C \lim_{t \rightarrow T-0} x_{m+1}(t) = d, \quad (11)$$

$$\lim_{t \rightarrow \theta_s-0} x_s(t) = x_{s+1}(\theta_s), s = 1: m, \quad (12)$$

where (12) are conditions for matching the solution at the interior points of partition.

The solution of problem (10) - (12) is a system of functions $x^*[t] = (x_1^*(t), x_2^*(t), \dots, x_{m+1}^*(t)) \in C([0, T], \theta_m, R^{n(m+1)})$, where the functions $x_r^*(t), r = \overline{1, m+1}$, are continuously differentiable on $[\theta_{r-1}, \theta_r]$, which satisfies system (10) and conditions (11), (12).

Problems (5), (6) and (10)-(12) are equivalent. If a system of functions $\tilde{x}[t] = (\tilde{x}_1(t), \tilde{x}_2(t), \dots, \tilde{x}_{m+1}(t)) \in C([0, T], \theta_m, R^{n(m+1)})$ is a solution of problem (10)-(12), then the function $\tilde{x}(t)$ defined by the equalities $\tilde{x}(t) = \tilde{x}_r(t), t \in [\theta_{r-1}, \theta_r], r = 1: (m+1)$, $\tilde{x}(T) = \lim_{t \rightarrow T-0} \tilde{x}_{m+1}(t)$ is a solution of the original problem (5), (6). Conversely, if $x(t)$ is a solution of problem (5), (6), then the system of functions $x[t] = (x_1(t), x_2(t), \dots, x_{m+1}(t))$, where $x_r(t) = x(t), t \in [\theta_{r-1}, \theta_r], r = 1: (m+1)$, and $\lim_{t \rightarrow T-0} x_{m+1}(t) = x(T)$, is a solution of problem (10)-(12).

Introducing the additional parameters $\lambda_r = x_r(\theta_{r-1}), r = 1: (m+1)$, and performing a replacement of the function $u_r(t) = x_r(t) - \lambda_r$ on each r -th interval $[\theta_{r-1}, \theta_r], r = 1: (m+1)$, we obtain the boundary value problem with parameters

$$\frac{du_r}{dt} = A(t)[u_r + \lambda_r] + \sum_{i=0}^m D_i(t)\lambda_{i+1} + F(t), \quad (13)$$

$$t \in [\theta_{r-1}, \theta_r], r = 1: (m+1),$$

$$u_r(\theta_{r-1}) = 0, r = 1: (m+1), \quad (14)$$

$$B\lambda_1 + C\lambda_{m+1} + C \lim_{t \rightarrow T-0} u_{m+1}(t) = d, \quad (15)$$

$$\lambda_s + \lim_{t \rightarrow \theta_s-0} u_s(t) = \lambda_{s+1}, s = 1: m. \quad (16)$$

A pair $(u^*[t], \lambda^*)$ with elements $u^*[t] = (u_1^*(t), u_2^*(t), \dots, u_{m+1}^*(t)) \in C([0, T], \theta_m, R^{n(m+1)})$, $\lambda^* = (\lambda_1^*, \lambda_2^*, \dots, \lambda_{m+1}^*) \in R^{n(m+1)}$ is said to be a solution to problem (13)-(16) if the functions $u_r^*(t), r = 1: (m+1)$, are continuously differentiable on $[\theta_{r-1}, \theta_r]$ and satisfy (13) and additional conditions (15), (16) with $\lambda_j = \lambda_j^*, j = 1: (m+1)$, and initial conditions (14).

Problems (5), (6) and (13)-(16) are equivalent. If the $x^*(t)$ is a solution of problem (5), (6), then the pair $(u^*[t], \lambda^*)$, where $u^*[t] = (x^*(t) - x^*(\theta_0), x^*(t) - x^*(\theta_1), \dots, x^*(t) - x^*(\theta_m))$, and $\lambda^* = (x^*(\theta_0), x^*(\theta_1), \dots, x^*(\theta_m))$, is a solution of problem (13)-(16). Conversely, if a pair $(\tilde{u}[t], \tilde{\lambda})$ with elements $\tilde{u}[t] = (\tilde{u}_1(t), \tilde{u}_2(t), \dots, \tilde{u}_{m+1}(t)) \in C([0, T], \theta_m, R^{n(m+1)})$, $\tilde{\lambda} = (\tilde{\lambda}_1, \tilde{\lambda}_2, \dots, \tilde{\lambda}_{m+1}) \in R^{n(m+1)}$, is a solution of (13)-(16), then the function $\tilde{x}(t)$ defined by the equalities $\tilde{x}(t) = \tilde{u}(t) + \tilde{\lambda}_r, t \in [\theta_{r-1}, \theta_r], r = 1: (m+1)$, will be the solution of the original problem (5), (6).

Using the fundamental matrix $X_r(t)$ of differential equation $\frac{dx}{dt} = A(t)x$ on $t \in [\theta_{r-1}, \theta_r], r = 1: (m+1)$, we reduce the Cauchy problem for the system of ordinary differential equations with

parameters (13), (14) to the equivalent system of integral equations

$$u_r(t) = X_r(t) \int_{\theta_{r-1}}^t X_r^{-1}(\tau)A(\tau)d\tau \lambda_r + X_r(t) \int_{\theta_{r-1}}^t X_r^{-1}(\tau) \sum_{i=0}^m D_i(\tau)\lambda_{i+1} d\tau + X_r(t) \int_{\theta_{r-1}}^t X_r^{-1}(\tau)F(\tau)d\tau, t \in [\theta_{r-1}, \theta_r], r = 1: (m + 1). \tag{17}$$

Substituting the corresponding right-hand sides of (17) into the conditions (15), (16), we obtain a system of linear algebraic equations with respect to the parameters $\lambda_r, r = 1: (m + 1)$

$$B\lambda_1 + C\lambda_{m+1} + CX_{m+1}(T) \int_{\theta_m}^T X_{m+1}^{-1}(\tau) \left\{ A(\tau)\lambda_{m+1} + \sum_{i=0}^m D_i(\tau)\lambda_{i+1} \right\} d\tau = d - CX_{m+1}(T) \int_{\theta_m}^T X_{m+1}^{-1}(\tau)F(\tau)d\tau, \tag{18}$$

$$\lambda_s + X_s(\theta_s) \int_{\theta_{s-1}}^{\theta_s} X_s^{-1}(\tau)A(\tau)d\tau \lambda_s + X_s(\theta_s) \int_{\theta_{s-1}}^{\theta_s} X_s^{-1}(\tau) \sum_{i=0}^m D_i(\tau)\lambda_{i+1} d\tau - \lambda_{s+1} = -X_s(\theta_s) \int_{\theta_{s-1}}^{\theta_s} X_s^{-1}(\tau)F(\tau)d\tau, s = 1: m. \tag{19}$$

We denote the matrix corresponding to the left side of the system of equations (18), (19) by $Q_*(\theta)$ and write the system in the form

$$Q_*(\theta)\lambda = F_*(\theta), \lambda \in R^{n(m+1)}, \tag{20}$$

where

$$F_*(\theta) = \left(d - CX_{m+1}(T) \int_{\theta_m}^T X_{m+1}^{-1}(\tau)F(\tau)d\tau, -X_1(\theta_1) \int_{\theta_0}^{\theta_1} X_1^{-1}(\tau)F(\tau)d\tau, \dots, -X_m(\theta_m) \int_{\theta_{m-1}}^{\theta_m} X_m^{-1}(\tau)F(\tau)d\tau \right)'$$

It is not difficult to establish that the solvability of the boundary value problem (5), (6) is equivalent to the solvability of the system (20). The solution of the system (20) is a vector $\lambda^* = (\lambda_1^*, \lambda_2^*, \dots, \lambda_{m+1}^*) \in R^{n(m+1)}$ consists of the values of the solutions of the original problem (5), (6) in the initial points of subintervals, i.e. $\lambda_r^* = x^*(\theta_{r-1}), r = 1: (m + 1)$.

Further we consider the Cauchy problems for ordinary differential equations on subintervals

$$\frac{dz}{dt} = A(t)z + P(t), z(\theta_{r-1}) = 0, t \in [\theta_{r-1}, \theta_r], r = 1: (m + 1), \tag{21}$$

where $P(t)$ is either $(n \times n)$ matrix, or n vector, both continuous on $[\theta_{r-1}, \theta_r], r = 1: (m + 1)$. Consequently, solution to problem (21) is a square matrix or a vector of dimension n . Denote by $a(P, t)$ the solution to the Cauchy problem (21). Obviously,

$$a(P, t) = X_r(t) \int_{\theta_{r-1}}^t X_r^{-1}(\tau)P(\tau)d\tau, t \in [\theta_{r-1}, \theta_r],$$

where $X_r(t)$ is a fundamental matrix of differential equation (21) on the r -th interval.

An algorithm for solving problem (1), (2).

We offer the following numerical implementation of algorithm based on the Runge–Kutta method of 4th order.

1. Suppose we have a partition: $0 = \theta_0 < \theta_1 < \dots < \theta_m < \theta_{m+1} = T$. Divide each r -th interval $[\theta_{r-1}, \theta_r], r = 1: (m + 1)$, into N_r parts with step $h_r = (\theta_r - \theta_{r-1})/N_r$. Assume on each interval $[\theta_{r-1}, \theta_r]$ the variable $\hat{\theta}$ takes its discrete values: $\hat{\theta} = \theta_{r-1}, \hat{\theta} = \theta_{r-1} + h_r, \dots, \hat{\theta} = \theta_{r-1} + (N_r -$

1) $h_r, \hat{\theta} = \theta_r$, and denote by $\{\theta_{r-1}, \theta_r\}$ the set of such points.

2. Solving the Cauchy problems for ordinary differential equations

$$\begin{aligned} \frac{dz}{dt} &= A(t)z + A(t), z(\theta_{r-1}) = 0, t \in [\theta_{r-1}, \theta_r], \\ \frac{dz}{dt} &= A(t)z + D_i(t), z(\theta_{r-1}) = 0, t \in [\theta_{r-1}, \theta_r], i = 0: m, \\ \frac{dz}{dt} &= A(t)z + F(t), z(\theta_{r-1}) = 0, t \in [\theta_{r-1}, \theta_r], r = 1: (m + 1), \end{aligned}$$

by using again the Runge–Kutta method of 4th order, we find the values of $(n \times n)$ matrices $a_r(A, \hat{\theta}), a_r(D_i, \hat{\theta}), i = 0: m$, and n vector $a_r(f, \hat{\theta})$ on $\{\theta_{r-1}, \theta_r\}, r = 1: (m + 1)$

3. Construct the system of linear algebraic equations with respect to parameters

$$Q_*^{\tilde{h}}(\theta)\lambda = -F_*^{\tilde{h}}(\theta), \lambda \in R^{n(m+1)}, \quad (22)$$

Solving the system (22), we find $\lambda^{\tilde{h}}$. As noted above, the elements of $\lambda^{\tilde{h}} = (\lambda_1^{\tilde{h}}, \lambda_2^{\tilde{h}}, \dots, \lambda_{m+1}^{\tilde{h}})$ are the values of approximate solution to problem (5), (6) in the starting points of subintervals: $x^{\tilde{h}r}(\theta_{r-1}) = \lambda_r^{\tilde{h}}, r = 1: (m + 1)$.

4. To define the values of approximate solution at the remaining points of set $\{\theta_{r-1}, \theta_r\}$, we solve the Cauchy problems

$$\begin{aligned} \frac{dx}{dt} &= A(t)x + \sum_{i=0}^m D_i(t)\lambda_{r+1}^{\tilde{h}} + F(t), \\ x(\theta_{r-1}) &= \lambda_r^{\tilde{h}}, t \in [\theta_{r-1}, \theta_r], r = 1: (m + 1). \end{aligned}$$

And the solutions to Cauchy problems are found by the Runge–Kutta method of 4th order. Thus, the algorithm allows us to find the numerical solution to the problem (5), (6).

We can see that the solution of boundary value problem (5), (6) also is the solution of boundary value problem (1), (2), when the matrix $[I - K(\theta_0)]$ is invertible.

To illustrate the proposed approach for the numerical solving linear two-point boundary value problem for essentially loaded differential equations (1), (2) on the basis of parameterization method, let us consider the following example.

Example. We consider a linear boundary value problem for essentially loaded differential equations

$$\frac{dx}{dt} = A(t)x + \sum_{i=0}^3 M_i(t)x(\theta_i) + K(t)\dot{x}(\theta_0) + f(t), t \in (0,1), \quad (23)$$

$$Bx(0) + Cx(T) = d, d \in R^2, x \in R^2. \quad (24)$$

Here

$$\theta_0 = 0, \theta_1 = \frac{1}{4}, \theta_2 = \frac{1}{2}, \theta_3 = \frac{3}{4}, \theta_4 = T = 1, A(t) = \begin{pmatrix} t+2 & t^3 \\ t^2 & t-1 \end{pmatrix},$$

$$M_0(t) = \begin{pmatrix} 6 & t \\ t^2 & 4t \end{pmatrix}, M_1(t) = \begin{pmatrix} 3t & 5 \\ t & t^2 \end{pmatrix}, M_2(t) = \begin{pmatrix} t^2 & t+3 \\ 0 & 11 \end{pmatrix},$$

$$M_3(t) = \begin{pmatrix} 9 & t-3 \\ t^2+5 & t^3 \end{pmatrix}, K(t) = \begin{pmatrix} t^2+3 & 5t \\ 2 & t-3 \end{pmatrix}, B = \begin{pmatrix} 4 & -3 \\ -5 & -1 \end{pmatrix},$$

$$C = \begin{pmatrix} 9 & 8 \\ -9 & 3 \end{pmatrix}, d = \begin{pmatrix} 18 \\ -43 \end{pmatrix}, f(t) = \begin{pmatrix} -t^5 + t^4 - \frac{37t^2}{8} + \frac{601t}{64} - \frac{1179}{64} \\ -t^5 - \frac{17t^3}{16} + \frac{177t^2}{64} + \frac{719t}{64} + \frac{57}{64} \end{pmatrix}.$$

We find $\dot{x}(0)$ from (23):

$$[I - K(0)]\dot{x}(0) = A(0)x(0) + \sum_{i=0}^3 M_i(0)x(\theta_i) + f(0).$$

The matrix $[I - K(0)]$ is invertible. Then

$$\dot{x}(0) = \begin{pmatrix} -0.5 & 0 \\ -0.25 & 0.25 \end{pmatrix} \left\{ A(0)x(0) + \sum_{i=0}^m M_i(0)x(\theta_i) + f(0) \right\}.$$

We consider a linear boundary value problem for loaded differential equations

$$\frac{dx}{dt} = A(t)x + \sum_{i=0}^3 D_i(t)x(\theta_i) + F(t), t \in (0,1),$$

$$Bx(0) + Cx(T) = d, d \in R^2, x \in R^2,$$

where

$$D_0(t) = M_0(t) + K(t)[I - K(0)]^{-1}\{A(0) + M_0(0)\} = \begin{pmatrix} -4t^2 - 10t - 6 & -\frac{t}{4} \\ t^2 - 2t - 2 & \frac{15t}{4} + \frac{3}{4} \end{pmatrix}$$

$$D_1(t) = M_1(t) + K(t)[I - K(0)]^{-1}M_1(0) = \begin{pmatrix} 3t & -\frac{5}{4}(2t^2 + 5t + 2) \\ t & t^2 - \frac{5t}{4} - \frac{5}{4} \end{pmatrix}$$

$$D_2(t) = M_2(t) + K(t)[I - K(0)]^{-1}M_2(0) = \begin{pmatrix} t^2 & 11t - \frac{2t^2}{3} - \frac{3}{2} \\ 0 & 2t + 2 \end{pmatrix}$$

$$D_3(t) = M_3(t) + K(t)[I - K(0)]^{-1}M_3(0) = \begin{pmatrix} -\frac{9t^2}{2} - 5t - \frac{9}{2} & \frac{3t^2}{2} + \frac{19t}{4} + \frac{3}{2} \\ t^2 - t - 1 & t^3 + \frac{3t}{4} + \frac{3}{4} \end{pmatrix}$$

$$F(t) = K(t)[I - K(0)]^{-1}f(0) + f(t) = \begin{pmatrix} -t^5 + t^4 + \frac{587t^2}{128} + \frac{1073t}{32} + \frac{1179}{128} \\ -t^5 - \frac{17t^3}{16} + \frac{177t^2}{64} + \frac{257t}{16} + \frac{309}{64} \end{pmatrix}.$$

We use the numerical implementation of algorithm. Accuracy of solution depends on the accuracy of solving the Cauchy problem on subintervals and evaluating definite integrals. We provide the results of the numerical implementation of algorithm by partitioning the subintervals $[0, 0.25]$, $[0.25, 0.5]$, $[0.5, 0.75]$, $[0.75, 1]$ with step $h = 0.025$.

Solving the system of equations (22), we obtain the numerical values of the parameters

$$\lambda_1^{\tilde{h}} = \begin{pmatrix} -0.000000041 \\ -2.00000002 \end{pmatrix}, \lambda_2^{\tilde{h}} = \begin{pmatrix} 0.765624981 \\ -2.437500012 \end{pmatrix},$$

$$\lambda_3^{\tilde{h}} = \begin{pmatrix} 1.625000009 \\ -2.750000008 \end{pmatrix}, \lambda_4^{\tilde{h}} = \begin{pmatrix} 2.671875034 \\ -2.937500004 \end{pmatrix}.$$

We find the numerical solutions at the other points of the subintervals using Runge-Kutta method of the 4-th order to the following Cauchy problems

$$\frac{d\tilde{x}_r}{dt} = A(t)\tilde{x}_r + \sum_{i=0}^3 D_i(t)\lambda_{r+1}^{\tilde{h}} + F(t),$$

$$x(\theta_{r-1}) = \lambda_r^{\tilde{h}}, t \in [\theta_{r-1}, \theta_r], r = 1:4.$$

Exact solution of the problem (23), (24) is $x^*(t) = \begin{pmatrix} t^3 + 3t \\ t^2 - 3t - 2 \end{pmatrix}$.

The results of calculations of numerical solutions at the partition points are presented in the following table:

t	$\tilde{x}_1(t)$	$\tilde{x}_2(t)$	t	$\tilde{x}_1(t)$	$\tilde{x}_2(t)$
0	-0.000000041	-2.000000002	0.5	1.625000009	-2.750000008
0.025	0.075015586	-2.049375019	0.525	1.719703137	-2.774375007
0.05	0.150124963	-2.097500018	0.55	1.816375014	-2.797500007
0.075	0.22542184	-2.144375017	0.575	1.915109392	-2.819375007
0.1	0.300999967	-2.190000016	0.6	2.01600002	-2.840000006
0.125	0.376953094	-2.234375015	0.625	2.119140648	-2.859375006
0.15	0.453374971	-2.277500014	0.65	2.224625025	-2.877500006
0.175	0.530359349	-2.319375014	0.675	2.332546903	-2.894375005
0.2	0.607999976	-2.360000013	0.7	2.44300003	-2.910000005
0.225	0.686390603	-2.399375012	0.725	2.556078157	-2.924375005
0.25	0.765624981	-2.437500012	0.75	2.671875034	-2.937500004
0.275	0.845796858	-2.474375011	0.775	2.79048441	-2.949375004
0.3	0.926999986	-2.510000011	0.8	2.912000037	-2.960000004
0.325	1.009328114	-2.54437501	0.825	3.036515662	-2.969375004
0.35	1.092874991	-2.57750001	0.85	3.164125037	-2.977500004
0.375	1.177734369	-2.60937501	0.875	3.294921912	-2.984375005
0.4	1.263999997	-2.640000009	0.9	3.429000036	-2.990000005
0.425	1.351765625	-2.669375009	0.925	3.566453159	-2.994375006
0.45	1.441125003	-2.697500008	0.95	3.707375031	-2.997500007
0.475	1.532171881	-2.724375008	0.975	3.851859402	-2.999375009
0.5	1.625000009	-2.750000008	1	4.000000021	-3.000000011

For the difference of the corresponding values of the exact and constructed solutions of the problem the following estimate is true:

$$\max_{j=0,40} \|x^*(t_j) - \tilde{x}(t_j)\| < 0.00000004.$$

Conclusion. In this work, we propose a numerical implementation of parametrization method for finding solutions to linear two-point boundary value problem for system of essentially loaded differential equations. Using the parametrization method, we reduce the considered problem to the equivalent boundary value problem with parameters. The example illustrating the numerical algorithms of parametrization method are provided.

**Ж. М. Кадирбаева^{1,2}, Э. А. Бакирова^{1,3},
А. Ш. Даулетбаева³, А. А. Касымғали³**

¹Математика және математикалық модельдеу институты, Алматы қаласы, Қазақстан;

²Халықаралық ақпараттық технологиялар университеті, Алматы қаласы, Қазақстан;

³Қазақ Ұлттық қыздар педагогикалық университеті, Алматы қаласы, Қазақстан

ЕЛЕУЛІ ТҮРДЕ ЖҮКТЕЛГЕН ДИФФЕРЕНЦИАЛДЫҚ ТЕНДЕУЛЕР ҮШІН ШЕТТІК ЕСЕПТІ ШЕШУ АЛГОРИТМІ

Аннотация. Елеулі түрде жүктелген дифференциалдық теңдеулер үшін сызықтық шеттік есеп қарастырылады. Біз қарастырылып отырған есепті елеулі түрде жүктелген дифференциалдық теңдеу қасиеттерін пайдалана отырып жүктелген дифференциалдық теңдеу үшін екі нүктелі шеттік есептерге келтіреміз. Аталған есеп параметрлеу әдісі арқылы зерттеледі. Жүктелген дифференциалдық теңдеулер жүйесі үшін шеттік есептің шешімін табудың алгоритмі ұсынылады. Бұл алгоритм жәй дифференциалдық теңдеулер жүйесі үшін Коши есептерін сандық шешуді және алгебралық теңдеулер жүйесін шешуді қамтиды. Коши есептерін сандық түрде шешу үшін төртінші ретті Рунге-Куттаның әдісі қолданылады. Ұсынылып отырған сандық жүзеге асырылу мысалмен көрсетіледі.

Түйін сөздер: елеулі түрде жүктелген дифференциалдық теңдеу, сандық жуықталған әдіс, алгоритм.

Ж. М. Кадирбаева^{1,2}, Э. А. Бакирова^{1,3}, А. Ш. Даулетбаева³, А. А. Касымгали³

¹Институт математики и математического моделирования, Алматы, Казахстан;

²Международный Университет Информационных Технологий, Алматы, Казахстан;

³Казахский Национальный Женский Педагогический Университет, Алматы, Казахстан

АЛГОРИТМ РЕШЕНИЯ КРАЕВОЙ ЗАДАЧИ ДЛЯ СУЩЕСТВЕННО НАГРУЖЕННЫХ ДИФФЕРЕНЦИАЛЬНЫХ УРАВНЕНИЙ

Аннотация. Рассматривается линейная краевая задача для существенно нагруженных дифференциальных уравнений. Используя свойства существенно нагруженного дифференциального уравнения, мы сводим рассматриваемую задачу к двухточечной краевой задаче для нагруженных дифференциальных уравнений. Данная задача исследуется методом параметризации. Предлагается алгоритм нахождения решения краевой задачи для системы нагруженных дифференциальных уравнений. Данный алгоритм включает численное решение задач Коши для системы обыкновенных дифференциальных уравнений и решение линейной системы алгебраических уравнений. Для численного решения задачи Коши применяется метод Рунге-Кутты четвертого порядка. Предлагаемая численная реализация иллюстрируется примером.

Ключевые слова: существенно нагруженное дифференциальное уравнение, численно приближенный метод, алгоритм.

Information about authors:

Kadirbayeva Zhazira Muratbekovna, Institute of Mathematics and Mathematical Modeling, Institute of Information and Computational Technologies, Chief researcher, Candidate of Physical and Mathematical Sciences, International Information Technology University, assistant professor, zh.kadirbayeva@gmail.com; <https://orcid.org/0000-0001-8861-4100>;

Bakirova Elmira Ainabekovna, Institute of Mathematics and Mathematical Modeling, leading scientific researcher, Candidate of Physical and Mathematical Sciences, ass. professor, Kazakh National Women's Teacher Training University, professor, bakirova1974@mail.ru; <https://orcid.org/0000-0002-3820-5373>;

Dauletbayeva Arshat Sharapatkyzy, Kazakh National Women's Teacher Training University, master, arshat0107.d@mail.ru; <https://orcid.org/0000-0002-7498-196X>;

Kassymgali Akerke Aibynkyzy, Kazakh National Women's Teacher Training University, master, kasyngalieva99@mail.ru; <https://orcid.org/0000-0002-8945-8004>

REFERENCES

- [1] Nakhushiev A.M. (2012) Loaded equations and their applications [Nagruzhennyye uravneniya i ih primeneniye]. Moscow, Nauka (in Russian).
- [2] Nakhushiev A.M. (1995) Equations of mathematical biology [Uravneniya matematicheskoy biologii]. Moscow, Vyshaiya shkola (in Russian).
- [3] Nakhushiev A.M. (1982) An approximation method for solving boundary value problems for differential equations with applications to the dynamics of soil moisture and groundwater [Ob odnom priblizhennom metode resheniya kraevykh zadach dlja differencial'nykh uravnenij i ego prilozheniya k dinamike pochvennoj vlagi gruntovykh vod]. Differential equations. V. 18, No. 1. P. 72-81 (in Russian).
- [4] Nakhushiev A.M. (2006) Displacement problems for partial differential equations. [Zadachi so smeshheniem dlja uravnenij v chastnykh proizvodnykh]. Moscow, Nauka (in Russian).
- [5] Dzhenaliev M.T., Ramazanov M.I. (2010) Loaded equations as perturbations of differential equations [Nagruzhennyye uravneniya kak vozmushheniya differencial'nykh uravnenij]. Almaty: Gylym (in Russian).
- [6] Dzhenaliev M.T. (2001) Loaded equations with periodic boundary conditions. Differential equations. V. 37, No. 1. P. 51-57.
- [7] Abdullaev V.M., Aida-zade K.R. (2014) Numerical method of solution to loaded nonlocal boundary value problems for ordinary differential equations. Computational Mathematics and Mathematical Physics. V.54, No 7. P. 1096-1109.
- [8] Aida-zade K.R., Abdullaev V.M. (2014) On the numerical solution to loaded systems of ordinary differential equations with non-separated multipoint and integral conditions. Numerical Analysis and Applications. V. 7, No. 1. P. 1-14.
- [9] Aida-zade K.R., Abdullaev V.M. (2004) On a numerical solution of loaded differential equations. Journal of computational mathematics and mathematical physics. V. 44, No. 9. P. 1585-1595.
- [10] Alikhanov A.A., Berezkov A.M., Shkhanukhov-Lafishev M.Kh. (2005) Boundary value problems for certain classes of loaded differential equations and solving them by finite difference methods. Computational Mathematics and Mathematical Physics. V. 48. P. 1581-1590.

- [11] Bakirova E.A., Kadirbayeva Zh.M. **(2016)** On a solvability of linear multipoint boundary value problem for the loaded differential equations [O razreshimosti linejnoj mnogotochečnoj kraevoj zadachi dlja nagruzhennyh differencial'nyh uravnenij]. Izvestiya NAN PK. Ser. fiz.-mat. V. 5, No 309. P. 168-175. (in Russian).
- [12] Assanova A. T., Imanchiyev A. E., Kadirbayeva Zh. M. **(2018)** Numerical solution of systems of loaded ordinary differential equations with multipoint conditions. Computational Mathematics and Mathematical Physics. V.58, No.4. P.508–516.
- [13] Assanova A.T., Kadirbayeva Zh.M. **(2018)** On the numerical algorithms of parametrization method for solving a two-point boundary-value problem for impulsive systems of loaded differential equations. Computational and Applied Mathematics. V. 37, No 4. P. 4966-4976.
- [14] Bakirova E.A., Tleulesova A.B., Kadirbayeva Zh.M. **(2017)** On one algorithm for finding a solution to a two-point boundary value problem for loaded differential equations with impulse effect. Bulletin of the Karaganda university. Mathematics series. No 3(87). P. 43-50.
- [15] Kadirbayeva Zh.M. **(2017)** On the method for solving linear boundary-value problem for the system of loaded differential equations with multipoint integral condition. Mathematical Journal. V. 17, No 4(66). P. 50-61.
- [16] Assanova A.T., Bakirova E.A., Kadirbayeva Zh.M. **(2019)** Numerical implementation of solving a boundary value problem for a system of loaded differential equations with parameter // News of the NAS RK. Phys.-Math. Series. V3, No. 325. P. 77-84. <https://doi.org/10.32014/2019.2518-1726.27> ISSN 2518-1726 (Online), ISSN 1991-346X (Print)
- [17] Dzhumabayev D.S. **(1989)** Criteria for the unique solvability of a linear boundary-value problem for an ordinary differential equation [Priznaki odnoznachnoj razreshimosti linejnoj kraevoj zadachi dlja obyknovennogo differencial'nogo uravnenija]. U.S.S.R. Computational Mathematics and Mathematical Physics. V. 29, No. 1. P. 50-66. (in Russian).
- [18] Dzhumabaev D.S. **(2010)** A method for solving the linear boundary value problem for an integro-differential equation. Computational Mathematics and Mathematical Physics. V. 50, No. 7. P. 1150-1161.
- [19] Dzhumabaev D.S. **(2016)** On one approach to solve the linear boundary value problems for Fredholm integro-differential equations. Journal of Computational and Applied Mathematics. V. 294. P. 342-357.
- [20] Assanova A.T., Bakirova E.A., Kadirbayeva Zh.M. **(2020)** Numerical solution to a control problem for integro-differential equations // Computational mathematics and mathematical physics, 2020 Vol. 60, №2. P. 203-221.
- [21] Assanova A.T., Bakirova E.A., Kadirbayeva Zh.M., Uteshova R.E. **(2020)** A computational method for solving a problem with parameter for linear systems of integro-differential equations // Computational and Applied Mathematics, 2020. Vol. 39, №3. Article number: 248. <https://doi.org/10.1007/s40314-020-01298-1>

Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

www.nauka-nanrk.kz

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы: *М. С. Ахметова, Д. С. Аленов, Р.Ж. Мрзабаева*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 15.04.2021.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.

11,6 п.л. Тираж 300. Заказ 2.