

ISSN 1991-346X

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

**ФИЗИКА-МАТЕМАТИКА
СЕРИЯСЫ**



СЕРИЯ

ФИЗИКО-МАТЕМАТИЧЕСКАЯ



**PHYSICO-MATHEMATICAL
SERIES**

2 (300)

НАУРЫЗ – СӘУІР 2015 ж.

МАРТ – АПРЕЛЬ 2015 г.

MARCH – APRIL 2015

1963 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С ЯНВАРЯ 1963 ГОДА
PUBLISHED SINCE JANUARY 1963

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

АЛМАТЫ, ҚР ҰҒА
АЛМАТЫ, НАН РК
ALMATY, NAS RK

Б а с р е д а к т о р

ҚР ҰҒА академигі,

Мұтанов Г. М.

Р е д а к ц и я а л қ а с ы:

физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Әшімов А.А.**; техн. ғ. докторы, проф., ҚР ҰҒА академигі **Байғұнчекөв Ж.Ж.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Жұмаділдаев А.С.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Қалменов Т.Ш.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Мұқашев Б.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Өтелбаев М.О.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Тәкібаев Н.Ж.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Харин С.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Әбішев М.Е.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Жантаев Ж.Ш.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Қалимолдаев М.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Косов В.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Мұсабаев Т.А.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Ойнаров Р.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Рамазанов Т.С.** (бас редактордың орынбасары); физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Темірбеков Н.М.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Өмірбаев У.У.**

Р е д а к ц и я к е ñ е с і:

Украинаның ҰҒА академигі **И.Н. Вишневский** (Украина); Украинаның ҰҒА академигі **А.М. Ковалев** (Украина); Беларусь Республикасының ҰҒА академигі **А.А. Михалевич** (Беларусь); Әзірбайжан ҰҒА академигі **А. Пашаев** (Әзірбайжан); Молдова Республикасының ҰҒА академигі **И. Тигиняну** (Молдова); мед. ғ. докторы, проф. **Иозеф Банас** (Польша)

Главный редактор

академик НАН РК

Г. М. Мутанов

Редакционная коллегия:

доктор физ.-мат. наук, проф., академик НАН РК **А.А. Ашимов**; доктор техн. наук, проф., академик НАН РК **Ж.Ж. Байгунчеков**; доктор физ.-мат. наук, проф., академик НАН РК **А.С. Джумадильдаев**; доктор физ.-мат. наук, проф., академик НАН РК **Т.Ш. Кальменов**; доктор физ.-мат. наук, проф., академик НАН РК **Б.Н. Мукашев**; доктор физ.-мат. наук, проф., академик НАН РК **М.О. Отелбаев**; доктор физ.-мат. наук, проф., академик НАН РК **Н.Ж. Такибаев**; доктор физ.-мат. наук, проф., академик НАН РК **С.Н. Харин**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **М.Е. Абишев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Ж.Ш. Жантаев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **М.Н. Калимолдаев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **В.Н. Косов**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Т.А. Мусабаев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Р. Ойнаров**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Т.С. Рамазанов** (заместитель главного редактора); доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Н.М. Темирбеков**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **У.У. Умирбаев**

Редакционный совет:

академик НАН Украины **И.Н. Вишневский** (Украина); академик НАН Украины **А.М. Ковалев** (Украина); академик НАН Республики Беларусь **А.А. Михалевич** (Беларусь); академик НАН Азербайджанской Республики **А. Пашаев** (Азербайджан); академик НАН Республики Молдова **И. Тигиняну** (Молдова); д. мед. н., проф. **Иозеф Банас** (Польша)

«Известия НАН РК. Серия физико-математическая». ISSN 1991-346X

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,

www.nauka-nanrk.kz / physics-mathematics.kz

© Национальная академия наук Республики Казахстан, 2015

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

Editor in chief

G. M. Mutanov,
academician of NAS RK

Editorial board:

A.A. Ashimov, dr. phys-math. sc., prof., academician of NAS RK; **Zh.Zh. Baigunchekov**, dr. eng. sc., prof., academician of NAS RK; **A.S. Dzhumadildayev**, dr. phys-math. sc., prof., academician of NAS RK; **T.S. Kalmenov**, dr. phys-math. sc., prof., academician of NAS RK; **B.N. Mukhashev**, dr. phys-math. sc., prof., academician of NAS RK; **M.O. Otelbayev**, dr. phys-math. sc., prof., academician of NAS RK; **N.Zh. Takibayev**, dr. phys-math. sc., prof., academician of NAS RK; **S.N. Kharin**, dr. phys-math. sc., prof., academician of NAS RK; **M.Ye. Abishev**, dr. phys-math. sc., prof., corr. member of NAS RK; **Zh.Sh. Zhantayev**, dr. phys-math. sc., prof., corr. member of NAS RK; **M.N. Kalimoldayev**, dr. phys-math. sc., prof., corr. member of NAS RK; **V.N. Kosov**, dr. phys-math. sc., prof., corr. member of NAS RK; **T.A. Mussabayev**, dr. phys-math. sc., prof., corr. member of NAS RK; **R. Oinarov**, dr. phys-math. sc., prof., corr. member of NAS RK; **T.S. Ramazanov**, dr. phys-math. sc., prof., corr. member of NAS RK (deputy editor); **N.M. Temirbekov**, dr. phys-math. sc., prof., corr. member of NAS RK; **U.U. Umirbayev**, dr. phys-math. sc., prof., corr. member of NAS RK

Editorial staff:

I.N. Vishnievski, NAS Ukraine academician (Ukraine); **A.M. Kovalev**, NAS Ukraine academician (Ukraine); **A.A. Mikhalevich**, NAS Belarus academician (Belarus); **A. Pashayev**, NAS Azerbaijan academician (Azerbaijan); **I. Tighineanu**, NAS Moldova academician (Moldova); **Joseph Banas**, prof. (Poland).

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.
ISSN 1991-346X

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

www.nauka-nanrk.kz / physics-mathematics.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2015

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

Volume 2, Number 300 (2015), 89 – 93

UDC 517.956.224, 517.956.225, 517.956.227

ON THE SECOND CHARACTERISTIC NUMBER OF THE NEWTON POTENTIAL

D. Suragan

Institute of mathematics and mathematical modeling MES RK, Almaty, Kazakhstan, e-mail: suragan@list.ru

Keywords: characteristic number, Newton potential, minimizer domain.

Abstract. In this note we prove that the minimum of the second characteristic number of the Newton potential among bounded open sets of \mathbb{R}^d with given volume is achieved by the union of two identical balls. The Newton potential can be related to a nonlocal boundary value problem for the Laplacian, so we obtain results on the second eigenvalue of the nonlocal Laplacian as well.

1 Introduction. Let Ω be a bounded open domain in \mathbb{R}^d , $d \geq 3$. Consider the Newton potential operator $N: L_2(\Omega) \rightarrow L_2(\Omega)$

$$Nf := \int_{\Omega} \varepsilon_d(x-y)f(y)dy \quad (1)$$

where

$$\varepsilon_d(x-y) = \frac{1}{(d-2)\sigma_d|x-y|^{d-2}}, \quad d \geq 3, \quad (2)$$

and $\sigma_d = \frac{2\pi^{d/2}}{\Gamma(\frac{d}{2})}$ is the surface area of the unit sphere in \mathbb{R}^d .

Since ε_d is real and symmetric function N is self-adjoint operator. Therefore, all characteristic numbers are real. In addition, it is easy to check that the operator N is positive. This means all its eigenvalues are positive. The characteristic numbers N of may be enumerated in ascending order,

$$\mu_1 \leq \mu_2 \leq \dots$$

where μ_i is repeated in this series according to its multiplicity. We denote the corresponding eigenfunctions by u_i, u_i, \dots , so that for each characteristic number μ_i there is one and only one, corresponding eigenfunction u_i ,

$$u_i = \mu_i N u_i, \quad i = 1, 2, \dots$$

In a bounded domain Ω of the Euclidean space \mathbb{R}^d , it is very well known that the solution to the Laplacian equation

$$-\Delta u(x) = f(x), \quad x \in \Omega, \quad (3)$$

is given by the Newton potential formula

$$u(x) = \int_{\Omega} \varepsilon_d(x-y)f(y)dy, \quad x \in \Omega, \quad (4)$$

for suitable functions f supported in Ω . An interesting question having several important applications is what boundary conditions can be put on u on the (smooth) boundary $\partial\Omega$ so that equation (3) complemented by this boundary condition would have the solution in still given by the same formula (4), with the same kernel ε_d given by (2). It turns out that the answer to this question is the integral boundary condition [4]

$$-\frac{u(x)}{2} + \int_{\partial\Omega} \frac{\partial \varepsilon_d(x-y)}{\partial n_y} u(y) dy - \int_{\partial\Omega} \varepsilon_d(x-y) \frac{\partial u(y)}{\partial n_y} dy = 0, x \in \partial\Omega, \quad (5)$$

where $\frac{\partial}{\partial n_y}$ denotes the outer normal derivative at a point y on $\partial\Omega$. A converse question to the one above would be to determine the trace of the Newton potential (4) on the boundary surface $\partial\Omega$, and one can use the potential theory to show that it has to be given by (5).

In [4] by using the boundary condition (5) the eigenvalues and eigenfunctions of the Newton potential were explicitly calculated in the 2-disk and in the 3-ball. In general, the boundary value problem (3)-(5) has various interesting properties and applications (see, for example, Kac [2, 3] and Saito [8]). The boundary value problem (3)-(5) can also be generalized for higher degrees of the Laplacian, see [5]. In this paper we are interested in some spectral geometry questions of N .

Historically, for the first time in the scientific literature, in Rayleigh’s famous book “Theory of Sound” (first published in 1877), by using some explicit computation and physical interpretations, he stated that a circle minimizes (among all domains of the same area) the first eigenvalue of the Dirichlet Laplacian. The musical interpretation of this result could be: among all drums of given area, the circular drum is the one which produces the deepest bass note. The proof of this conjecture was obtained after some decades later, simultaneously (and independently) by G.Faber and E.Krahn. Nowadays, the Rayleigh-Faber-Krahn inequality has been expanded many other operators; see [6] for further references.

In Section 2 we prove the following Rayleigh-Faber-Krahn theorem for the Newton potential N , i.e. it is proved that a ball is minimizer of the first characteristic number of the Newton potential N among all domains of given volume in R^d .

In Section 3 we are interested in minimizing the second characteristic number of N among open sets of given volume. We show that the minimizer is no longer one ball, but two! The similar result for the Dirichlet Laplacian called Krahn-Szego theorem, that is, the minimum of the second eigenvalue of the Dirichlet Laplacian among bounded open sets of R^d with given volume is achieved by the union of two identical balls. See, for example, [1] for further references.

2 Rayleigh-Faber-Krahn theorem. In this section we prove the following analogy of the Rayleigh-Faber-Krahn theorem for the Newton potential N .

Theorem 1. *A ball Ω^* is minimizer of the first characteristic number of the Newton potential N among all domains of given volume, i.e.*

$$\mu_1(\Omega^*) \leq \mu_1(\Omega) \quad (6)$$

for an arbitrary bounded open domain $\Omega \subset R^d$ with $|\Omega| = |\Omega^*|$.

We will use this result later in the proof of Theorem 2.

Proof of Theorem 1. Slightly different statement of Lemma 1 is called Jentsch’s theorem in [9]. However, for completeness of this note we restate and give its proof below.

Lemma 1. *The smallest characteristic number μ_1 of N is simple; the corresponding eigenfunction u_1 is positive and any other eigenfunction $u_i, i \neq 1$ is sign changing in Ω .*

Proof. The eigenfunctions of N may be chosen to be real as its kernel is real. First let us prove that u_1 cannot change sign in the domain Ω , that is,

$$u_1(x)u_1(y) = |u_1(x)u_1(y)|, \quad x, y \in \Omega.$$

In fact, in the opposite case, by virtue of the continuity of the function $u_1(x)$, there would be neighborhoods $U(x_0, r) \subset \Omega$ and $U(y_0, r) \subset \Omega$ such that

$$|u_1(x)u_1(y)| > u_1(x)u_1(y), x \in U(x_0, r) \subset \Omega, y \in U(y_0, r) \subset \Omega.$$

And so, by virtue of

$$\int_{\Omega} \varepsilon_d(x-\xi)\varepsilon_d(\xi-y)d\xi > 0. \quad (7)$$

We obtain

$$\begin{aligned} \frac{(N^2|u_1|, |u_1|)}{\| |u_1| \|^2} &= \frac{1}{\| |u_1| \|^2} \int_{\Omega} \int_{\Omega} \int_{\Omega} \varepsilon_d(x-\xi)\varepsilon_d(\xi-y) d\xi |u_1(x)||u_1(y)| dx dy \\ &> \frac{1}{\| |u_1| \|^2} \int_{\Omega} \int_{\Omega} \int_{\Omega} \varepsilon_d(x-\xi)\varepsilon_d(\xi-y) d\xi u_1(x)u_1(y) dx dy = \frac{1}{\mu_1^2}. \end{aligned} \quad (8)$$

μ_1^2 is the smallest characteristic number of N^2 and u_1 is the eigenfunction corresponding to μ_1^2 , i.e.

$$u_1 = \mu_1^2 N^2 u_1.$$

Therefore, by the variational principle we have

$$\frac{1}{\mu_1^2} = \sup_{f \in L^2(\Omega)} \frac{(N^2 f, f)}{\|f\|^2}. \tag{9}$$

This means that the strong inequality (8) contradicts the variational principle (9).

Now we shall prove that the eigenfunction $u_1(x)$ cannot become zero in Ω and therefore can be chosen positive in Ω .

In fact, in the opposite case there will be a point $x_0 \in \Omega$ such that

$$u_1(x_0) = \mu_1^2 \int_{\Omega} \int_{\Omega} \varepsilon_d(x_0 - \xi) \varepsilon_d(\xi - y) d\xi u_1(y) dy = 0$$

from which, by virtue of the condition (7), the contradiction follows: $u_1(y) = 0, \forall y \in \Omega$.

Since u_1 is positive it follows that μ_1 is a simple. In fact, if there were an eigenfunction \tilde{u}_1 linearly independent of u_1 and corresponding to μ_1 , then for all real c linear combination $u_1 + c\tilde{u}_1$ also would be eigenfunction corresponding to μ_1 and therefore, by what has been proved, it could not become zero in Ω . As c is arbitrary, this is impossible.

Finally, we show that the other eigenfunction $u_i, i = 2, 3, \dots$, are sign changing in Ω . If $u_i \geq 0, i \neq 1$, or $u_i \leq 0, i \neq 1$ then

$$\int_{\Omega} u_1 u_i \neq 0$$

as $u_1(x) > 0$ in Ω . This contradicts the orthogonality of the eigenfunction family $\{u_i\}$ of N .

Lemma 1 is proved.

Let Ω be a bounded measurable set in R^d . Its symmetric rearrangement Ω^* is an open ball originated at 0 with a measure equal to the measure of Ω , i.e. $|\Omega^*| = |\Omega|$. Let u be a nonnegative measurable function in Ω , in the sense that all its positive level sets have finite measure,

$$Vol(\{x|u(x) > t\}) < \infty, (\forall t > 0).$$

In the definition of the symmetric-decreasing rearrangement of u can be used the layer-cake decomposition [7], which expresses a nonnegative function u in terms of its level sets as

$$u(x) = \int_0^{\infty} X\{u(x) > t\} dt$$

where X is the characteristic function of the corresponding domain.

Definition 1. [7] Let u be a nonnegative measurable function in Ω . A function

$$u^*(x) = \int_0^{\infty} X\{u(x) > t\}^* dt$$

is called a symmetric-decreasing rearrangement of a nonnegative measurable function u .

By Lemma 1 the first characteristic number μ_1 of the operator N is positive and simple; the corresponding eigenfunction u_1 can be chosen positive in Ω . Recalling Riesz' inequality [7] and the fact that $\varepsilon_d(x - y)$ is a symmetric-decreasing function, i.e. ε_d and ε_d^* have the same formula, we obtain

$$\int_{\Omega} \int_{\Omega} u_1(y) \varepsilon_d(x - y) u_1(x) dy dx \leq \int_{\Omega^*} \int_{\Omega^*} u_1^*(y) \varepsilon_d(x - y) u_1^*(x) dy dx. \tag{10}$$

In addition, for each nonnegative function $u \in L^2(\Omega)$ we have

$$\|u\|_{L^2(\Omega)} = \|u^*\|_{L^2(\Omega^*)}. \tag{11}$$

Therefore, from (10), (11) and the variational principle for $\mu_1(\Omega^*)$, we get

$$\begin{aligned} \mu_1(\Omega) &= \frac{\int_{\Omega} |u_1(x)|^2 dx}{\int_{\Omega} \int_{\Omega} u_1(y) \vartheta(x-y) u_1(x) dy dx} \geq \frac{\int_{\Omega^*} |u_1^*(x)|^2 dx}{\int_{\Omega^*} \int_{\Omega^*} u_1^*(y) \vartheta(x-y) u_1^*(x) dy dx} \geq \\ &= \frac{\int_{\Omega^*} |\vartheta(x)|^2 dx}{\inf_{\vartheta \in L^2(\Omega^*)} \int_{\Omega^*} \int_{\Omega^*} \vartheta(y) \vartheta(x-y) \vartheta(x) dy dx} = \mu_1(\Omega^*). \end{aligned}$$

Theorem 1 is proved.

3Krahn-Szego theorem. In this section we are interested in minimizing the second characteristic number of the Newton potential N among open sets of given volume. As in case of the Dirichlet Laplacian, the minimizer is no longer one ball, but two!

Theorem 2. *The minimum of $\mu_2(\Omega)$ among bounded open sets of R^d with given volume is achieved by the union of two identical balls.*

Similar result for the Dirichlet Laplacian is called the Krahn-Szego theorem. See, for example, [1] for further references.

Proof of Theorem 2. Lemma 1 says that among eigenfunctions of N only the first eigenfunction is positive

$$u_1(x) > 0, \forall x \in \Omega.$$

Therefore,

$$u_2(x) > 0, \forall x \in \Omega^+ \subset \Omega, \Omega^+ \neq \{0\}.$$

$$u_2(x) < 0, \forall x \in \Omega^- \subset \Omega, \Omega^- \neq \{0\}.$$

We have

$$u_2(x) = \mu_2(\Omega) \int_{\Omega} \varepsilon_d(x-y)u_2(y)dy, x \in \Omega.$$

Taking

$$u_2^+(x) = \begin{cases} u_2(x) & \text{in } \Omega^+, \\ 0 & \text{otherwise,} \end{cases} \quad (12)$$

and

$$u_2^-(x) = \begin{cases} u_2(x) & \text{in } \Omega^-, \\ 0 & \text{otherwise,} \end{cases}$$

we obtain

$$u_2(x) = \mu_2(\Omega) \int_{\Omega^+} \varepsilon_d(x-y)u_2^+(y)dy + \mu_2(\Omega) \int_{\Omega^-} \varepsilon_d(x-y)u_2^-(y)dy, x \in \Omega.$$

Multiplying by $u_2^+(x)$ and integrating over Ω^+ we get

$$\begin{aligned} \int_{\Omega^+} |u_2^+(x)|^2 dx &= \mu_2(\Omega) \int_{\Omega^+} u_2^+(x) \int_{\Omega^+} \varepsilon_d(x-y)u_2^+(y)dydx + \\ &\mu_2(\Omega) \int_{\Omega^+} u_2^+(x) \int_{\Omega^-} \varepsilon_d(x-y)u_2^-(y)dydx, \quad x \in \Omega. \end{aligned}$$

The second term in the right hand side is negative as we know sign of all integrands. Therefore, one has

$$\int_{\Omega^+} |u_2^+(x)|^2 dx \leq \mu_2(\Omega) \int_{\Omega^+} u_2^+(x) \int_{\Omega^+} \varepsilon_d(x-y)u_2^+(y)dydx,$$

that is,

$$\frac{\int_{\Omega^+} |u_2^+(x)|^2 dx}{\int_{\Omega^+} u_2^+(x) \int_{\Omega^+} \varepsilon_d(x-y)u_2^+(y)dydx} \leq \mu_2(\Omega)$$

From here by using the variational principle one obtains

$$\begin{aligned} \mu_2(\Omega^+) &= \inf_{\vartheta \in L^2(\Omega^+)} \frac{\int_{\Omega^+} |\vartheta(x)|^2 dx}{\int_{\Omega^+} \vartheta(x) \int_{\Omega^+} \varepsilon_d(x-y)\vartheta(y)dydx} \\ &\leq \frac{\int_{\Omega^+} |u_2^+(x)|^2 dx}{\int_{\Omega^+} u_2^+(x) \int_{\Omega^+} \varepsilon_d(x-y)u_2^+(y)dydx} \leq \mu_2(\Omega). \end{aligned}$$

Similarly, we get

$$\mu_1(\Omega^-) \leq \mu_2(\Omega).$$

So we have

$$\mu_1(\Omega^+) \leq \mu_2(\Omega), \mu_1(\Omega^-) \leq \mu_2(\Omega). \quad (13)$$

We now introduce B^+ and B^- , balls of the same volume as Ω^+ and Ω^- , correspondingly. According to Theorem 1, we have

$$\mu_1(B^+) \leq \mu_1(\Omega^+), \quad \mu_1(B^-) \leq \mu_1(\Omega^-). \quad (14)$$

Let us introduce a new open set $\tilde{\Omega}$ defined as $\tilde{\Omega} = B^+ \cup B^-$. Since $\tilde{\Omega}$ is disconnected, we obtain its eigenvalues by gathering and reordering the eigenvalues of B^+ and B^- . Therefore,

$$\mu_2(\tilde{\Omega}) \leq \max(\mu_1(B^+), \mu_1(B^-)).$$

According to (13) and (14) we have

$$\mu_2(\tilde{\Omega}) \leq \max(\mu_1(B^+), \mu_1(B^-)) \leq \max(\mu_1(\Omega^+), \mu_1(\Omega^-)) \leq \mu_2(\Omega).$$

This shows that, in any case, the minimum of μ_2 is to be sought among the union of balls. But, if the two balls would have different radii, we would decrease the second eigenvalue by shrinking the largest one and dilating the smaller one (without changing the total volume). Therefore, the minimum is achieved by the union of two identical balls.

Theorem 2 is proved.

REFERENCES

- [1] Benguria R.D., Linde H., Loewe B. *Isoperimetric inequalities for eigenvalues of the Laplacian and the Schrodinger operator*, Bulletin of Mathematical Sciences, 2012, Vol. 2, Issue 1, 1-56.
- [2] Кас М. *On some connections between probability theory and differential and integral equations*, Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability, 1950, University of California Press, Berkeley and Los Angeles, 1951, 189-215.
- [3] Кас М. *Integration in function spaces and some of its applications*, Lezioni Fermiane. Accademia Nazionale dei Lincei, Pisa, 1980. 82 P.
- [4] Kalmenov T.Sh., Suragan D. *To spectral problems for the volume potential*, Doklady Mathematics, 2009, Vol. 80, 646-649.
- [5] Kalmenov T.Sh., Suragan D. *A boundary condition and spectral Problems for the Newton Potentials*, Operator Theory: Advances and Applications, 2011, Vol. 216, 187-210.
- [6] Henrot A. *Extremum problems for eigenvalues of elliptic operators*, Birkhauser, Basel, 2006.
- [7] Lieb E., Loss M. *Analysis*, 2nd ed., Amer. Math. Soc., 2001.
- [8] Saito N. *Data analysis and representation on a general domain using eigenfunctions of Laplacian*, Appl. Comput. Harmon. Anal. 2008, Vol. 25, 68-97.
- [9] Vladimirov V.S. *Equations of mathematical physics*, 2nd ed., Mir Publishers, Moscow 1983.

НЬЮТОН ПОТЕНЦИАЛЫНЫҢ ЕКІНШІ СИПАТТЫҚ САНЫ ЖАЙНДА

Д. Сұраған

ҚР БҒМ Математика және математикалық үлгілеу институты, Алматы, Қазақстан

Тірек сөздер: Лаплас-Бельтрами операторы, өткізгіш шекара, қисынды шектік есеп.

Аннотация. Жұмыста Ньютон потенциалының екінші сипаттық саны берілген көлемді R^d -дегі шенелген ашық жиындар арасында өзінің ең аз шамасын бір-келкі екі шардың біргеінде қабылдайтынын дәлелдейміз. Ньютон потенциалы Лапласианның локальды емес шекаралық есебіне қатасты болып табылатындықтан алынатын нәтижелер Лапласианның локальды емес шекаралық есебіне екінші меншікті мәнінде тиісті.

НЬЮТОН ПОТЕНЦИАЛЫНЫҢ ЕКІНШІ СИПАТТЫҚ САНЫ ЖАЙНДА

Д. Сураган

Институт математики и математического моделирования, МОН РК, Алматы, Казахстан

Тірек сөздер: Лаплас-Бельтрами операторы, өткізгіш шекара, қисынды шектік есеп.

Аннотация. Жұмыста Ньютон потенциалының екінші сипаттық саны берілген көлемді R^d -дегі шенелген ашық жиындар арасында өзінің ең аз шамасын бір-келкі екі шардың біргеінде қабылдайтынын дәлелдейміз. Ньютон потенциалы Лапласианның локальды емес шекаралық есебіне қатасты болып табылатындықтан алынатын нәтижелер Лапласианның локальды емес шекаралық есебіне екінші меншікті мәнінде тиісті.

Поступила 17.03.2015 г.

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

physics-mathematics.kz

Редактор *М. С. Ахметова*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 20.03.2015.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
10,5 п.л. Тираж 300. Заказ 2.