

ISSN 1991-346X

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

**ФИЗИКА-МАТЕМАТИКА
СЕРИЯСЫ**



СЕРИЯ

ФИЗИКО-МАТЕМАТИЧЕСКАЯ



**PHYSICO-MATHEMATICAL
SERIES**

3 (301)

МАМЫР – МАУСЫМ 2015 ж.

МАЙ – ИЮНЬ 2015 г.

MAY – JUNE 2015

1963 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН

ИЗДАЕТСЯ С ЯНВАРЯ 1963 ГОДА

PUBLISHED SINCE JANUARY 1963

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ

ВЫХОДИТ 6 РАЗ В ГОД

PUBLISHED 6 TIMES A YEAR

АЛМАТЫ, ҚР ҰҒА
АЛМАТЫ, НАН РК
ALMATY, NAS RK

Б а с р е д а к т о р

ҚР ҰҒА академигі,

Мұтанов Г. М.

Р е д а к ц и я а л қ а с ы:

физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Әшімов А.А.**; техн. ғ. докторы, проф., ҚР ҰҒА академигі **Байғұнчечков Ж.Ж.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Жұмаділдаев А.С.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Қалменов Т.Ш.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Мұқашев Б.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Өтелбаев М.О.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Тәкібаев Н.Ж.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Харин С.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Әбішев М.Е.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Жантаев Ж.Ш.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Қалимолдаев М.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Косов В.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Мұсабаев Т.А.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Ойнаров Р.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Рамазанов Т.С.** (бас редактордың орынбасары); физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Темірбеков Н.М.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Өмірбаев У.У.**

Р е д а к ц и я к ең е с і:

Украинаның ҰҒА академигі **И.Н. Вишневский** (Украина); Украинаның ҰҒА академигі **А.М. Ковалев** (Украина); Беларусь Республикасының ҰҒА академигі **А.А. Михалевич** (Беларусь); Әзірбайжан ҰҒА академигі **А. Пашаев** (Әзірбайжан); Молдова Республикасының ҰҒА академигі **И. Тигиняну** (Молдова); мед. ғ. докторы, проф. **Иозеф Банас** (Польша)

Главный редактор

академик НАН РК

Г. М. Мутанов

Редакционная коллегия:

доктор физ.-мат. наук, проф., академик НАН РК **А.А. Ашимов**; доктор техн. наук, проф., академик НАН РК **Ж.Ж. Байгунчеков**; доктор физ.-мат. наук, проф., академик НАН РК **А.С. Джумадильдаев**; доктор физ.-мат. наук, проф., академик НАН РК **Т.Ш. Кальменов**; доктор физ.-мат. наук, проф., академик НАН РК **Б.Н. Мукашев**; доктор физ.-мат. наук, проф., академик НАН РК **М.О. Отелбаев**; доктор физ.-мат. наук, проф., академик НАН РК **Н.Ж. Такибаев**; доктор физ.-мат. наук, проф., академик НАН РК **С.Н. Харин**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **М.Е. Абишев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Ж.Ш. Жантаев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **М.Н. Калимолдаев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **В.Н. Косов**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Т.А. Мусабаев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Р. Ойнаров**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Т.С. Рамазанов** (заместитель главного редактора); доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Н.М. Темирбеков**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **У.У. Умирбаев**

Редакционный совет:

академик НАН Украины **И.Н. Вишневский** (Украина); академик НАН Украины **А.М. Ковалев** (Украина); академик НАН Республики Беларусь **А.А. Михалевич** (Беларусь); академик НАН Азербайджанской Республики **А. Пашаев** (Азербайджан); академик НАН Республики Молдова **И. Тигиняну** (Молдова); д. мед. н., проф. **Иозеф Банас** (Польша)

«Известия НАН РК. Серия физико-математическая». ISSN 1991-346X

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,

www.nauka-nanrk.kz / physics-mathematics.kz

© Национальная академия наук Республики Казахстан, 2015

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

Editor in chief

G. M. Mutanov,
academician of NAS RK

Editorial board:

A.A. Ashimov, dr. phys-math. sc., prof., academician of NAS RK; **Zh.Zh. Baigunchekov**, dr. eng. sc., prof., academician of NAS RK; **A.S. Dzhumadildayev**, dr. phys-math. sc., prof., academician of NAS RK; **T.S. Kalmenov**, dr. phys-math. sc., prof., academician of NAS RK; **B.N. Mukhashev**, dr. phys-math. sc., prof., academician of NAS RK; **M.O. Otelbayev**, dr. phys-math. sc., prof., academician of NAS RK; **N.Zh. Takibayev**, dr. phys-math. sc., prof., academician of NAS RK; **S.N. Kharin**, dr. phys-math. sc., prof., academician of NAS RK; **M.Ye. Abishev**, dr. phys-math. sc., prof., corr. member of NAS RK; **Zh.Sh. Zhantayev**, dr. phys-math. sc., prof., corr. member of NAS RK; **M.N. Kalimoldayev**, dr. phys-math. sc., prof., corr. member of NAS RK; **V.N. Kosov**, dr. phys-math. sc., prof., corr. member of NAS RK; **T.A. Mussabayev**, dr. phys-math. sc., prof., corr. member of NAS RK; **R. Oinarov**, dr. phys-math. sc., prof., corr. member of NAS RK; **T.S. Ramazanov**, dr. phys-math. sc., prof., corr. member of NAS RK (deputy editor); **N.M. Temirbekov**, dr. phys-math. sc., prof., corr. member of NAS RK; **U.U. Umirbayev**, dr. phys-math. sc., prof., corr. member of NAS RK

Editorial staff:

I.N. Vishnievski, NAS Ukraine academician (Ukraine); **A.M. Kovalev**, NAS Ukraine academician (Ukraine); **A.A. Mikhalevich**, NAS Belarus academician (Belarus); **A. Pashayev**, NAS Azerbaijan academician (Azerbaijan); **I. Tighineanu**, NAS Moldova academician (Moldova); **Joseph Banas**, prof. (Poland).

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.
ISSN 1991-346X

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

www.nauka-nanrk.kz / physics-mathematics.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2015

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

Volume 3, Number 301 (2015), 62 – 65

UDC 510.532

MINIMAL ELEMENTS AND MINIMAL COVERS IN ROGERS SEMILATTICES

As. A. Issakhov

Al-Farabi Kazakh National University, Almaty, Kazakhstan.

E-mail: asylissakhov@mail.ru

Key words: minimal numbering, positive numbering, decidable numbering, minimal cover.

Abstract. This article is devoted to studying algebraic properties of Rogers semilattices of generalized computable families of total functions. Indeed, it is a continuation of attempts to find the elementary properties of the corresponding Rogers semilattices, which differ from properties of the classical Rogers semilattices for families of computable functions. Families of total A -computable functions, where $\emptyset^1 \leq_T A$, will be considered. The basic idea was taken from [1].

Some definitions and notations. For a family F of total functions a numbering $\alpha : \omega \rightarrow F$ is called computable if the binary function $\lambda x \lambda y \alpha(x)(y)$ is total recursive. Let $\alpha : \omega \rightarrow F$ and $\beta : \omega \rightarrow F$ be two numberings of the same set F . We say that the numbering α is reducible to the numbering β , if there is a computable function f such that $\alpha = \beta \circ f$, and we write this symbolically as: $\alpha \leq \beta$. If $\alpha \leq \beta$ and $\beta \leq \alpha$ then the numberings α and β are said to be equivalent, written as $\alpha \equiv \beta$. Denote by $\text{deg}(\alpha)$ the degree of α , that is, the set $\{\beta \mid \beta \equiv \alpha\}$ of numberings. The reducibility of numberings is a pre-order relation on the set of all computable numberings of a family F , which we denote by $\text{Com}(F)$, and it induces a partial order relation on a set of degrees of the numberings in $\text{Com}(F)$, which we also denote by \leq . The partially ordered set $\mathfrak{R}(F) = \langle \{\text{deg}(\alpha) \mid \alpha \in \text{Com}(F)\}, \leq \rangle$ is an upper semilattice, which we call the Rogers semilattice of the family F .

A numbering α of a set F is called minimal if, for any numbering β of F , $\beta \leq \alpha$ implies that $\alpha \leq \beta$. A computable one-to-one numbering is called a Friedberg numbering. The numerical equivalence θ_α of a numbering α is defined as follows: $\theta_\alpha = \{(x, y) \mid \alpha(x) = \alpha(y)\}$. An equivalence relation η is said to be decidable (positive) if η is computable (computably enumerable). For the further undefined notions we refer to [2, 3].

Let F be a family of total functions which are computable by an oracle A . A numbering $\alpha : \omega \rightarrow F$ is called A -computable if the binary function $\alpha(n)(x)$ is A -computable, [4]. A family F is called A -computable if it has an A -computable numbering. If A is a recursive set, then we are dealing with a family of computable functions and its classical computable numberings. The partially ordered set $\mathfrak{R}_A(F) = \langle \{\text{deg}(\alpha) \mid \alpha \in C_A(F)\}, \leq \rangle$, where $C_A(F)$ denotes the set of all A -computable numberings of the family F , is called the Rogers semilattice of the family F , [4].

It is known that the Rogers semilattice of every infinite Σ_{n+2}^0 -computable family contains an infinite number of minimal elements, and if a family $S \subseteq \Sigma_{n+2}^0$ has a Σ_{n+2}^0 -computable Friedberg numbering

then S has an infinite number of non-equivalent positive undecidable Σ_{n+2}^0 -computable numberings and an infinite number of non-equivalent Σ_{n+2}^0 -computable minimal non-positive numberings, [1]. Generalizing these results according to [4] and doing some simplifying in their proofs we have the next statements.

Theorem 1 [5, 6]. Let F be an infinite A -computable family of total functions, where $\emptyset' \leq_T A$. Then F has infinitely many pairwise nonequivalent A -computable Friedberg numberings.

This theorem is obtained after generalization theorem 1 from [5].

Theorem 2. Let F be an infinite A -computable family of total functions, where $\emptyset' \leq_T A$. Then F has infinitely many pairwise nonequivalent positive non-decidable A -computable numberings.

Proof. By theorem 1, suppose that α is an A -computable Friedberg numbering of the family F . Let M be a maximal set and $\overline{M} = \{\overline{m}_0 < \overline{m}_1 < \overline{m}_2 < \dots\}$. For all $k \in \omega$ we construct numberings β_k of the family F by the following way:

$$\beta_k(\overline{m}_i) = \alpha(i) \text{ for any } i \in \omega, \text{ and } \beta_k(x) = \alpha(k) \text{ for any } x \in M.$$

Then

$$y = \beta_k(n)(x) \Leftrightarrow (n \in M \ \& \ y = \alpha(k)(x)) \text{ or } (n \in \overline{M} \ \& \ \exists i(\overline{m}_i = n \ \& \ y = \alpha(i)(x))),$$

where the relations $n \in M$, $n \in \overline{M}$ and $\overline{m}_i = n$ are \emptyset' -computable. Therefore β_k is an A -computable numbering of the family F . Also

$$\beta_k(x) = \beta_k(y) \Leftrightarrow x = y \text{ or } (x \in M \cup \{p\} \ \& \ y \in M \cup \{p\}),$$

where $p \in \overline{M}$ is a β_k -index of the function $\alpha(k)$. So, obviously, θ_{β_k} is a c.e. equivalence. Since the set $\beta_k^{-1}(\alpha(k)) = M \cup \{p\}$ is not computable the numbering β_k is not decidable.

Now we will show that if $k \neq l$ then $\beta_k \not\leq \beta_l$. Assume $\beta_k(x) = \beta_l(f(x))$ for some computable function f . Since $\beta_l(x) \neq \alpha(k)$ for any $x \in M$, it follows that $f(M) \subseteq f(\beta_k^{-1}(\alpha(k))) \subseteq \overline{M}$. And note that the set $\overline{M} \setminus f(\beta_k^{-1}(\alpha(k)))$ contains β_l -indices of all functions in F distinct from $\alpha(k)$, therefore this set is infinite. So, since M is maximal set, the c.e. set $f(M)$ is finite. Moreover, by computability of the set $f^{-1}(f(M)) = \{y | f(y) \in f(M)\}$, the set M is computable, which contradicts with maximality of the set M .

Theorem 3. Let F be an infinite A -computable family of total functions, where $\emptyset' \leq_T A$. Then F has infinitely many pairwise nonequivalent minimal non-positive A -computable numberings.

Proof. By theorem 1, suppose that α is an A -computable Friedberg numbering of the family F . Let M be a maximal set and $\overline{M} = \{\overline{m}_0 < \overline{m}_1 < \overline{m}_2 < \dots\}$. By theorem 2, we know that if we define numbering β of the family F by the following way:

$$\beta(\overline{m}_i) = \alpha(i) \text{ for any } i \in \omega, \text{ and } \beta(x) = \alpha(p) \text{ for any } x \in M \text{ with fixed } p \in \omega,$$

then β is positive non-decidable A -computable numbering of F such that $\beta^{-1}(\alpha(p))$ is a non-computable c.e. set. For all $k \in \omega$ we construct numberings γ_k of the family F by the next way:

$$\gamma_k(\overline{m}_i) = \beta(i) \text{ for any } i \in \omega, \text{ and } \gamma_k(x) = \beta(k) \text{ for any } x \in M.$$

We will show that if $\beta(k) \neq \alpha(p)$ then γ_k is a minimal non-positive A -computable numbering of F . First, we prove that γ_k is a minimal numbering. Let γ be a numbering of F which is reducible to γ_k via some computable function f , where $\text{rng}(f) = W$. Since F is an infinite family and $\gamma_k(x) = \beta(k)$ for any $x \in M$, it follows that the set $W \cap \overline{M}$ is infinite. Then, by M is a maximal set,

$\overline{M} \setminus W$ is a finite set. Now we construct a computable function g reducing γ_k to γ . We enumerate the sets M and W simultaneously, and for every $x \in M \cup W$ if x appears in M first then $g(x)$ is equal to some fixed γ -index of the function $\beta(k)$ (for example, $g(x) = c = \mu_m(\mu_z(\hat{f}(z) = f(m)))$), where \hat{f} is computable function such that $\text{rng}(\hat{f}) = M$, otherwise $g(x) = \mu_y(f(y) = x)$. Finally if x belongs to the finite set $\overline{M} \setminus W$ then again $g(x)$ is equal to some fixed γ -index of the function $\beta(k)$, which completes the construction.

Now it remains to show that if $\beta(k) \neq \alpha(p)$ then γ_k is a non-positive A -computable numbering. $\gamma_k(x) = \alpha(p)$ for all $x \in \widehat{W} \subseteq \overline{M}$, since $\gamma_k(x) = \beta(k) \neq \alpha(p)$ for all $x \in M$, where $\widehat{W} = \gamma_k^{-1}(\alpha(p))$ is infinite set. If γ_k were positive numbering then the set \widehat{W} would be c.e. set, it follows that $\omega \setminus \widehat{W}$ is an infinite set since $M \subseteq \omega \setminus \widehat{W}$, and $\widehat{W} \setminus M$ is an infinite set since $\widehat{W} \subseteq \overline{M}$, which contradicts with the affirmation that M is maximal set.

Theorem 4 [7]. Let F be an A -computable family of total functions, where $\emptyset' \leq_T A$. If F contains at least two functions then F has no A -computable principal numbering.

This theorem is obtained after generalization theorem 3 from [5].

If for any non-principal A -computable numbering α of a family S we can find an A -computable numbering β of S such that $\text{deg}(\beta) > \text{deg}(\alpha)$, and for every degree $\text{deg}(\gamma)$, if $\text{deg}(\alpha) \leq \text{deg}(\gamma) \leq \text{deg}(\beta)$ then either $\text{deg}(\gamma) = \text{deg}(\alpha)$ or $\text{deg}(\gamma) = \text{deg}(\beta)$, then $\text{deg}(\beta)$ is called a minimal cover of $\text{deg}(\alpha)$.

It is known that if a family $S \subseteq \Sigma_{n+2}^0$ contains at least two sets C and D , and α be a Σ_{n+2}^0 -computable numbering of S , and if the set $\alpha^{-1}(C)$ is \emptyset' -computable, then $\text{deg}(\alpha)$ has a minimal cover, [1]. But we have the next statement for an A -computable family of total functions.

Theorem 5. Let F be an A -computable family of total functions and let α be an A -computable numbering of F , where $\emptyset' \leq_T A$. If F contains at least two functions then $\text{deg}(\alpha)$ has a minimal cover.

Proof. By theorem 4, we know that any numbering of F is not principal. Let M be a maximal set and $\overline{M} = \{\overline{m}_0 < \overline{m}_1 < \overline{m}_2 < \dots\}$. Then there exists an one-to-one computable function f such that $f(\omega) = M$. If we define $\beta(x) = \alpha(f^{-1}(x))$ for all $x \in M$ then $\alpha \leq \beta$ via f and β is an A -computable since $f^{-1}(x)$ is \emptyset' -computable set. Now we take $f_1, f_2 \in F$ and find a number $a \in \omega$ such that $f_1(a) \neq f_2(a)$. We complete construction β by defining values $\beta(x)$ for $x \in \overline{M}$. If $\varphi_e(\overline{m}_e) \downarrow$ and $\alpha(\varphi_e(\overline{m}_e))(a) = f_1(a)$ then we define $\beta(\overline{m}_e) = f_2$, otherwise $\beta(\overline{m}_e) = f_1$. It is clear that β is still an A -computable. We will show that $\beta \not\leq \alpha$. Suppose $\beta(x) = \alpha(\varphi_e(x))$, $x \in \omega$, for some computable function φ_e . Put $x = \overline{m}_e$. If $\alpha(\varphi_e(\overline{m}_e))(a) = f_1(a)$ then by construction $\beta(\overline{m}_e)(a) = f_2(a) \neq f_1(a) = \alpha(\varphi_e(\overline{m}_e))(a)$, contradiction. If $\alpha(\varphi_e(\overline{m}_e))(a) \neq f_1(a)$ then by construction $\beta(\overline{m}_e)(a) = f_1(a) \neq \alpha(\varphi_e(\overline{m}_e))(a)$, contradiction again. Therefore $\text{deg}(\alpha) < \text{deg}(\beta)$.

Now let γ be an A -computable numbering of F such that $\alpha \leq \gamma$ and $\gamma \leq \beta$. Suppose $\gamma \leq \beta$ via computable function h where $\text{rng}(h) = V$ then $\gamma = \beta h$, also we know that $\alpha = \beta f$ where $\text{rng}(f) = M$. Since $\beta f \leq \beta h \leq \beta$, it follows that we can assume $M \subseteq V \subseteq \omega$ (otherwise we can redefine the function h such that the condition is satisfied). Since the set M is maximal, either $\omega \setminus V$ or $V \setminus M$ is finite. In the former case (for obvious reasons) $\gamma \equiv \beta$, in the latter case $\gamma \equiv \alpha$.

REFERENCES

- [1] Badaev S.A., Goncharov S.S. *Rogers semilattices of families of arithmetic sets*, Algebra and logic, **2001**, vol. 40, N 5, 283-291 (in Eng.).
- [2] Ershov Yu.L. *Theory of numerations*, M.: Nauka, **1977**, 416 p. (in Russ.).
- [3] Goncharov S.S., Sorbi A. *Generalized computable numerations and nontrivial Rogers semilattices*, Algebra and logic, **1997**, vol. 36, N 6, 621-641 (in Eng.).
- [4] Badaev S.A., Goncharov S.S. *Generalized computable universal numberings*, Algebra and logic, **2014**, vol. 53, N 5, 355-364 (in Eng.).
- [5] Issakhov A. *Some computable numberings of the families of total functions in the arithmetical hierarchy*, Bulletin of Symbolic Logic, **2014**, vol. 20, N 2, 230-231 (in Eng.).
- [6] Issakhov As.A. *Friedberg numberings of the family of total functions in the arithmetical hierarchy*, News of the National Academy of Sciences of the Republic of Kazakhstan, series of phys. and math., **2014**, vol. 293, N 1, 8-11 (in Russ.).
- [7] Issakhov As.A. *Some properties of computable numberings of the families of total functions in the arithmetical hierarchy*, KazNU Bulletin, series of math., mech., inf., **2014**, vol. 81, N 2, 62-65 (in Russ.).

РОДЖЕРС ЖАРТЫТОРЛАРЫНДАҒЫ МИНИМАЛ ЭЛЕМЕНТТЕР ЖӘНЕ МИНИМАЛ ЖАБУЛАР

Ас. А. Исахов

Әл-Фараби атындағы Қазақ ұлттық университеті, Алматы, Қазақстан

Тірек сөздер: минимал нөмірлеу, позитив нөмірлеу, шешілімді нөмірлеу, минимал жабу.

Аннотация. Мақала барлық жерде анықталған функциялардың жалпыланған есептелімді үйірлерінің Роджерс жартыторларының алгебралық қасиеттерін зерттеуге арналған. Мұнда айтылған Роджерс жартыторларының, есептелімді функциялар үйірлерінің классикалық Роджерс жартыторларынан айырмашылығын көрсететін элементар қасиеттерін іздеу жалғасуда. $\emptyset' \leq_T A$ болатындай барлық жерде анықталған A -есептелімді функциялар үйірлері қарастырылады. Мақаланың негізгі идеялары [1]-ден алынған.

МИНИМАЛЬНЫЕ ЭЛЕМЕНТЫ И МИНИМАЛЬНЫЕ ПОКРЫТИЯ В ПОЛУРЕШЕТКАХ РОДЖЕРСА

Ас. А. Исахов

Казахский национальный университет им. аль-Фараби, Алматы, Казахстан

Ключевые слова: минимальная нумерация, позитивная нумерация, разрешимая нумерация, минимальное покрытие.

Аннотация. Статья посвящена изучению алгебраических свойств полурешеток Роджерса, обобщенно вычислимых семейств всюду определенных функций. На самом деле, это продолжение попыток найти элементарные свойства соответствующих полурешеток Роджерса, отличных от свойств классических полурешеток Роджерса для семейств вычислимых функций. Будут рассматриваться семейства всюду определенных A -вычислимых функций, где $\emptyset' \leq_T A$. Основные идеи взяты из [1].

Поступила 25.02.2015 г.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

physics-mathematics.kz

Редактор *М. С. Ахметова*

Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 9.06.2015.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.

15,7 п.л. Тираж 300. Заказ 3.