

ISSN 1991-346X

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

**ФИЗИКА-МАТЕМАТИКА
СЕРИЯСЫ**



СЕРИЯ

ФИЗИКО-МАТЕМАТИЧЕСКАЯ



**PHYSICO-MATHEMATICAL
SERIES**

4 (308)

ШІЛДЕ – ТАМЫЗ 2016 ж.

ИЮЛЬ – АВГУСТ 2016 г.

JULY – AUGUST 2016

1963 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С ЯНВАРЯ 1963 ГОДА
PUBLISHED SINCE JANUARY 1963

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

АЛМАТЫ, ҚР ҰҒА
АЛМАТЫ, НАН РК
ALMATY, NAS RK

Б а с р е д а к т о р

ҚР ҰҒА академигі,

Мұтанов Г. М.

Р е д а к ц и я а л қ а с ы:

физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Әшімов А.А.**; техн. ғ. докторы, проф., ҚР ҰҒА академигі **Байғұнчекөв Ж.Ж.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Жұмаділдаев А.С.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Қалменов Т.Ш.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Мұқашев Б.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Өтелбаев М.О.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Тәкібаев Н.Ж.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА академигі **Харин С.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Әбішев М.Е.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Жантаев Ж.Ш.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Қалимолдаев М.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Косов В.Н.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Мұсабаев Т.А.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Ойнаров Р.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Рамазанов Т.С.** (бас редактордың орынбасары); физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Темірбеков Н.М.**; физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Өмірбаев У.У.**

Р е д а к ц и я к ең е с і:

Украинаның ҰҒА академигі **И.Н. Вишневский** (Украина); Украинаның ҰҒА академигі **А.М. Ковалев** (Украина); Беларусь Республикасының ҰҒА академигі **А.А. Михалевич** (Беларусь); Әзірбайжан ҰҒА академигі **А. Пашаев** (Әзірбайжан); Молдова Республикасының ҰҒА академигі **И. Тигиняну** (Молдова); мед. ғ. докторы, проф. **Иозеф Банас** (Польша)

Главный редактор

академик НАН РК

Г. М. Мутанов

Редакционная коллегия:

доктор физ.-мат. наук, проф., академик НАН РК **А.А. Ашимов**; доктор техн. наук, проф., академик НАН РК **Ж.Ж. Байгунчеков**; доктор физ.-мат. наук, проф., академик НАН РК **А.С. Джумадильдаев**; доктор физ.-мат. наук, проф., академик НАН РК **Т.Ш. Кальменов**; доктор физ.-мат. наук, проф., академик НАН РК **Б.Н. Мукашев**; доктор физ.-мат. наук, проф., академик НАН РК **М.О. Отелбаев**; доктор физ.-мат. наук, проф., академик НАН РК **Н.Ж. Такибаев**; доктор физ.-мат. наук, проф., академик НАН РК **С.Н. Харин**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **М.Е. Абишев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Ж.Ш. Жантаев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **М.Н. Калимолдаев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **В.Н. Косов**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Т.А. Мусабаев**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Р. Ойнаров**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Т.С. Рамазанов** (заместитель главного редактора); доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Н.М. Темирбеков**; доктор физ.-мат. наук, проф., чл.-корр. НАН РК **У.У. Умирбаев**

Редакционный совет:

академик НАН Украины **И.Н. Вишневский** (Украина); академик НАН Украины **А.М. Ковалев** (Украина); академик НАН Республики Беларусь **А.А. Михалевич** (Беларусь); академик НАН Азербайджанской Республики **А. Пашаев** (Азербайджан); академик НАН Республики Молдова **И. Тигиняну** (Молдова); д. мед. н., проф. **Иозеф Банас** (Польша)

«Известия НАН РК. Серия физико-математическая». ISSN 1991-346X

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,

www.nauka-nanrk.kz / physics-mathematics.kz

© Национальная академия наук Республики Казахстан, 2016

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

Editor in chief

G. M. Mutanov,
academician of NAS RK

Editorial board:

A.A. Ashimov, dr. phys-math. sc., prof., academician of NAS RK; **Zh.Zh. Baigunchekov**, dr. eng. sc., prof., academician of NAS RK; **A.S. Dzhumadildayev**, dr. phys-math. sc., prof., academician of NAS RK; **T.S. Kalmenov**, dr. phys-math. sc., prof., academician of NAS RK; **B.N. Mukhashev**, dr. phys-math. sc., prof., academician of NAS RK; **M.O. Otelbayev**, dr. phys-math. sc., prof., academician of NAS RK; **N.Zh. Takibayev**, dr. phys-math. sc., prof., academician of NAS RK; **S.N. Kharin**, dr. phys-math. sc., prof., academician of NAS RK; **M.Ye. Abishev**, dr. phys-math. sc., prof., corr. member of NAS RK; **Zh.Sh. Zhantayev**, dr. phys-math. sc., prof., corr. member of NAS RK; **M.N. Kalimoldayev**, dr. phys-math. sc., prof., corr. member of NAS RK; **V.N. Kosov**, dr. phys-math. sc., prof., corr. member of NAS RK; **T.A. Mussabayev**, dr. phys-math. sc., prof., corr. member of NAS RK; **R. Oinarov**, dr. phys-math. sc., prof., corr. member of NAS RK; **T.S. Ramazanov**, dr. phys-math. sc., prof., corr. member of NAS RK (deputy editor); **N.M. Temirbekov**, dr. phys-math. sc., prof., corr. member of NAS RK; **U.U. Umirbayev**, dr. phys-math. sc., prof., corr. member of NAS RK

Editorial staff:

I.N. Vishnievski, NAS Ukraine academician (Ukraine); **A.M. Kovalev**, NAS Ukraine academician (Ukraine); **A.A. Mikhalevich**, NAS Belarus academician (Belarus); **A. Pashayev**, NAS Azerbaijan academician (Azerbaijan); **I. Tighineanu**, NAS Moldova academician (Moldova); **Joseph Banas**, prof. (Poland).

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.
ISSN 1991-346X

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

www.nauka-nanrk.kz / physics-mathematics.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2016

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

Volume 3, Number 307 (2016), 214 – 225

УДК 517.957.6

ТОЧНОЕ И ПРИБЛИЖЕННОЕ РЕШЕНИЕ ДВУХФАЗОВОЙ ОБРАТНОЙ ЗАДАЧИ СТЕФАНА

М.М. Сарсенгельдин, С. Касабек, Б.М. Сагидолла

Университет им. Сулеймана Демиреля, Каскелен, Казахстан

Ключевые слова: Метод интегральной функция ошибок, Метод интегрального теплового баланса, Двухфазовая обратная задача Стефана.

Аннотация: Основная идея является нахождение точных решений двухфазной обратной задачи Стефана для вырожденной области с движущейся границей. Отслеживание ответов этих вопросов будут организованы следующим образом. Для нахождения аналитического решения мы в основном следуем методом, предложенным С.Н. Харин применяя формулу Фаа Ди Бруно для интегральных функций ошибок. В продолжении раздела интегральные функции ошибки и ее свойств, необходимых для разработки новых методов.

Задача Стефана очень сложна для точного решения. Поэтому мы пытаемся найти приближенное решение путем присвоения температурных профилей с изменением дифференциального уравнения на уравнения теплового баланса, которые получаются путем интегрирования по X и t . Для нахождения приближенного решения мы в основном следуем методом, предложенным С.Н. Харин, применения метода интегрального теплового баланса.

EXACT AND APPROXIMATE SOLUTIONS OF TWO PHASE INVERSE STEFAN PROBLEM

M. Sarsengeldin, S. Kassabek, B. Sagidolla

Department of mathematics and natural sciences, Suleyman Demirel University,
Almaty, Qaskelen, Kazakhstan

E-mail: sarsengeldin.merey@sdu.edu.kz, kassabek@gmail.com, bizhigit.sagidolla@gmail.com

Keywords: Integral Error Functions, Integral Power Balance Method, Two Phase Inverse Stefan Problem.

Abstract. The main idea is finding the exact solution of two phase inverse Stefan problem for degenerate domain with moving boundary.

Tracking answers of these questions will be organized as following. For finding analytical solution we mainly follow the method proposed by S.N. Kharin applying Faa Di Bruno's formula for Integral Error Functions. In the continuation of this section Integral Error Functions and its properties necessary for elaboration of new methods are presented.

The Stefan problem is very complicated for exact solution. Therefore we try to find an approximate solution by profile assignment of the temperature with change of differential equation by heat balance equations, which are obtained by integration with respect to X and t . For finding approximate solution we mainly follow the method proposed by S.N. Kharin applying Integral Power Balance Method.

Problem formulation

In the following two sections we will deal to find the exact and approximate solutions of heat equations:

$$\frac{\partial u_1}{\partial t} = a_1^2 \frac{\partial^2 u_1}{\partial x^2}, \quad 0 < x < \alpha(t), \quad t > 0 \quad (1)$$

$$\frac{\partial u_2}{\partial t} = a_2^2 \frac{\partial^2 u_2}{\partial x^2}, \quad \alpha(t) < x < \infty, \quad t > 0 \quad (2)$$

Subjected to the following conditions:

$$u_1(0, 0) = 0 \quad (3)$$

$$u_2(x, 0) = f(x) \quad \alpha(t) < x < \infty \quad (4)$$

$$-\lambda_1 \left. \frac{\partial u_1}{\partial x} \right|_{x=0} = P(t), \quad t > 0 \quad (5)$$

$$u_1(\alpha(t), t) = u_2(\alpha(t), t) = u_m \quad (6)$$

The Stefan's condition:

$$-\lambda_1 \left. \frac{\partial u_1}{\partial x} \right|_{x=\alpha(t)} = -\lambda_2 \left. \frac{\partial u_2}{\partial x} \right|_{x=\alpha(t)} + L\gamma \frac{d(\alpha(t))}{dt} \quad (7)$$

$$u_2|_{x=\infty} = 0 \quad (8)$$

It is necessary to find temperature distribution $u_1(x, t)$ and $u_2(x, t)$ also it is required to reconstruct the boundary function $P(t)$ if the free boundary $\alpha(t)$ is given. Such problem is called two phase inverse Stefan Problem. The heat spread in the solid is negligible because of the physical properties of contact material. This condition is valid for refractory metals like wolfram. After getting the solutions we try to discuss given both solutions.

I. EXACT SOLUTION OF TWO PHASE INVERSE STEFAN PROBLEM

1.1 Introduction

The first analytical solution of two phase inverse Stefan problem, which describes the dynamics of soil freezing has been published by Lamé and Clayperon.

Despite the quite extensive list of problems in literature which lead to the necessity to solve Stefan type problems see: e.g., [1–9] and a long bibliography [9] on methods for solving these problems lead to additional difficulties which occur due to the degeneracy of domains. In some specific cases particularly for free moving boundaries it is possible to construct Heat potentials and a problem can be reduced to the system of integral equations [2], [3], however in the case of degeneracy, singularity in integral equations occur, and method of successive approximations is inapplicable in general. Moreover, the use of numerical methods is problematic when the number of parameters is great. Therefore, development of new analytical methods is very important especially for various applications because it enables one to analyze an interrelationship of different input parameters and their influence on the dynamics of investigating phenomena.

As for applications: a wide range of electric contact phenomena, in particular, the phenomena occurring at the interaction of electrical arc with electrode can be described in dynamic use of the presented method see e.g., [9] for very short arc duration (nanosecond diapason), when experimental

investigation is very difficult. In this study we will try to find solution of two Phase Inverse Stefan problem for degenerate domain with moving boundary.

1.2 Problem Solution

Let us consider this problem at the suggestion that $\alpha(t) = \sum_{n=1}^{\infty} \alpha_n t^{n/2}$ is given.

The unknown functions are $u_1(x, t), u_2(x, t), P(t)$ we represent in the form of Heat polynomials and Integral Error Functions:

$$u_1(x, t) = \sum_{n=0}^{\infty} \left[(2a_1 \sqrt{t})^{2n} A_{2n} i^{2n} \operatorname{erfc} \left(\frac{x}{2a_1 \sqrt{t}} \right) + (2a_1 \sqrt{t})^{2n+1} A_{2n+1} i^{2n+1} \operatorname{erfc} \left(-\frac{x}{2a_1 \sqrt{t}} \right) \right] \quad (9)$$

$$u_2(x, t) = \sum_{n=0}^{\infty} \left[(2a_2 \sqrt{t})^{2n} B_{2n} i^{2n} \operatorname{erfc} \left(\frac{x}{2a_2 \sqrt{t}} \right) + (2a_2 \sqrt{t})^{2n+1} B_{2n+1} i^{2n+1} \operatorname{erfc} \left(-\frac{x}{2a_2 \sqrt{t}} \right) \right] \quad (10)$$

$$P(t) = \sum_{n=0}^{\infty} P_n t^{\frac{n}{2}} \quad (11)$$

Where the coefficients $A_{2k}, A_{2k+1}, B_{2k}, B_{2k+1}, P(t)$ should be found satisfying the boundary conditions.

Let us satisfy first the initial condition (4). It is clear that:

$$u_2(x, 0) = \lim_{t \rightarrow 0} u_2(x, t) = \sum_{n=0}^{\infty} \left[\lim_{t \rightarrow 0} (2a_2 \sqrt{t})^{2n} B_{2n} i^{2n} \operatorname{erfc} \left(\frac{x}{2a_2 \sqrt{t}} \right) + \lim_{t \rightarrow 0} (2a_2 \sqrt{t})^{2n+1} B_{2n+1} i^{2n+1} \operatorname{erfc} \left(-\frac{x}{2a_2 \sqrt{t}} \right) \right] \quad (12)$$

by properties of Integral Error Functions:

$$\lim_{x \rightarrow \infty} \frac{i^n \operatorname{erfc}(x)}{x^n} = 0$$

by using L'Hospital Rule to find limit:

$$\lim_{x \rightarrow \infty} \frac{i^n \operatorname{erfc}(-x)}{x^n} = \lim_{x \rightarrow \infty} \frac{i^{n-1} \operatorname{erfc}(-x)}{nx^{n-1}} = \lim_{x \rightarrow \infty} \frac{i^{n-2} \operatorname{erfc}(-x)}{n(n-1)x^{n-2}} = \dots = \lim_{x \rightarrow \infty} \frac{\operatorname{erfc}(-x)}{n!} = \frac{2}{n!}$$

So therefore,

$$u_2(x, 0) = \sum_{n=0}^{\infty} B_{2n+1} \frac{2}{(2n+1)!} x^{2n+1} = f(x) \quad (13)$$

If we expand $f(x)$ into Taylor series:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(2n+1)}(0)}{(2n+1)!} x^{2n+1} \quad (14)$$

$$\sum_{n=0}^{\infty} B_{2n+1} \frac{2}{(2n+1)!} x^{2n+1} = \sum_{n=0}^{\infty} \frac{f^{(2n+1)}(0)}{(2n+1)!} x^{2n+1}$$

$$B_{2n+1} = \frac{1}{2} f^{(2n+1)}(0), \quad (n = 0, 1, 2, \dots) \quad (15)$$

From the condition (6) for $u_1(\alpha(t), t) = u_m$:

$$u_2(x, t) = \sum_{n=0}^{\infty} \left[(2a_2 \sqrt{t})^{2n} B_{2n} i^{2n} \operatorname{erfc} \left(\frac{x}{2a_2 \sqrt{t}} \right) + (2a_2 \sqrt{t})^{2n+1} B_{2n+1} i^{2n+1} \operatorname{erfc} \left(-\frac{x}{2a_2 \sqrt{t}} \right) \right] \Big|_{x=\alpha(t)} = u_m$$

The main problem is to find the remaining unknown coefficients we have, also if $\sqrt{t} = \tau$, then our expression will get:

$$\begin{aligned} \frac{\partial^{2k}}{\partial \tau^{2k}} \left(\sum_{n=0}^{\infty} \left[(2a_2 \tau)^{2n} B_{2n} i^{2n} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_2 \tau} \right) + (2a_2 \tau)^{2n+1} B_{2n+1} i^{2n+1} \operatorname{erfc} \left(-\frac{\alpha(\tau^2)}{2a_2 \tau} \right) \right] \right) &= \frac{\partial^{2k}}{\partial \tau^{2k}} (u_m) = 0 \\ \sum_{n=0}^{\infty} \sum_{l=0}^{2k} (2a_2)^{2n} B_{2n} \binom{2k}{l} \frac{(2n)!}{(2n-2k+l)!} \cdot \tau^{2n-2k+l} \cdot \frac{\partial^{2k}}{\partial \tau^{2k}} \left(i^{2n} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_2 \tau} \right) \right) &+ \\ + (2a_2)^{2n+1} B_{2n+1} \binom{2k}{l} \frac{(2n)!}{(2n+1-2k+l)!} \cdot \tau^{2n+1-2k+l} \cdot \frac{\partial^{2k}}{\partial \tau^{2k}} \left(i^{2n+1} \operatorname{erfc} \left(-\frac{\alpha(\tau^2)}{2a_2 \tau} \right) \right) &= 0 \end{aligned}$$

After taking both parts 2k-derivatives we can see following expression:

$$\begin{aligned} \sum_{n=0}^{\infty} (2a_2)^{2n} B_{2n} \frac{(2k)!}{(2k-2n)!} \frac{\partial^{2k-2n}}{\partial \tau^{2k-2n}} \left(i^{2n} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_2 \tau} \right) \right) \Big|_{\tau=0} &+ \\ + (2a_2)^{2n+1} B_{2n+1} \frac{(2k)!}{(2k-2n-1)!} \frac{\partial^{2k-2n-1}}{\partial \tau^{2k-2n-1}} \left(i^{2n+1} \operatorname{erfc} \left(-\frac{\alpha(\tau^2)}{2a_2 \tau} \right) \right) \Big|_{\tau=0} &= 0 \end{aligned}$$

For this purpose we use the Faa Di Bruno formula (Arbogast formula) for a derivative of a composite function:

$$\begin{aligned} \frac{\partial^{2k-2n}}{\partial \tau^{2k-2n}} \left(i^{2n} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_2 \tau} \right) \right) &= \sum_{j=0}^{2k-2n} i^{2n} \operatorname{erfc}^{(j)}(\delta) \beta_{2k-2n,j}(\delta', \delta'', \dots, \delta^{2k-2n-j}) \\ i^{2n} \operatorname{erfc}^{(j)}(\delta) &= \frac{d^j}{d\tau^j} \left(i^{2n} \operatorname{erfc}(\delta) \right) \Big|_{\tau=0} = (-1)^{j-2n-1} \frac{2}{\sqrt{\pi}} H_{j-2n-1}(\delta) \exp(-\delta^2) \end{aligned}$$

$$\begin{aligned} \frac{\partial^{2k-2n-1}}{\partial \tau^{2k-2n-1}} \left(i^{2n+1} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_2 \tau} \right) \right) &= \sum_{j=0}^{2k-2n-1} i^{2n+1} \operatorname{erfc}^{(j)}(\delta) \beta_{2k-2n-1,j}(\delta', \delta'', \dots, \delta^{2k-2n-1-j}) \\ i^{2n+1} \operatorname{erfc}^{(j)}(\delta) &= \frac{d^j}{d\tau^j} \left(i^{2n+1} \operatorname{erfc}(\delta) \right) \Big|_{\tau=0} = (-1)^{j-2n-2} \frac{2}{\sqrt{\pi}} H_{j-2n-2}(\delta) \exp(-\delta^2) \end{aligned}$$

Let us denote our boundaries in the following form to get rid of the singularity:

$$\begin{aligned} \gamma(t) = \alpha(t) - \alpha_0 &= \gamma_1 t^{1/2} + \gamma_2 t + \dots = \sum_{n=0}^{\infty} \gamma_{n+1} t^{(n+1)/2} \\ \alpha(\tau^2) &= \gamma_1 \tau + \gamma_2 \tau^2 + \dots + \gamma_{n+1} \tau^{n+1} = \sum_{n=0}^{\infty} \gamma_{n+1} \tau^{(n+1)} \end{aligned}$$

Then we can easily find our boundaries:

$$\delta|_{\tau=0} = \frac{\sum_{n=0}^{\infty} \gamma_{n+1} \tau^{n+1}}{2a_2 \tau} \Bigg|_{\tau=0} = \frac{\gamma_1}{2a_2}$$

$$\delta|_{\tau=0} = -\frac{\sum_{n=0}^{\infty} \gamma_{n+1} \tau^{n+1}}{2a_2 \tau} \Bigg|_{\tau=0} = -\frac{\gamma_1}{2a_2}$$

Finally, we get the following sum:

$$\sum_{n=0}^{\infty} B_{2n} \frac{(2a_2)^{2n} (2k)!}{(2k-2n)!} \sum_{j=0}^{2k-2n} (-1)^{j-2n-1} \frac{2}{\sqrt{\pi}} H_{j-2n-1} \left(\frac{\gamma_1}{2a_2} \right) \exp\left(-\frac{\gamma_1^2}{4a_2^2}\right) \beta_{2k-2n,j}(\delta', \delta'', \dots, \delta^{2k-2n-j}) +$$

$$+ \sum_{n=0}^{\infty} B_{2n+1} \frac{(2a_2)^{2n+1} (2k)!}{(2k-2n-1)!} \sum_{j=0}^{2k-2n-1} (-1)^{j-2n} \frac{2}{\sqrt{\pi}} H_{j-2n-2} \left(-\frac{\gamma_1}{2a_2} \right) \exp\left(-\frac{\gamma_1^2}{4a_2^2}\right) \beta_{2k-2n-1,j}(\delta', \delta'', \dots, \delta^{2k-2n-1-j}) = 0$$

Here let us denote the known coefficients in the following form:

$$K_{n,1} = \sum_{n=0}^{\infty} \frac{(2a_2)^{2n} (2k)!}{(2k-2n)!} \sum_{j=0}^{2k-2n} (-1)^{j-2n-1} \frac{2}{\sqrt{\pi}} H_{j-2n-1} \left(\frac{\gamma_1}{2a_2} \right) \exp\left(-\frac{\gamma_1^2}{4a_2^2}\right) \beta_{2k-2n,j}(\delta', \delta'', \dots, \delta^{2k-2n-j})$$

$$K_{n,2} = \sum_{n=0}^{\infty} \frac{(2a_2)^{2n+1} (2k)!}{(2k-2n-1)!} \sum_{j=0}^{2k-2n-1} (-1)^{j-2n} \frac{2}{\sqrt{\pi}} H_{j-2n-2} \left(-\frac{\gamma_1}{2a_2} \right) \exp\left(-\frac{\gamma_1^2}{4a_2^2}\right) \beta_{2k-2n-1,j}(\delta', \delta'', \dots, \delta^{2k-2n-1-j})$$

After we can easily find our unknown coefficient:

$$B_{2n} \cdot K_{n,1} = B_{2n+1} \cdot K_{n,2}$$

$$B_{2n} = \frac{B_{2n+1} \cdot K_{n,2}}{K_{n,1}} = \frac{f^{(2n+1)}(0) \cdot K_{n,2}}{2K_{n,1}} \tag{16}$$

From the condition (6) for $u_1(x, t)$ will be the same situation for $u_1(\alpha(t), t) = u_m$:

$$A_{2n} = \frac{A_{2n+1} \cdot K_{n,2}}{K_{n,1}} \tag{17}$$

By using the condition (7):

$$dt = 2\tau d\tau$$

$$\left(\frac{L\gamma}{2\tau} \frac{d\alpha(\tau)}{d\tau} \right) \Bigg|_{\tau=0}^{(k)} = \left(\frac{L\gamma}{2} \sum_{n=0}^{\infty} \frac{\alpha^{(n)}(0)}{(2n)!} \tau^{2n-1} \right) \Bigg|_{\tau=0}^{(k)}$$

It is easy to see that $k = 2n - 1$, therefore:

$$\frac{L\gamma}{2} \sum_{n=0}^{\infty} \frac{\alpha^{(n)}(0)}{(2n)!} (\tau^{2n-1}) \Bigg|_{\tau=0}^{(k)} = \frac{L\gamma}{2} \sum_{n=0}^{\infty} \frac{\alpha^{(n)}(0)}{2n}$$

If we will take $2n - 1$ derivatives in both parts by using Leibniz and Faa di Bruno formulas:

$$\begin{aligned} & \left(-\lambda_1 \frac{\partial u_1}{\partial x} \Big|_{x=\alpha(\tau^2)} \right) \Big|_{\tau=0}^{(2n-1)} = \left(-\lambda_2 \frac{\partial u_2}{\partial x} \Big|_{x=\alpha(\tau^2)} \right) \Big|_{\tau=0}^{(2n-1)} + \frac{L\gamma}{4} \sum_{n=0}^{\infty} \frac{\alpha^{(n)}(0)}{n} \\ & \left[\lambda_1 \sum_{n=0}^{\infty} (2a_1)^{2n-1} A_{2n} (2n-1)! \left(i^{2n-1} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_1\tau} \right) \right) \Big|_{\tau=0} - \right. \\ & \left. - \lambda_1 \sum_{n=0}^{\infty} (2a_1)^{2n} A_{2n+1} \cdot \tau \cdot (2n-1)! \left(i^{2n} \operatorname{erfc} \left(-\frac{\alpha(\tau^2)}{2a_1\tau} \right) \right) \Big|_{\tau=0} \right] = \\ & \lambda_1 \sum_{n=0}^{\infty} (2a_1)^{2n-1} A_{2n} (2n-1)! \left(i^{2n-1} \operatorname{erfc} \left(\frac{\gamma_1 + \gamma_2\tau + \dots}{2a_1} \right) \right) \Big|_{\tau=0} = A_{2n} \sum_{n=0}^{\infty} \lambda_1 (2a_1)^{2n-1} (2n-1)! \left(i^{2n-1} \operatorname{erfc} \left(\frac{\gamma_1}{2a_1} \right) \right) \\ & \sum_{n=0}^{\infty} A_{2n} C_{n,1} = \sum_{n=0}^{\infty} B_{2n} C_{n,2} + \frac{L\gamma}{4} \sum_{n=0}^{\infty} \frac{\alpha^{(n)}(0)}{n} \end{aligned} \tag{18}$$

where

$$\begin{aligned} C_{n,1} &= \lambda_1 (2a_1)^{2n-1} (2n-1)! i^{2n-1} \operatorname{erfc} \left(\frac{\gamma_1}{2a_1} \right) \\ C_{n,2} &= \lambda_2 (2a_2)^{2n-1} (2n-1)! i^{2n-1} \operatorname{erfc} \left(\frac{\gamma_1}{2a_2} \right) \end{aligned}$$

From (17) and (18) we can easily find two unknowns in the following form:

$$A_{2n} = \frac{4n \cdot C_{n,2} \cdot f^{(2n+1)}(0) \cdot K_{n,2} + L\gamma \cdot \alpha^{(n)}(0) K_{n,1}}{8n \cdot K_{n,1} \cdot C_{n,1}} \tag{19}$$

$$A_{2n+1} = \frac{4n \cdot C_{n,2} \cdot f^{(2n+1)}(0) \cdot K_{n,2} + L\gamma \cdot \alpha^{(n)}(0) K_{n,1}}{8n \cdot K_{n,2} \cdot C_{n,1}} \tag{20}$$

By using the last condition (5) we get:

$$\lambda_1 \sum_{n=0}^{\infty} (2a_1)^{2n-1} \left[A_{2n}(\tau)^{2n-1} i^{2n-1} \operatorname{erfc}(0) - A_{2n+1}(\tau)^{2n} i^{2n} \operatorname{erfc}(0) \right] = \sum_{n=0}^{\infty} P_n \tau^n \tag{21}$$

The main problem is to find the remaining unknown coefficient we have:

$$\begin{aligned} & \frac{\partial^k}{\partial \tau^k} \left(\lambda_1 \sum_{n=0}^{\infty} \left[(2a_1\tau)^{2n-1} A_{2n} i^{2n-1} \operatorname{erfc} \left(\frac{x}{2a_1\tau} \right) + (2a_1\tau)^{2n} A_{2n+1} i^{2n} \operatorname{erfc} \left(-\frac{x}{2a_1\tau} \right) \right] \right) = \frac{\partial^k}{\partial \tau^k} \left(\sum_{n=0}^{\infty} P_n \tau^n \right) \\ & \lambda_1 \sum_{n=0}^{\infty} \sum_{l=0}^k \left[(2a_1)^{2n-1} A_{2n} \binom{k}{l} \frac{(2n-1)!}{(2n-1-k+l)!} \cdot \tau^{2n-1-k+l} \cdot \frac{\partial^k}{\partial \tau^k} \left(i^{2n-1} \operatorname{erfc} \left(\frac{x}{2a_1\tau} \right) \right) - \right. \\ & \left. - (2a_1)^{2n} A_{2n+1} \binom{k}{l} \frac{(2n)!}{(2n-k+l)!} \cdot \tau^{2n-k+l} \cdot \frac{\partial^k}{\partial \tau^k} \left(i^{2n} \operatorname{erfc} \left(-\frac{x}{2a_1\tau} \right) \right) \right] = \frac{\partial^k}{\partial \tau^k} \left(\sum_{n=0}^{\infty} P_n \tau^n \right) \end{aligned}$$

After taking both parts k-derivatives we can see following expression:

$$\lambda_1 \left[\sum_{n=0}^{\infty} (2a_1)^{2n-1} A_{2n} \frac{(k)!}{(k-2n+1)!} \frac{\partial^{k-2n+1}}{\partial \tau^{k-2n+1}} \left(i^{2n-1} \operatorname{erfc} \left(\frac{x}{2a_1 \tau} \right) \right) \right]_{\tau=0} -$$

$$- (2a_1)^{2n} A_{2n+1} \frac{(k)!}{(k-2n)!} \frac{\partial^{k-2n}}{\partial \tau^{k-2n}} \left(i^{2n} \operatorname{erfc} \left(-\frac{x}{2a_2 \tau} \right) \right) \Big|_{\tau=0} = \left(\sum_{n=0}^{\infty} P_n \frac{k!}{(k-n)!} \right)$$

For this purpose we use the Faa Di Bruno formula (Arbogast formula) for a derivative of a composite function:

$$\frac{\partial^{k-2n+1}}{\partial \tau^{k-2n+1}} \left(i^{2n-1} \operatorname{erfc} \left(\frac{x}{2a_2 \tau} \right) \right) = \sum_{j=0}^{k-2n+1} i^{2n-1} \operatorname{erfc}^{(j)}(\delta) \beta_{k-2n+1,j}(\delta', \delta'', \dots, \delta^{k-2n+1-j})$$

$$i^{2n-1} \operatorname{erfc}^{(j)}(\delta) = \frac{d^j}{d\tau^j} \left(i^{2n-1} \operatorname{erfc}(\delta) \right) \Big|_{\tau=0} = (-1)^{j-2n} \frac{2}{\sqrt{\pi}} H_{j-2n}(\delta) \exp(-\delta^2)$$

$$\delta \Big|_{x=0} = \frac{x}{2a_2 \tau} \Big|_{x=0} = 0 \quad \exp(0) = 1$$

$$\frac{\partial^{k-2n}}{\partial \tau^{k-2n}} \left(i^{2n} \operatorname{erfc} \left(-\frac{x}{2a_2 \tau} \right) \right) = \sum_{j=0}^{k-2n} i^{2n} \operatorname{erfc}^{(j)}(\delta) \beta_{k-2n,j}(\delta', \delta'', \dots, \delta^{k-2n-j})$$

$$i^{2n} \operatorname{erfc}^{(j)}(\delta) = \frac{d^j}{d\tau^j} \left(i^{2n} \operatorname{erfc}(\delta) \right) \Big|_{\tau=0} = (-1)^{j-2n-1} \frac{2}{\sqrt{\pi}} H_{j-2n-1}(\delta) \exp(-\delta^2)$$

Finally, we get following sum:

$$\lambda_1 \left[\sum_{n=0}^{\infty} A_{2n} \frac{(2a_1)^{2n-1} (k)!}{(k-2n+1)!} \sum_{j=0}^{k-2n+1} (-1)^{j-2n} \frac{2}{\sqrt{\pi}} H_{j-2n}(0) \beta_{k-2n+1,j}(\delta', \delta'', \dots, \delta^{k-2n+1-j}) - \right.$$

$$\left. - \sum_{n=0}^{\infty} A_{2n+1} \frac{(2a_1)^{2n} (k)!}{(k-2n)!} \sum_{j=0}^{k-2n} (-1)^{j-2n-1} \frac{2}{\sqrt{\pi}} H_{j-2n-1}(0) \beta_{k-2n,j}(\delta', \delta'', \dots, \delta^{k-2n-j}) \right] = \sum_{n=0}^{\infty} P_n \frac{k!}{(k-n)!}$$

After we can easily find our unknown coefficient:

$$A_{2n} \cdot K_{n,3} - A_{2n+1} \cdot K_{n,4} = P_n \tag{22}$$

1.3 Conclusion

Main result namely coefficients of function A_{2k} , A_{2k+1} (9), B_{2k} , B_{2k+1} (10) and the function $P(t)$ are obtained from (19), (20), (16), (15) and (21-22) respectively. It worth nothing that deviation of solution can provide maximum principle. Moreover if convergence of (9) and (10) proved, wide class of heat equations that describe diverse phenomena can be solved both analytically and numerically.

II. APPROXIMATE SOLUTION OF TWO PHASE INVERSE STEFAN PROBLEM

2.1. Introduction

Exact solutions of heat conduction problems are rather cumbersome and time-consuming. In addition, they are practically absent in the problems of the radial heat flux in spherical coordinates with

the change in the aggregate state [11], 12]. Therefore, to solve practical problems are commonly used charts, obtained by numerical or approximate methods [13]. One of the approximate analytical methods is an integral method of heat balance, which primarily attracted to its physical clarity, simplicity and precision of the results is high enough that demonstrates T. Goodman [14] on numerous examples. The main difficulty to be faced when using the integral method of heat balance is the setting right temperature profile, which according to T. Goodman, significantly affects the accuracy of the results.

There are several approaches in selecting the temperature profiles. In [15] A.I. Veinik offers to use temperature profiles in the form of ordinary polynomials for problems of any geometry, which should simplify the solution of the problem.

In this study we will try to find approximate solution of two Phase Inverse Stefan problem for degenerate domain with moving boundary.

2.2. Problem Solution

Let us consider this problem at the suggestion that $\alpha(t) = \sum_{n=1}^{\infty} \alpha_n t^{n/2}$ is given.

To find the unknown $u_2(x, t)$, we identify the temperature profile corresponding following conditions:

$$u_2(\alpha(t), t) = u_m \tag{23}$$

$$u_2|_{x=\beta(t)} = 0 \tag{24}$$

$$\frac{\partial u_2}{\partial x} \Big|_{x=\beta(t)} = 0 \tag{25}$$

$$u_2|_{x=\infty} = 0 \tag{26}$$

Therefore the temperature profile will be:

$$u_2(x, t) = \begin{cases} u_m \left[\frac{x - \beta(t)}{\alpha(t) - \beta(t)} \right]^2, & \alpha(t) < x < \beta(t) \\ 0, & x > \beta(t) \end{cases} \tag{27}$$

Here we need to find the $\beta(t)$, where $\beta(0) = 0$ by using the equation (2) to use the integral of power balance:

$$\int_{\alpha(t)}^{\beta(t)} \frac{\partial u_2}{\partial t} dx = a_2^2 \frac{\partial u_2}{\partial x} \Big|_{\alpha(t)}^{\beta(t)} \tag{28}$$

$$\int_{\alpha(t)}^{\beta(t)} \frac{\partial u_2}{\partial t} dx = a_2^2 \left[\frac{\partial u_2}{\partial x} \Big|_{x=\beta(t)} - \frac{\partial u_2}{\partial x} \Big|_{x=\alpha(t)} \right]$$

$$\int_{\alpha(t)}^{\beta(t)} \frac{\partial u_2}{\partial t} dx = a_2^2 \left[0 - 2u_m \left[\frac{x - \beta(t)}{\alpha(t) - \beta(t)} \right] \left(\frac{1}{\alpha(t) - \beta(t)} \right) \Big|_{x=\alpha(t)} \right]$$

$$\int_{\alpha(t)}^{\beta(t)} \frac{\partial u_2}{\partial t} dx = -a_2^2 \frac{2u_m}{\alpha(t) - \beta(t)}$$

For the right side we use the Leibniz integral rule, also differentiation under the integral sign:

$$\frac{d}{dx} \left(\int_{\alpha(x)}^{\beta(x)} f(x,t) dt \right) = f(x, \beta(x)) \cdot \beta'(x) - f(x, \alpha(x)) \cdot \alpha'(x) + \int_{\alpha(x)}^{\beta(x)} \frac{\partial}{\partial x} f(x,t) dt$$

Therefore,

$$\int_{\alpha(t)}^{\beta(t)} \frac{\partial}{\partial t} f(x,t) dx = \frac{d}{dt} \left(\int_{\alpha(t)}^{\beta(t)} f(x,t) dx \right) - f(\beta(t), t) \cdot \beta'(t) + f(\alpha(t), t) \cdot \alpha'(t) \quad (29)$$

By using Leibniz's method we get:

$$\left[\frac{d}{dt} \left(\int_{\alpha(t)}^{\beta(t)} u_2 dx \right) - u_2 |_{x=\beta(t)} \frac{d\beta(t)}{dt} + u_2 |_{x=\alpha(t)} \frac{d\alpha(t)}{dt} \right] = -a_2^2 \frac{2u_m}{\alpha(t) - \beta(t)}$$

The solution of given differential equation has been shown in the paper [24].

$$\frac{d\beta(t)}{dt} (\beta(t) - \alpha(t)) + 2 \frac{d\alpha(t)}{dt} (\beta(t) - \alpha(t)) - 6a_2^2 \quad (30)$$

To solve the equation (30) we have used the mathematical program Mathcad 13. To find the solution we considered three particular cases and got following graphs:

$$\frac{d}{dt} y(t)(y(t) - \alpha(t)) + 2 \frac{d\alpha(t)}{dt} (y(t) - \alpha(t)) - 6a_2^2 = 0 \quad (31)$$

Equation (31) have been plotted in the following way for the first particular case when $t = 100$:




Figure 1 $\beta(t)$ at $t = 100$

Equation (31) have been plotted in the following way for the second particular case when $t = 300$:




Figure 2 $\beta(t)$ at $t = 300$

Equation (31) have been plotted in the following way for the third particular case when $t = 500$:




Figure 3 $\beta(t)$ at $t = 500$

This given plots approximately show the value of $\beta(t)$. Therefore, we can consider that we know the value of $\beta(t)$ and consider other cases.

For $u_1(x, t)$ we have the temperature profile $A(t)x^2 + B(t)x + C(t)$ as it is linear. We will use the conditions (5),(6) for $u_1(x, t)$, (7) and (1):

$$u_1(x, t) = A(t)x^2 + B(t)x + C(t), \quad 0 < x < \alpha(t)$$

By using the condition (5) we get:

$$\left. \frac{\partial u_1}{\partial x} \right|_{x=0} = B(t) = -\frac{P(t)}{\lambda_1} \quad (32)$$

By using the condition (6) for $u_1(x, t)$ we have:

$$u_1(\alpha(t), t) = A(t)\alpha^2(t) + B(t)\alpha(t) + C(t) = u_m \quad (33)$$

By using the Stefan condition (7) we get:

$$-\lambda_1(2A(t)\alpha(t) + B(t)) = -\lambda_2 \left(\frac{2u_m}{\alpha(t) - \beta(t)} \right) + L\gamma \frac{d\alpha(t)}{dt}$$

If we equalize the right side of the equation above to $Q(t)$, we will have following expression:

$$2A(t)\alpha(t) + B(t) = -\frac{Q(t)}{\lambda_1} \quad (34)$$

Let us use the last equation (1) and get an integral of power balance:

$$\int_0^{\alpha(t)} \frac{\partial^2 u_1}{\partial x^2} dx = \left. \frac{\partial u_1}{\partial x} \right|_0^{\alpha(t)}$$

$$a_1^2 \left[\left. \frac{\partial u_1}{\partial x} \right|_{x=\alpha(t)} - \left. \frac{\partial u_1}{\partial x} \right|_{x=0} \right] = \int_0^{\alpha(t)} \frac{\partial u_1}{\partial t} dx$$

$$a_1^2 [2A(t) \cdot \alpha(t) + B(t) - B(t)] = a_1^2 [2A(t) \cdot \alpha(t)] = \int_0^{\alpha(t)} \frac{\partial u_1}{\partial t} dx$$

The solutions has been shown in the paper [10].

$$A'(t) [2\alpha^2(t) + 4\alpha'(t) \cdot \alpha(t)] + A(t) [4\alpha'(t) + 4(\alpha''(t) \cdot \alpha(t) + (\alpha'(t))^2) - 12a_1^2] +$$

$$+ \left[\frac{d}{dt} \left[\frac{3\alpha'(t)Q(t)}{\lambda_1} + \frac{3\alpha(t)Q'(t)}{\lambda_1} \right] - u_m \frac{\alpha'(t)}{\alpha(t)} \right] = 0 \quad (35)$$

If we denote the functions that is multiplied to $A'(t)$, $A(t)$ and some functions by $Z_1(t)$, $Z_2(t)$ and $Z_3(t)$ respectively, then we get:

$$A'(t) \cdot Z_1(t) + A(t) \cdot Z_2(t) + Z_3(t) = 0$$

We can easily rewrite our differential equation in the following form:

$$y'(x) + y(x) \cdot S(x) + T(x) = 0$$

By solving our differential equation by Integrating Factor method that has been shown in the paper [10], we can show the solution of givendifferential equation:

$$A(t) = -e^{-\int S(t)dt} \cdot \left(\int T(t) \cdot e^{\int S(t)dt} dt \right) + c \cdot e^{-\int S(t)dt} \quad (36)$$

where $S(t) = Z_2(t) / Z_1(t)$ and $T(t) = Z_3(t) / Z_1(t)$. Let us denote the right side of above expression as $K(t)$, therefore we can show the temperature profile for $u_1(x, t)$ and reconstruct the boundary function $P(t)$:

$$u_1(x, t) = K(t)x^2 - \left[\frac{Q(t)}{\lambda_1} + 2K(t)\alpha(t) \right] x + \left[u_m + K(t)\alpha^2(t) + \frac{Q(t)\alpha(t)}{\lambda_1} \right] \quad (37)$$

$$P(t) = Q(t) + \frac{2K(t)\alpha(t)}{\lambda_1} \quad (38)$$

2.4. Conclusion

In this chapter for solving problems of heat conduction in spherical geometry by integral method of power balance for the first time evaluated the effectiveness of the temperature type profiles: $T(r,t) = \text{polynomial}$ and $T(r,t) = \text{polynomial} / r$ with polynomials of degree n . The results show that in dealing with problems of thermal conductivity for different geometry approximate integral method of power balance is not always possible to achieve sufficient accuracy when using temperature profiles of the same type.

REFERENCES

- [1] Alexiades V. and Solomon A. D. *Mathematical Modeling of Melting and Freezing Processes*, Taylor and Francis, Washington, DC, 1993.
- [2] Grassi M., Grassi G. Mathematical modelling and controlled drug delivery: matrix systems, *Current Drug Delivery* 2 (2005) 97–116.
- [3] Kharin S.N., Sarsengeldin M. Influence of Contact Materials on Phenomena in a Short Electrical Arc. TransTechPublications. – Switzerland.
- [4] Tarzia D. A. *A bibliography on moving-free boundary problems for the heat-diffusion equation. The Stefan and related problems*, 2000.
- [5] Tikhonov A.N., Samarski A.A. *Equations of Mathematical Physics*.
- [6] Friedman A. *Free boundary problems for parabolic equations I. Melting of solids*, *J. Math. Mech.*, 8 (1959), pp. 499–517.
- [7] Rubinstein L. I. *The Stefan Problem*, Transl. Math. Monogr. 1971.
- [8] Sarsengeldin M.M. “Solution Of Heat Equation With Moving Boundaries Obtained By Hartree Functions”, International scientific conference.
- [9] Kharin S.N., Sarsengeldin M.M., Temirkul A. “Analytical Solution Of The Third Boundary-Value Problem For The Heat Equation By IEF Method”.
- [10] Kassabek M. S., Sagidolla B.M. “Approximate Solution of Two phase Inverse Stefan problem” (accepted), 2016.
- [11] Carslaw G., Eger D. *Thermal conductivity of solids*, 1964. – 488 p.
- [12] ЛукoвA.V. *Теория теплопроводности*, 1967. – 600 p.
- [13] Pehovich A.I., Zhidkih *Расчеты теплового режима твердых тел*, 1976.
- [14] Goodman T. *Application of integral method in nonlinear problems of non-stationary heat exchange*, 1967. – p. 47–96.
- [15] Veinik A.I. *Приближенный расчет процессов теплопроводности*, 1959.

НАҚТЫ ЖӘНЕ ЖУЫҚТАЛҒАН ЕКІ ФАЗАЛЫ КЕРІ СТЕФАН МӘСЕЛЕЛЕРІНІҢ ШЕШІМІ

М.М. Сарсенгельдин, С. Қасабек, Б.М. Сағидолла

Сүлеймен Демирел атындағы университет, Қаскелең, Қазақстан

Тірек сөздер: Интегралды функциялар қателіктері әдісі, Интегралды жылулық тепе-теңдік әдісі, Екі фазалы кері Стефан мәселесі.

Аннотация. Берілген жұмыстың негізгі идеясы, қозғалатын шекаралары бар нұқсанды өңір үшін екі фазалық кері Стефан мәселенің нақты шешімдерін табу болып табылады.

Төменде осы мәселелерді жауап қадағалау ұйымдастырылады. Аналитикалық шешім табу үшін біз негізінен С.Н. Харин жұмыстарында ұсынған әдістері Фаа Ди Бруно формуласын интегралды қателіктері функцияларына қолданамыз. Интегралды қателіктері функциясы және оның жаңа әдістерінің қасиеттері де анық көрінген.

Стефан мәселесі нақты шешім үшін күрделі болып табылады. Сондықтан, біз шамамен шешімін табу үшін, X және t арқылы интегралдап алынған дифференциалдық теңдеулерге температуралық профильдер тағайындаймыз. Бұл шамамен шешімін табу үшін С.Н. Харинның ұсынған әдістерін қолдана отырып, интегралды жылулық тепе-теңдік әдісін қолдандық.

Поступила 17.06.2016 г.

СОДЕРЖАНИЕ

Процессы в околоземном космическом пространстве

<i>Яковец А.Ф., Гордиенко Г.И., Жумабаев Б.Т., Литвинов Ю.Г., Абдрахманов Н.</i> Статистика ночных увеличений электронной концентрации в максимуме F2-слоя.....	5
<i>Сомских В.М.</i> О природе бифуркации динамических систем.....	11
<i>Жантаев Ж.Ш., Грищенко В.Ф., Мукушев А.</i> Схемотехническое моделирование защиты электронной аппаратуры от электростатического разряда.....	15
<i>Антонова В.П., Крюков С.В., Луценко В.Ю., Чубенко А.П.</i> Эффекты землетрясений в интенсивности нейтронов тепловых энергий на высокогорной станции Северного Тянь-Шаня.....	20
<i>Салихов Н.М.</i> Новый метод регистрации динамики вспышек ионизации в ионосфере аппаратно-программным комплексом доплеровских измерений на наклонной радиотрассе.....	27

Наземно-космические методы исследования геодинимических процессов в земной коре

<i>Вилев А.В., Жантаев Ж.Ш., Стихарный А.П.</i> Динамика сезонных движений GPS станций на территории Северного Тянь-Шаня.....	34
<i>Хачикян Г.Я., Жумабаев Б.Т., Тойшиев Н.С., Калдыбаев А., Нуракынов С.</i> Вариации солнечной активности и пространственно-временное распределение сильных землетрясений ($M \geq 7.0$) на территории Евразии в 1973-2014 гг.....	40
<i>Бибосинов А.Ж., Шигаев Д.Т., Калдыбаев А.А., Нуракынов С.М., Бреусов Н.Г., Мамырбек Г.Б.</i> Исследование Шардаринского гидрокомплекса методом георадиолокации.....	46
<i>Бибосинов А.Ж., Нуракынов С.М., Калдыбаев А.А., Шигаев Д.Т.</i> Эффективность применения георадиолокационного метода при изучении инженерно-геологических условий на участках Алматинского метрополитена приповерхностного залегания.....	50
<i>Шигаев Д.Т., Мунсызбай Т.М.</i> Маломощная солнечная теплоэлектростанция с максимальным использованием энергии Солнца.....	56
<i>Жантаев Ж.Ш., Хачикян Г.Я., Кайраткызы Д., Андреев А.</i> Долговременные тренды в вариациях продолжительности земных суток и частоты возникновения на планете землетрясений.....	62
<i>Хачикян Г.Я., Жумабаев Б.Т., Сералиев А., Хасанов Э.</i> Пространственное распределение характеристик главного геомагнитного поля и эпицентров глубокофокусных ($h > 350$ км) землетрясений по данным 1973-2014 гг.....	67

<i>Исанова М.К., Коданова С.К., Рамазанов Т.С., Бастыкова Н.Х., Габдуллин М.Т., Молдабеков Ж.А.</i> Сечение рассеяния и тормозная способность в плотной плазме: влияние эффектов дифракции и динамического экранирования.....	73
<i>Кудайкулов А.А., Жозеранд К., Калтаев А.</i> Численное исследование процесса пальцеобразования при течении двух не смешивающихся жидкостей в канале.....	86
<i>Ахметов Б.С., Корченко А.А., Жумангалиева Н.К.</i> Модель решающих правил для обнаружения аномалий в информационных системах.....	91
<i>Бапаев К.Б., Сламжанова С.С., Исаева Г.Б.</i> О дискретных неравенствах.....	101
<i>Боос Э.Г., Альменова А.М., Жуков В.В., Садыков Т.Х., Степанов А., Таутаев Е.М.</i> Исследование взаимодействий частиц космического излучения методом радиоизлучения на высоте 3340 метров над уровнем моря.....	110
<i>Джакупов К.Б.</i> О моделировании динамики вязкой жидкости уравнениями ротора скорости и функции тока.....	117
<i>Джакупов К.Б.</i> Эффективное применение уравнений максвелла и закона ома в численном моделировании двухфазных процессов магнитной гидродинамики.....	124
<i>Исадыков А.Н., Иванов М.А., Сахиев С.К., Жаугашиева С.А., Нурбакова Г.С., Мукушев Б.А.</i> Вычисление ширины распада $\omega(782)$ мезона для реакции $\omega \rightarrow \pi^+ + \pi^- + \pi^0$ в ковариантной модели кварков.....	135
<i>Калмурзаев Б.С.</i> О полурешетках роджерса двухэлементных семейств разностей в п. множеств.....	141
<i>Кошеров Т.С., Жумабекова Г.Е.</i> Исследование структуры и фазового состава поверхности кремния при температурном и лазерном воздействии.....	147
<i>Кошеров Т.С., Көшікінбай Б.Қ.</i> Особенности напряженного состояния пластин кремния в процессе термического отжига.....	156
<i>Курманбаев Д.М.</i> Солитонная деформация поверхности энепера третьего порядка.....	163
<i>Майлебаева Д., Тилегенова Д.</i> Метод параметризации при решении трансцендентных уравнений.....	168
<i>Мамаев Ш.М., Даниярбек Р.Н.</i> Ұзындығы шектелген стерженде пластикалық облыстың және кернеуді жеңілдету толқындарының құрылуын торлық-характеристика әдісімен зерттеу.....	173
<i>Оңгарбаева А.Д.</i> Электрондық білім беру ресурстарын оқу процесінде болашақ мұғалімдерді оқытуда қолдану.....	184
<i>Сүйменбаев Б.Т., Алексеева Л.А., Сүйменбаева Ж.Б., Гусейнов С.Р.</i> Моделирование динамики космического аппарата в гравимагнитном поле земли в системе «MATLAB SIMULINK».....	188
<i>Туленбаев К.М., Шаймарданова Ж.Н., Габдуллин Б.</i> Структурные свойства (α, β) – коммутативных алгебр.....	208
<i>Сарсенгельдин М.М., Касабек С., Сагидолла Б.М.</i> Точное и приближенное решения двухфазовой обратной задачи Стефана.....	214