ҚАЗАҚСТАН РЕСПУБЛИКАСЫ ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ Әль-фараби атындағы Қазақ ұлттық университетінің

ХАБАРЛАРЫ

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК РЕСПУБЛИКИ КАЗАХСТАН Казахский национальный университет имени Аль-фараби

NEWS

OF THE NATIONAL ACADEMY OFSCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-farabi kazakh
national university

SERIES PHYSICO-MATHEMATICAL

1 (323)

JANUARY - FEBRUARY 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

Бас редакторы ф.-м.ғ.д., проф., ҚР ҰҒА академигі **Ғ.М. Мұтанов**

Редакция алкасы:

Жұмаділдаев А.С. проф., академик (Қазақстан)

Кальменов Т.Ш. проф., академик (Қазақстан)

Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)

Өмірбаев У.У. проф. корр.-мүшесі (Қазақстан)

Жусіпов М.А. проф. (Қазақстан)

Жұмабаев Д.С. проф. (Қазақстан)

Асанова А.Т. проф. (Қазақстан)

Бошкаев К.А. PhD докторы (Қазақстан)

Сураған Д. корр.-мүшесі (Қазақстан)

Quevedo Hernando проф. (Мексика),

Джунушалиев В.Д. проф. (Қырғыстан)

Вишневский И.Н. проф., академик (Украина)

Ковалев А.М. проф., академик (Украина)

Михалевич А.А. проф., академик (Белорус) **Пашаев А.** проф., академик (Әзірбайжан)

Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары

Тигиняну И. проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18, http://physics-mathematics.kz/index.php/en/archive

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)

Кальменов Т.Ш. проф., академик (Казахстан)

Жантаев Ж.Ш. проф., чл.-корр. (Казахстан)

Умирбаев У.У. проф. чл.-корр. (Казахстан)

Жусупов М.А. проф. (Казахстан)

Джумабаев Д.С. проф. (Казахстан)

Асанова А.Т. проф. (Казахстан)

Бошкаев К.А. доктор PhD (Казахстан)

Сураган Д. чл.-корр. (Казахстан)

Quevedo Hernando проф. (Мексика),

Джунушалиев В.Д. проф. (Кыргызстан)

Вишневский И.Н. проф., академик (Украина)

Ковалев А.М. проф., академик (Украина)

Михалевич А.А. проф., академик (Беларусь)

Пашаев А. проф., академик (Азербайджан)

Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.

Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: POO «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год. Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,

http://physics-mathematics.kz/index.php/en/archive

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

Editor in chief doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

Editorial board:

Dzhumadildayev A.S. prof., academician (Kazakhstan)

Kalmenov T.Sh. prof., academician (Kazakhstan)

Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)

Umirbayev U.U. prof. corr. member. (Kazakhstan)

Zhusupov M.A. prof. (Kazakhstan)

Dzhumabayev D.S. prof. (Kazakhstan)

Asanova A.T. prof. (Kazakhstan)

Boshkayev K.A. PhD (Kazakhstan)

Suragan D. corr. member. (Kazakhstan)

Quevedo Hernando prof. (Mexico),

Dzhunushaliyev V.D. prof. (Kyrgyzstan)

Vishnevskyi I.N. prof., academician (Ukraine)

Kovalev A.M. prof., academician (Ukraine)

Mikhalevich A.A. prof., academician (Belarus)

Pashayev A. prof., academician (Azerbaijan)

Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.

Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

http://physics-mathematics.kz/index.php/en/archive

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

— 4 —

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

https://doi.org/10.32014/2019.2518-1726.2

Volume 1, Number 323 (2019), 14 – 21

UDK 517.951 MRNTI 27.31.15

A.T.Assanova¹, A.A.Boichuk², Z.S.Tokmurzin³

¹ Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan; ² Institute of Mathematics NAS Ukraine, Kyiv, Ukraine; ³ K.Zhubanov Aktobe Regional State University, Aktobe, Kazakhstan assanova@math.kz; boichuk.aa@gmail.com; tokmurzinzh@gmail.com;

ON THE INITIAL-BOUNDARY VALUE PROBLEM FOR SYSTEM OF THE PARTIAL DIFFERENTIAL EQUATIONS OF FOURTH ORDER

Abstract. A initial-boundary value problem for system of the partial differential equations of fourth order is considered. We study the existence of classical solutions to the initial-boundary value problem for system of the partial differential equations of fourth order and offer the methods for finding its approximate solutions. Sufficient conditions for the existence and uniqueness of a classical solution to the initial-boundary value problem for system of the partial differential equations of fourth order are set. By introducing of a new unknown functions, we reduce the considered problem to an equivalent problem consisting of a nonlocal problem for the system of hyperbolic equations of second order with functional parameters and the integral relations. We offer the algorithm for finding an approximate solution to the investigated problem and prove its convergence. Sufficient conditions for the existence of unique solution to the equivalent problem with parameters are established. Conditions of unique solvability to the initial-boundary value problem for system of the partial differential equations of fourth order are obtained in the terms of initial data. Separately, the result is given for the initial-periodic in time boundary value problem.

Keywords: system of the partial differential equations of fourth order, initial-boundary value problem, nonlocal problem, system of the hyperbolic equations of second order, solvability, algorithm.

1. Introduction. Currently, the problems of mathematical physics connected with the description of wave motion of liquids of different nature are drawn by great attention. This interest is caused not only by big applied importance of these problems, but their new theoretical and mathematical content often do not have analogues in the classical mathematical physics. One of the important classes of such problems are the initial-boundary value problems for fourth order partial differential equations. To date, various methods for researching and solving the initial-boundary value problems for fourth order partial differential equations of hyperbolic and composite types are developed in [1-12]. In order to investigate various boundary value problems for fourth order partial differential equations along with the classical methods of mathematical physics (the Fourier method, the method of Green's functions, Poincare's metric concept) we apply the method of differential inequalities and other methods of qualitative theory of ordinary differential equations. Based on them, the conditions for solvability of considered boundary value problems are obtained, and the ways for finding their solutions are offered. Fourth order system of partial differential equations began to be studied relatively recently.

In the present work we consider system of the partial differential equations of fourth order at the rectangular domain. Boundary condition for time variable are specified as a combination of values from the partial derivatives of required solution on third orders by spatial variable. We investigate the questions of existence and uniqueness of the classical solution to initial-boundary value problem for system of the partial differential equations of fourth order and its applications.

2. Methods. For solve to considered problem we use a method of introduction additional functional parameters [13-29]. The original problem is reduced to an equivalent problem consisting of nonlocal problem for system of the hyperbolic equations of second order with functional parameters and integral relations. Sufficient conditions for the unique solvability to investigated problem are established in the terms of initial data. Algorithms for finding solution to the equivalent problem are constructed. Conditions of unique solvability to initial-boundary value problem for system of partial differential equations of fourth order are established in the terms of system's coefficients and boundary matrices. Separately, the result is given for the initial-periodic in time boundary value problem.

Note that, in [30, 31] a similar approach has been applied to the initial-boundary value problem and nonlocal problem for the system of partial differential equations of third order.

2. Statement of problem. At the domain $\Omega = [0,T] \times [0,\omega]$ we consider the initial-periodic boundary value problem for system of the partial differential equations of fourth order in the following form

$$\frac{\partial^{4} u}{\partial t \partial x^{3}} = A_{1}(t, x) \frac{\partial^{3} u}{\partial x^{3}} + A_{2}(t, x) \frac{\partial^{3} u}{\partial t \partial x^{2}} + A_{3}(t, x) \frac{\partial^{2} u}{\partial x^{2}} + A_{4}(t, x) \frac{\partial^{2} u}{\partial t \partial x} + A_{5}(t, x) \frac{\partial u}{\partial x} + A_{6}(t, x) \frac{\partial u}{\partial t} + A_{7}(t, x)u + f(t, x), \qquad (t, x) \in \Omega,$$
(1)

$$\frac{\partial^3 u(0,x)}{\partial x^3} = K(x) \frac{\partial^3 u(T,x)}{\partial x^3} + \varphi(x), \qquad x \in [0,\omega],$$
 (2)

$$u(t,0) = \psi_0(t), \qquad t \in [0,T],$$
 (3)

$$\frac{\partial u(t,x)}{\partial x}\Big|_{x=0} = \psi_1(t), \qquad t \in [0,T], \tag{4}$$

$$\frac{\partial^2 u(t,x)}{\partial x^2}\Big|_{x=0} = \psi_2(t), \qquad t \in [0,T], \tag{5}$$

where $u(t,x) = col(u_1(t,x), u_2(t,x), ..., u_n(t,x))$ is unknown function, the $n \times n$ -matrices $A_i(t,x)$, $i = \overline{1,7}$, and n-vector function f(t,x) are continuous on Ω , the $n \times n$ -matrix K(x) and n-vector-function $\varphi(x)$ are continuous on $[0,\omega]$, the n-vector-functions $\psi_0(t)$, $\psi_1(t)$ and $\psi_2(t)$ are continuously differentiable on [0,T]. The initial data satisfy the condition of approval.

A function
$$u(t,x) \in C(\Omega,R^n)$$
 having partial derivatives $\frac{\partial u(t,x)}{\partial x} \in C(\Omega,R^n)$, $\frac{\partial u(t,x)}{\partial t} \in C(\Omega,R^n)$, $\frac{\partial^2 u(t,x)}{\partial x^2} \in C(\Omega,R^n)$, $\frac{\partial^2 u(t,x)}{\partial t \partial x} \in C(\Omega,R^n)$, $\frac{\partial^3 u(t,x)}{\partial t \partial x} \in C(\Omega,R^n)$, $\frac{\partial^3 u(t,x)}{\partial t \partial x^3} \in C(\Omega,R^n)$, is called a classical solution to problem (1)--(5) if it satisfies system (1) for all $(t,x) \in \Omega$, and boundary conditions (2)--(5).

We will investigate the questions of existence and uniqueness of the classical solutions to the initial-boundary value problem for system of the partial differential equations of fourth order (1)--(5) and the approaches of constructing its approximate solutions. For this goals, we applied the method of introduction additional functional parameters proposed in [13-31] for solving of various nonlocal problems for systems of hyperbolic equations with mixed derivatives. Considered problem is provided to nonlocal problem for the system of hyperbolic equations of second order including additional functions and integral relation. The algorithm for finding the approximate solution of the investigated problem is proposed and its convergence proved. Sufficient conditions of the existence unique classical solution to problem (1)--(5) are obtained in the terms of initial data.

3. Scheme of the method and reduction to equivalent problem. We introduce a new unknown functions $v(t,x) = \frac{\partial u(t,x)}{\partial x}$, $w(t,x) = \frac{\partial^2 u(t,x)}{\partial x^2}$ and rewrite the problem (1)--(5) in the following form

$$\frac{\partial^2 w}{\partial t \partial x} = A_1(t, x) \frac{\partial w}{\partial x} + A_2(t, x) \frac{\partial w}{\partial t} + A_3(t, x) w + f(t, x) + \frac{\partial^2 w}{\partial t} +$$

$$+ A_4(t,x)\frac{\partial v(t,x)}{\partial t} + A_5(t,x)v(t,x) + A_6(t,x)\frac{\partial u(t,x)}{\partial t} + A_7(t,x)u(t,x), \qquad (t,x) \in \Omega,$$
 (6)

$$\frac{\partial w(0,x)}{\partial x} = K(x)\frac{\partial w(T,x)}{\partial x} + \varphi(x), \qquad x \in [0,\omega], \tag{7}$$

$$w(t,0) = \psi_2(t), \qquad t \in [0,T],$$
 (8)

$$v(t,x) = \psi_0(t) + \int_0^x w(t,\xi)d\xi, \quad u(t,x) = \psi_0(t) + \psi_1(t)x + \int_0^x \int_0^\xi w(t,\xi_1)d\xi_1d\xi, \quad (t,x) \in \Omega.$$
 (9)

Here the conditions (3) and (4) are taken into account in (9).

A triple functions (w(t,x),v(t,x),u(t,x)), where the function $w(t,x) \in C(\Omega,R^n)$ has partial derivatives $\frac{\partial w(t,x)}{\partial x} \in C(\Omega,R^n)$, $\frac{\partial w(t,x)}{\partial t} \in C(\Omega,R^n)$, $\frac{\partial^2 w(t,x)}{\partial t\partial x} \in C(\Omega,R^n)$, the functions $v(t,x) \in C(\Omega,R^n)$ and $u(t,x) \in C(\Omega,R^n)$ have partial derivatives $\frac{\partial v(t,x)}{\partial x} \in C(\Omega,R^n)$, $\frac{\partial^2 v(t,x)}{\partial x} \in C(\Omega,R^n)$, $\frac{\partial^2 v(t,x)}{\partial t\partial x} \in C(\Omega,R^n)$, $\frac{\partial^2 u(t,x)}{\partial t\partial x} \in C(\Omega,R^n)$, is called a solution to problem (6)-(9) if it satisfies of the system of hyperbolic equations of second order (6) for all $(t,x) \in \Omega$, the boundary conditions (7), (8), and the integral relations (9).

At fixed v(t,x) and u(t,x) the problem (6)--(8) is a nonlocal problem for the system of hyperbolic equations with respect to w(t,x) on Ω . The integral relations (9) allow us to determine the unknown functions v(t,x) and u(t,x) for all $(t,x) \in \Omega$.

4. Algorithm. The unknown function w(t,x) will be determined from nonlocal problem for the system of hyperbolic equations (6)--(8). The unknown functions v(t,x) and u(t,x) will be found from integral relations (9).

If we know the functions v(t,x) and u(t,x), then from nonlocal problem (6)--(8) find the function w(t,x). Conversely, if we known the functions v(t,x) and u(t,x), then from nonlocal problem (6)--(8) we find the function w(t,x). Since the functions v(t,x), u(t,x) and v(t,x) are unknowns together, for finding of the solution to problem (6)--(9) we use an iterative method. The solution to problem (6)--(9) is the triple functions $(w^*(t,x),v^*(t,x),u^*(t,x))$ we defined as a limit of sequence of triples $(w^{(k)}(t,x),v^{(k)}(t,x),u^{(k)}(t,x))$, k=0,1,2,..., according to the following algorithm:

Step 0. 1) Suppose in the right-hand part of the system (6)
$$\frac{\partial v(t,x)}{\partial t} = \dot{\psi}_0(t)$$
, $v(t,x) = \psi_0(t)$,

 $\frac{\partial u(t,x)}{\partial t} = \dot{\psi}_0(t) + \dot{\psi}_1(t)x, \text{ and } u(t,x) = \psi_0(t) + \psi_1(t)x, \text{ from nonlocal problem (6)--(8) we find the initial approximation } w^{(0)}(t,x) \text{ for all } (t,x) \in \Omega;$

2) From the integral relations (9) under $w(t,x) = w^{(0)}(t,x)$, we find the functions $v^{(0)}(t,x)$ and $u^{(0)}(t,x)$ for all $(t,x) \in \Omega$.

Step 1. 1) Suppose in the right-hand part of system (6) $\frac{\partial v(t,x)}{\partial t} = \frac{\partial v^{(0)}(t,x)}{\partial t}$, $v(t,x) = v^{(0)}(t,x)$,

 $\frac{\partial u(t,x)}{\partial t} = \frac{\partial u^{(0)}(t,x)}{\partial t}, \text{ and } u(t,x) = u^{(0)}(t,x), \text{ from nonlocal problem (6)--(8) we find the first approximation } w^{(1)}(t,x) \text{ for all } (t,x) \in \Omega.$

2) From the integral relations (9) under $w(t,x) = w^{(1)}(t,x)$, we find the functions $v^{(1)}(t,x)$ and $u^{(1)}(t,x)$ for all $(t,x) \in \Omega$.

And so on.

Step k. 1) Suppose in the right-hand part of system (6) $\frac{\partial v(t,x)}{\partial t} = \frac{\partial v^{(k-1)}(t,x)}{\partial t}$,

$$v(t,x) = v^{(k-1)}(t,x), \quad \frac{\partial u(t,x)}{\partial t} = \frac{\partial u^{(k-1)}(t,x)}{\partial t}, \text{ and } u(t,x) = u^{(k-1)}(t,x), \text{ from nonlocal problem (6)}$$

(8) we find the k-th approximation $w^{(k)}(t,x)$ for all $(t,x) \in \Omega$:

$$\frac{\partial^{2} w^{(k)}}{\partial t \partial x} = A_{1}(t, x) \frac{\partial w^{(k)}}{\partial x} + A_{2}(t, x) \frac{\partial w^{(k)}}{\partial t} + A_{3}(t, x) w^{(k)} + f(t, x) + A_{4}(t, x) \frac{\partial v^{(k-1)}(t, x)}{\partial t} + A_{5}(t, x) v^{(k-1)}(t, x) + A_{6}(t, x) \frac{\partial u^{(k-1)}(t, x)}{\partial t} + A_{7}(t, x) u^{(k-1)}(t, x), \quad (t, x) \in \Omega, \quad (10)$$

$$\frac{\partial w^{(k)}(0, x)}{\partial t} + K_{5}(t, x) v^{(k)}(T, x) + K_{5}(t, x) v^{(k)}(T, x) + K_{7}(t, x) u^{(k-1)}(t, x), \quad (t, x) \in \Omega, \quad (10)$$

$$\frac{\partial w^{(k)}(0,x)}{\partial x} = K(x)\frac{\partial w^{(k)}(T,x)}{\partial x} + \varphi(x), \qquad x \in [0,\omega],$$
(11)

$$w^{(k)}(t,0) = \psi_2(t), \qquad t \in [0,T].$$
 (12)

2) From the integral relations (9) under $w(t,x) = w^{(k)}(t,x)$, we find the functions $v^{(k)}(t,x)$ and $u^{(k)}(t,x)$ for all $(t,x) \in \Omega$:

$$v^{(k)}(t,x) = \psi_0(t) + \int_0^x w^{(k)}(t,\xi)d\xi , \quad u^{(k)}(t,x) = \psi_0(t) + \psi_1(t)x + \int_0^x \int_0^\xi w^{(k)}(t,\xi_1)d\xi_1d\xi , \quad (t,x) \in \Omega .$$
 (13) Here $k = 1,2,3,...$

5. The main results. The following theorem gives conditions of feasibility and convergence of the constructed algorithm and the conditions of the existence unique solution to problem (6)--(9).

Theorem 1. Suppose that

- i) the $n \times n$ -matrices $A_i(t,x)$, $i = \overline{1,7}$, and n-vector function f(t,x) are continuous on Ω ;
- ii) the $n \times n$ -matrix K(x) and n-vector-function $\varphi(x)$ are continuous on $[0, \omega]$;
- iii) the n-vector-functions $\psi_0(t)$, $\psi_1(t)$ and $\psi_2(t)$ are continuously differentiable on [0,T];
- iv) the $n \times n$ -matrix $Q(x) = I K(x) \left[I + \int_{0}^{T} A_{1}(\tau, x) d\tau \right]$ is invertible for all $x \in [0, \omega]$, where

I is unit matrix on dimension n.

Then the nonlocal problem for system of the hyperbolic equations with parameters (6)--(9) has a unique solution $(w^*(t,x),v^*(t,x),u^*(t,x))$ as a limit of sequences $(w^{(k)}(t,x),v^{(k)}(t,x),u^{(k)}(t,x))$

defining by the algorithm proposed above for k = 0,1,2,...

Proof. Let the conditions i - iv of the Theorem be satisfied. From the 0th step of the above algorithm and Theorem 1 from [21] it follows that the nonlocal problem for system of the hyperbolic equations

$$\frac{\partial^2 w}{\partial t \partial x} = A_1(t, x) \frac{\partial w}{\partial x} + A_2(t, x) \frac{\partial w}{\partial t} + A_3(t, x) w + f(t, x) + A_4(t, x) \dot{\psi}_0(t) + A_5(t, x) \psi_0(t) + A_5(t, x)$$

$$\frac{\partial w(0,x)}{\partial x} = K(x)\frac{\partial w(T,x)}{\partial x} + \varphi(x), \qquad x \in [0,\omega],$$
(15)

$$w(t,0) = \psi_2(t), \qquad t \in [0,T]$$
 (16)

has a unique classical solution $w^{(0)}(t,x)$ for all $(t,x) \in \Omega$.

Further we determine the functions $v^{(0)}(t,x)$ and $u^{(0)}(t,x)$ from the integral relations

$$v^{(0)}(t,x) = \psi_0(t) + \int_0^x w^{(0)}(t,\xi)d\xi, \quad u^{(0)}(t,x) = \psi_0(t) + \psi_1(t)x + \int_0^x \int_0^\xi w^{(0)}(t,\xi_1)d\xi_1d\xi$$

for all $(t,x) \in \Omega$. Functions $v^{(0)}(t,x)$ and $u^{(0)}(t,x)$ together with their partial derivatives $\frac{\partial v^{(0)}(t,x)}{\partial t}$ and $\frac{\partial u^{(0)}(t,x)}{\partial t}$, respectively, are continuous on Ω .

Continuing the iterative process according to the above algorithm, we define successive approximations $w^{(k)}(t,x)$, $v^{(k)}(t,x)$ and $u^{(k)}(t,x)$ for all $(t,x) \in \Omega$ and k = 1,2,...

The conditions i) – iv) of Theorem provide the uniform convergence on Ω of the sequences $\left\{w^{(k)}(t,x)\right\}$, $\left\{v^{(k)}(t,x)\right\}$ and $\left\{u^{(k)}(t,x)\right\}$ as $k\to\infty$ to functions $w^*(t,x)$, $v^*(t,x)$ and $u^*(t,x)$, respectively, for all $(t,x)\in\Omega$. In addition, there are finite limits of sequences of their partial derivatives as $k\to\infty$.

The triple founded functions $(w^*(t,x),v^*(t,x),u^*(t,x))$ has all the required continuous partial derivatives on Ω and be solution to problem (6)—(9). Uniqueness of solution to problem (6)—(9) is proved by method of contradiction.

Theorem 1 is proved.

From the equivalence of problems (6)—(9) and (1)—(5) it follows

Theorem 2. Suppose that the conditions i) - iv) of Theorem 1 are fulfilled.

Then the initial-periodic boundary value problem for system of the partial differential equations of fourth order (1)--(5) has a unique classical solution $u^*(t,x)$.

For K(x) = I and $\varphi(x) = 0$ we obtain the initial-periodic boundary value problem for system of the partial differential equations of fourth order (1), (3)-(5) with condition

$$\frac{\partial^3 u(0,x)}{\partial x^3} = \frac{\partial^3 u(T,x)}{\partial x^3}, \qquad x \in [0,\omega].$$
 (2')

Then the following assertion is true.

Theorem 3. Suppose that

- 1) the $n \times n$ -matrices $A_i(t,x)$, $i = \overline{1,7}$, and n-vector function f(t,x) are continuous on Ω ;
- 2) the *n*-vector-functions $\psi_0(t)$, $\psi_1(t)$ and $\psi_2(t)$ are continuously differentiable on [0,T];
- 3) the $n \times n$ -matrix $Q(x) = \int_{0}^{T} A_{1}(\tau, x) d\tau$ is invertible for all $x \in [0, \omega]$.

Then the initial-periodic boundary value problem for system of the partial differential equations of fourth order (1), (2'), (3)--(5) has a unique classical solution.

Funding. This results are partially supported by grant of the Ministry education and science of Republic Kazakhstan No. AP 05131220 for 2018-2020 years.

УДК 517.951 МРНТИ 27.31.15

А.Т. Асанова¹, А.А. Бойчук², Ж.С. Токмурзин³

¹Институт математики и математического моделирования, Алматы, Казахстан; ² Институт математики НАН Украины, Киев, Украина; ³ Актобинский региональный государственный университет им. К.Жубанова, Актобе, Казахстан

О НАЧАЛЬНО-КРАЕВОЙ ЗАДАЧЕ ДЛЯ СИСТЕМЫ ДИФФЕРЕНЦИАЛЬНЫХ УРАВНЕНИЙ В ЧАСТНЫХ ПРОИЗВОДНЫХ ЧЕТВЕРТОГО ПОРЯДКА

Аннотация. Рассматривается начально-краевая задача для системы дифференциальных уравнений в частных производных четвертого порядка. Исследуются вопросы существования классического решения начально-краевой задачи для системы дифференциальных уравнений в частных производных четвертого порядка и предлагаются методы нахождения их приближенных решений. Установлены достаточные условия существования и единственности классического решения начально-краевой задачи для системы дифференциальных уравнений в частных производных четвертого порядка. Путем введения новых неизвестных функций исследуемая задача сведена к эквивалентной задаче, состоящей из нелокальной задачи для системы гиперболических уравнений второго порядка с функциональными параметрами и интегральных соотношений. Предложены алгоритмы нахождения приближенного решения исследуемой задачи и доказана их сходимость. Установлены достаточные условия существования единственного решения эквивалентной задачи с параметрами. Условия однозначной разрешимости начально-краевой задачи для системы дифференциальных уравнений в частных производных четвертого порядка получены в терминах исходных данных. Отдельно приводится результат для начально-периодической по времени краевой задачи.

Ключевые слова: система дифференциальных уравнений в частных производных четвертого порядка, начально-краевая задача, нелокальная задача, система гиперболических уравнений второго порядка, разрешимость, алгоритм.

УДК 517.951 МРНТИ 27.31.15

А.Т. Асанова¹, А.А. Бойчук², Ж.С. Токмурзин³

¹Математика және математикалық моделдеу институты, Алматы, Қазақстан;
² Украина ҰҒА Математика институты, Киев, Украина;
³ Қ.Жұбанов ат. Ақтөбе өңірлік мемлекеттік университеті, Ақтөбе, Қазақстан

ТӨРТІНШІ РЕТТІ ДЕРБЕС ТУЫНДЫЛЫ ДИФФЕРЕНЦИАЛДЫҚ ТЕҢДЕУЛЕР ЖҮЙЕСІ ҮШІН БАСТАПҚЫ - ШЕТТІК ЕСЕП ТУРАЛЫ

Аннотация. Төртінші ретті дербес туындылы дифференциалдық теңдеулер жүйесі үшін бастапқышеттік есеп қарастырылады. Төртінші ретті дербес туындылы дифференциалдық теңдеулер жүйесі үшін бастапқы-шеттік есептің классикалық шешімінің бар болуы мәселелері мен олардың жуық шешімдерін табу әдістері зерттелген. Төртінші ретті дербес туындылы дифференциалдық теңдеулер жүйесі үшін бастапқышеттік есептің классикалық шешімінің бар болуы мен жалғыздығының жеткілікті шарттары тағайындалған. Жаңа белгісіз функциялар енгізу жолымен зерттеліп отырған есеп гиперболалық теңдеулер жүйесі үшін параметрлері бар бейлокал есептен және интегралдық қатынастардан тұратын пара-пар есепке келтірілген. Зерттеліп отырған есептің жуық шешімін табу алгоритмдері ұсынылған және олардың жинақтылығы дәлелденген. Параметрлері бар пара-пар есептің жалғыз шешімінің бар болуының жеткілікті шарттары тағайындалған. Төртінші ретті дербес туындылы дифференциалдық теңдеулер жүйесі үшін бастапқы-шеттік

есептің бірмәнді шешілімділігінің шарттары бастапқы берілімдер терминінде алынған. Бастапқы-уақыт бойынша периодты шеттік есеп үшін нәтиже жеке келтірілген.

Түйін сөздер: Төртінші ретті дербес туындылы дифференциалдық теңдеулер жүйесі, бастапқы-шеттік есеп, бейлокал есеп, екінші ретті гиперболалық теңдеулер жүйесі, шешілімділік, алгоритм.

Information about authors:

Assanova Anar Turmaganbetkyzy - the member of Editorial Board of journal "News of the NAS RK. Physico-Mathematical Series", Institute of Mathematics and Mathematical Modeling, Chief scientific researcher, Doctor of Physical and Mathematical Sciences, professor, anarasanova@list.ru, assanova@math.kz,

https://orcid.org/0000-0001-8697-8920;

Boichuk Alexander (Oleksandr) Andriyovych – Institute of Mathematics of the National Academy Sciences of the Ukraine, Corresponding member of the National Academy Sciences of the Ukraine, Head of laboratory of boundary value problems of differential equations theory, Doctor of Physical and Mathematical Sciences, professor, boichuk@imath.kiev.ua, boichuk@imath.kiev.ua, boichuk.aa@gmail.com, boichuk.aa@gmailto:boichuk.aa@gmailto:boichuk.aa@gmailto:boichuk.aa@gmailto:boichuk.aa@gmailto:boichuk.aa@gmailto:boichuk.aa@gmailto:b

Tokmurzin Zhanibek Syrlybayevich – K.Zhubanov Aktobe Regional State University, PhD -student, tokmurzinzh@gmail.com, https://orcid.org/0000-0002-3738-5923;

REFERENCES

- [1] Ptashnyck B.I. Ill-posed boundary value problems for partial differential equations, Naukova Dumka, Kiev (1984) (in Russian).
- [2] Ptashnyk B.Yo., Il'kiv V.S., Kmit' I. Ya., Polishchuk V.M. Nonlocal boundary value problems for partial differential equations, Naukova Dumka: Kyiv, Ukraine, 2002. (in Ukrainian)
- [3] Kiguradze T., Lakshmikantham V. On the Dirichlet problem for fourth order linear hyperbo-lic equations // Nonlinear Analysis, 49 (2002), No. 2, 197–219.
- [4] Kiguradze T., Lakshmikantham V. On Dirichlet problem in a characteristic rectangle for higher order linear hyperbolic equations // Nonlinear Anal., **50**:8 (**2002**), 1153—1178.
 - PII: S0362-546X(01)00806-9
- [5] Kiguradze T.I., Kusano T. Well-posedness of initial-boundary value problems for higher-order linear hyperbolic equations with two independent variables // Differential Equations, 39:4 (2003), 553--563.
- [6] Kiguradze T., Kusano T. On ill-posed initial-boundary value problems for higher order linear hyperbolic equations with two independent variables // Differential Equations, **39**:10 (**2003**), 1379–1394.
- [7] Dzhokhadze O.M. The Riemann function for higher-order hyperbolic equations and systems with dominated lower terms, Differential Equations, **39**:10 (**2003**), 1440–1453.

DOI: 0012-2661/03/3910-1440

- [8] Midodashvili B. Generalized Goursat problem for a spatial fourth order hyperbolic equation with dominated low terms // Proc. of A. Razmadze Math. Institute, **138** (2005), 43--54.
 - [9] Nakhushev A.M. Problems with shift for a partial differential equations, Nauka, Moskow (2006). ISBN: 5-02-034076-6
- [10] Kiguradze I., Kiguradze T. On solvability of boundary value problems for higher order nonlinear hyperbolic equations // Nonlinear Analysis, **69 (2008)**, 1914--1933.

doi:10.1016/j.na.2007.07.033

- [11] Kiguradze T. On solvability and well-posedness of boundary value problems for nonlinear hyperbolic equations of the fourth order // Georgian Mathematical Journal, 15 (2008), No. 3, pp. 555–569.
- [12] Kiguradze T. The Valle-Poussin problem for higher order nonlinear hyperbolic equations // Computers & Mathematics with Applications, **59** (**2010**), 994--1002.

doi:10.1016/j.camwa.2009.09.009

- [13] Asanova A.T. On the unique solvability of a nonlocal boundary value problem with data on intersecting lines for systems of hyperbolic equations, Differential Equations. **45**:3 (2009), 385--394. DOI: 10.1134/S0012266109030082
- [14] Asanova A.T. On a boundary-value problem with data on noncharacteristic intersecting lines for systems of hyperbolic equations with mixed derivative, Journal of Mathematical Sciences (United States). **187**:4 (**2012**), 375-386. 1072-3374/12/1874–0375
- [15] Asanova A.T. On a nonlocal boundary-value problem for systems of impulsive hyperbolic equations, Ukrainian Mathematical Journal. **65**:3 **(2013)**, 349--365.

0041-5995/13/6503-0349

- [16] Asanova A.T., Dzhumabaev D.S. Well-posedness of nonlocal boundary value problems with integral condition for the system of hyperbolic equations, Journal of Mathematical Analysis and Applications, **402**:1 (**2013**), 167--178. doi:10.1016/j.jmaa.2013.01.012
- [17] Asanova A.T. Well-posed solvability of a nonlocal boundary-value problem for systems of hyperbolic equations with impulse effects, Ukrainian Mathematical Journal. 67:3 (2015), 333-346. DOI: 10.1007/s11253-015-1083-3
- [18] Asanova A.T. On solvability of nonlinear boundary value problems with integral condition for the system of hyperbolic equations, Electronic Journal of Qualitative Theory of Differential Equations. 63 (2015), 1--13. doi: 10.14232/ejqtde.2015.1.63

- [19] Asanova A.T., Imanchiev A.E. On conditions of the solvability of nonlocal multi-point boundary value problems for quasi-linear systems of hyperbolic equations, Eurasian Mathematical Journal. 6:4 (2015), 19--28.
- [20] Asanova A.T. Multipoint problem for a system of hyperbolic equations with mixed derivative, Journal of Mathematical Sciences (United States), 212:3 (2016), 213--233.

DOI: 10.1007/s10958-015-2660-6

[21] Asanova A.T. Criteria of solvability of nonlocal boundary-value problem for systems of hyperbolic equations with mixed derivatives, Russian Mathematics. 60:1 (2016), 1-17.

DOI: 10.3103/S1066369X16050017

- [22] Assanova A.T. On the solvability of nonlocal boundary value problem for the systems of impulsive hyperbolic equations with mixed derivatives, Journal of Discontinuity, Nonlinearity and Complexity. 5:2 (2016), 153--165. DOI: 10.5890/DNC.2016.06.005
- [23] Assanova A.T. Periodic solutions in the plane of systems of second-order hyperbolic equations, Mathematical Notes. **101**:1 (**2017**), 39–47. DOI: 10.1134/S0001434617010047
- [24] Assanova A.T. Nonlocal problem with integral conditions for a system of hyperbolic equations in characteristic rectangle, Russian Mathematics. 61:5 (2017), 7–20.

DOI: 10.3103/S1066369X17050024

- [25] Asanova A.T., Kadirbaeva Zh. M., and Bakirova E. A. On the unique solvability of a nonlocal boundary-value problem for systems of loaded hyperbolic equations with impulsive actions, Ukrainian Mathematical Journal, **69**:8 (**2018**), 1175-1195.
 - DOI: 10.1007/s11253-017-1424-5
- [26] Assanova A.T. On a nonlocal problem with integral conditions for the system of hyperbolic equations // Differential Equations, **54**:2 (**2018**), 201–214.

DOI: 10.1134/S0012266118020076

- [27] Assanova A.T., Kadirbayeva Z.M. Periodic problem for an impulsive system of the loaded hyperbolic equations // Electronic Journal of Differential Equations, 2018:72 (2018), 1--8.
- [28] Assanova A.T., Imanchiyev A.E., and Kadirbayeva Zh.M. Numerical Solution of Systems of Loaded Ordinary Differential Equations with Multipoint Conditions // Computational Mathematics and Mathematical Physics, **58**:4 (**2018**), 508–516.DOI: 10.1134/S096554251804005X
- [29] Assanova A.T., Kadirbayeva Z.M. On the numerical algorithms of parametrization method for solving a two-point boundary-value problem for impulsive systems of loaded differential equations // Computational and Applied Mathematics, 37:4 (2018), 4966--4976.

DOI 10.1007/s40314-018-0611-9

- [30] Assanova A.T., Sabalakhova A.P., Toleukhanova Z.M. On the solving of initial-boundary value problem for system of partial differential equations of the third order, News of the National Academy of Sciences of the Republic of Kazakhstan. Physico-Mathematical Series. 3:319 (2018), 67 73.
- [31] Assanova A.T., Alikhanova B.Zh., Nazarova K.Zh. Well-posedness of a nonlocal problem with integral conditions for third order system of the partial differential equations, News of the National Academy of Sciences of the Republic of Kazakhstan. Physico-Mathematical Series. 5:321 (2018), 33 41. https://doi.org/10.32014/2018.2518-1726.5

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see http://www.elsevier.com/publishingethics and http://www.elsevier.com/journal-authors/ethics.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see http://www.elsevier.com/postingpolicy), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service http://www.elsevier.com/editors/plagdetect.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

www:nauka-nanrk.kz

http://physics-mathematics.kz/index.php/en/archive

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы М. С. Ахметова, Т.А. Апендиев, Д.С. Аленов Верстка на компьютере А.М. Кульгинбаевой

Подписано в печать 05.02.2019. Формат 60x881/8. Бумага офсетная. Печать – ризограф. $4,75\,$ п.л. Тираж 300. Заказ 1.

Национальная академия наук РК 050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19