

ISSN 2518-1726 (Online),  
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ  
Әль-фараби атындағы Қазақ ұлттық университетінің

# Х А Б А Р Л А Р Ы

---

---

## ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН  
Қазақстан Республикасының  
Ғылым Академиясының  
Әль-Фараби атындағы  
Қазақ ұлттық университетінің

## NEWS

OF THE NATIONAL ACADEMY OF SCIENCES  
OF THE REPUBLIC OF KAZAKHSTAN  
Al-farabi kazakh  
national university

**SERIES**  
**PHYSICO-MATHEMATICAL**

**1 (323)**

**JANUARY – FEBRUARY 2019**

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р ы  
ф.-м.ғ.д., проф., ҚР ҰҒА академигі **Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

**Жұмаділдаев А.С.** проф., академик (Қазақстан)  
**Кальменов Т.Ш.** проф., академик (Қазақстан)  
**Жантаев Ж.Ш.** проф., корр.-мүшесі (Қазақстан)  
**Өмірбаев У.У.** проф. корр.-мүшесі (Қазақстан)  
**Жүсіпов М.А.** проф. (Қазақстан)  
**Жұмабаев Д.С.** проф. (Қазақстан)  
**Асанова А.Т.** проф. (Қазақстан)  
**Бошқаев К.А.** PhD докторы (Қазақстан)  
**Сұраған Д.** корр.-мүшесі (Қазақстан)  
**Quevedo Hernando** проф. (Мексика),  
**Джунушалиев В.Д.** проф. (Қырғыстан)  
**Вишневский И.Н.** проф., академик (Украина)  
**Ковалев А.М.** проф., академик (Украина)  
**Михалевич А.А.** проф., академик (Белорус)  
**Пашаев А.** проф., академик (Әзірбайжан)  
**Такибаев Н.Ж.** проф., академик (Қазақстан), бас ред. орынбасары  
**Тигиняну И.** проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)  
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде  
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.  
Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор  
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Редакционная коллегия:

**Джумадильдаев А.С.** проф., академик (Казахстан)  
**Кальменов Т.Ш.** проф., академик (Казахстан)  
**Жантаев Ж.Ш.** проф., чл.-корр. (Казахстан)  
**Умирбаев У.У.** проф. чл.-корр. (Казахстан)  
**Жусупов М.А.** проф. (Казахстан)  
**Джумабаев Д.С.** проф. (Казахстан)  
**Асанова А.Т.** проф. (Казахстан)  
**Бошкаев К.А.** доктор PhD (Казахстан)  
**Сураган Д.** чл.-корр. (Казахстан)  
**Quevedo Hernando** проф. (Мексика),  
**Джунушалиев В.Д.** проф. (Кыргызстан)  
**Вишневский И.Н.** проф., академик (Украина)  
**Ковалев А.М.** проф., академик (Украина)  
**Михалевич А.А.** проф., академик (Беларусь)  
**Пашаев А.** проф., академик (Азербайджан)  
**Такибаев Н.Ж.** проф., академик (Казахстан), зам. гл. ред.  
**Тигиняну И.** проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов  
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

E d i t o r i n c h i e f  
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d:

**Dzhumadildayev A.S.** prof., academician (Kazakhstan)  
**Kalmenov T.Sh.** prof., academician (Kazakhstan)  
**Zhantayev Zh.Sh.** prof., corr. member. (Kazakhstan)  
**Umirbayev U.U.** prof. corr. member. (Kazakhstan)  
**Zhusupov M.A.** prof. (Kazakhstan)  
**Dzhumabayev D.S.** prof. (Kazakhstan)  
**Asanova A.T.** prof. (Kazakhstan)  
**Boshkayev K.A.** PhD (Kazakhstan)  
**Suragan D.** corr. member. (Kazakhstan)  
**Quevedo Hernando** prof. (Mexico),  
**Dzhunushaliyev V.D.** prof. (Kyrgyzstan)  
**Vishnevskiy I.N.** prof., academician (Ukraine)  
**Kovalev A.M.** prof., academician (Ukraine)  
**Mikhalevich A.A.** prof., academician (Belarus)  
**Pashayev A.** prof., academician (Azerbaijan)  
**Takibayev N.Zh.** prof., academician (Kazakhstan), deputy editor in chief.  
**Tiginyanu I.** prof., academician (Moldova)

**News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.**

**ISSN 2518-1726 (Online), ISSN 1991-346X (Print)**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN  
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.5>

Volume 1, Number 323 (2019), 38 – 45

UDC 517.946:517.588

IRSTI 27.29.21; 27.23.25

Zh.N. Tasmambetov<sup>1</sup>, N. Rajabov<sup>2</sup>, A.A. Issenova<sup>3</sup>

<sup>1</sup> Aqtobe Regional Zhubanov State University, <sup>2</sup> Tajik National University,

<sup>3</sup> Aqtobe Regional Zhubanov State University  
tasmam@rambler.ru, nusrat38@mail.ru, akkenje\_ia@mail.ru

## THE CONSTRUCTION OF A SOLUTION OF A RELATED SYSTEM OF THE LAGUERRE TYPE

**Abstract.** The aim of the work is to study the system of Laguerre type obtained from the degenerate of Horn system by direct selection of parameters, as well as using an exponential transformation. Such a system consisting of two partial differential equations of second order is called related to the basic Laguerre system. The difficulties of studying are that if in the ordinary case there is one degenerate of Kummer's equation and only one degenerate hypergeometric function satisfying it, then in the case of two variables there 20 degenerate systems and 20 degenerate hypergeometric functions of two variables satisfying them appear. It is not known how many systems of Laguerre type exist, and with which of the 20 degenerate systems it links to. There is no general method of a research In this work Frobenius-Latysheva's method which is generalized in this case by Zh.N. Tasmambetov is applied to creation of their normal and regular solution depending on Laguerre's polynomial of two variables. The classification of singular curves using rank and antirank is given, as well as basic information about the features of constructing normal-regular solutions of such systems. The main theorem on the existence of four linearly independent partial solutions is proved. Solutions are determined by the degenerate hypergeometric function of M.P. Humbert in the form of normal-regular ranks of two variables depending on Laguerre's polynomials. The conclusions indicate the relationship of such systems with overridden systems and some representations of Laguerre's polynomial of two variables.

**Key words:** Related, system, Laguerre-type system, Horn system, normal-regular solution, special curves, rank, antirank, overdetermined.

### Introduction

The degenerate hypergeometric function is the root of many well-known functions, and through it all orthogonal polynomials of one variable are expressed [1]-[2]. Indeed, if  $\gamma$  is not an integer, is  $\alpha$  a negative integer or zero, then the series

$$G(\alpha, \gamma; x) = 1 + \frac{\alpha}{\gamma} x + \frac{\alpha(\alpha+1)}{2!\gamma(\gamma+1)} x^2 + \dots \quad (1)$$

representing the degenerate hypergeometric function terminates, and we obtain a polynomial  $G(-n, \gamma; x)$  in particular expressing the polynomial of Laguerre. In the theory of orthogonal polynomials, there are several differential equations solutions of which are Laguerre's polynomials and various applications in the problems of mathematical physics, as well as in the theory of the hydrogen atom, etc. [3, c.226]- [4]- [5,115-118]. The generalization of this theory to Laguerre's polynomials of two variables and systems of partial differential equations of the second order, which they satisfy, has not reached this level. The study is complicated by the fact that if in the ordinary case there is only one degenerate hypergeometric equation, then in the case of two variables there are 20 degenerate systems and 20 degenerate hypergeometric functions of two variables satisfying them [6, c.226-230]-[7]. It is not yet known how

many systems of the Laguerre type are there and with which of the 20 degenerate systems they are connected. Apparently, this was influenced by the insufficient development of the analytical theory of such systems. Therefore, another direction for studying orthogonal polynomials of two variables as eigenfunctions of linear partial differential operators of the second order was developed [8]-[9]-[10].

In the works of Zh.N.Tasmambetov and A.A.Issanova the system of Horn ( $\Psi_2$ ) was selected as a binding system and the connection between the degenerate hypergeometric function of Humbert  $\Psi_2(\alpha, \gamma, \gamma'; x, y)$  and the Laguerre's function of two variables  $L_{n,m}^{(\alpha, \beta)}(x, y)$  was studied.

In [11] it was shown that from the system of Horn

$$\left. \begin{aligned} xZ_{xx} + (\gamma - x)Z_x - yZ_y + nZ &= 0 \\ yZ_{yy} + (\gamma - y)Z_y - xZ_x + nZ &= 0 \end{aligned} \right\} \quad (2)$$

when  $\gamma = \alpha + 1, \gamma' = \beta + 1, \alpha \neq 0, \beta \neq 0, \lambda = -n$  the basic system of Laguerre is obtained

$$\left. \begin{aligned} xZ_{xx} + (\alpha + 1 - x)Z_x - yZ_y + nZ &= 0 \\ yZ_{yy} + (\beta + 1 - y)Z_y - xZ_x + nZ &= 0 \end{aligned} \right\} \quad (3)$$

solution of which is a polynomial

$$\Psi_2(-n, \alpha + 1, \beta + 1; x, y) = \sum_{\mu, \nu=0}^{\infty} \frac{(-n)_{\mu+\nu}}{(\alpha + 1)_{\mu} (\beta + 1)_{\nu}} \cdot \frac{x^{\mu}}{\mu!} \cdot \frac{y^{\nu}}{\nu!}. \quad (4)$$

By analogy, this polynomial is called the generalized Laguerre's polynomials of two variables and is denoted by

$$L_{n,m}^{(\alpha, \beta)} = \Psi_2(-n, \alpha + 1, \beta + 1; x, y). \quad (5)$$

### Basic information

According to the general theory of systems of the form (2), when the condition of compatibility and integrality is performed [12], it has up to four linearly independent solutions  $Z_i (i = \overline{1,4})$ , and the general solution depends on arbitrary constants and is represented as a sum

$$Z(x, y) = C_1 Z_1(x, y) + C_2 Z_2(x, y) + C_3 Z_3(x, y) + C_4 Z_4(x, y) \quad (6)$$

where  $C_i (i = \overline{1,4})$  are arbitrary constants,  $Z = Z(x, y)$  is a general unknown.

The system has a regular  $(0,0)$  singularity and an irregular  $(\infty, \infty)$  singularity. To classify regular and irregular singularities K.Ya. Latysheva used the notion of rank

$$p = 1 + k, k = \max_{(1 \leq s \leq n)} \frac{\beta_s - \beta_0}{s} \quad (7)$$

introduced by H. Poincare and antirank

$$m = -1 - \chi, \chi = \min_{(1 \leq s \leq n)} \frac{\pi_s - \pi_0}{s} \quad (8)$$

introduced by L. Tome.

These concepts were generalized to the case of the studied system of (2) Zh.N.Tasmambetov[13].

If the rank is  $p \leq 0$ , then the special curve  $(x = \infty, y = \infty)$  is regular, when  $p > 0$  the special curve is irregular. When  $m \leq 0$  a special curve  $(0,0)$  is regular, and if  $m > 0$  special is irregular [13].

**Definition 1.** If the rank  $p > 0$  and antirank  $m \leq 0$ , then system (2) has a solution

$$Z(x, y) = \exp Q(x, y) \cdot x^{\rho_i} y^{\sigma_i} \sum_{\mu, \nu=0}^{\infty} A_{\mu, \nu}^{(i)} x^{\mu} y^{\nu}, A_{0,0} \neq 0, \quad (9)$$

where  $\rho_i, \sigma_i (i = \overline{1,4}), A_{\mu, \nu}^{(i)} (\mu, \nu = 0, 1, 2, \dots)$  - unknown constants;  $Q(x, y)$  - polynomial of two variables

$$Q(x, y) = \frac{\alpha_{p0}}{p} x^p + \frac{\alpha_{0p}}{p} y^p + \dots + \alpha_{11} xy + \alpha_{10} x + \alpha_{01} y, \quad (10)$$

with unknown coefficients  $\alpha_{p0}, \alpha_{0p}, \dots, \alpha_{11}, \alpha_{10}, \alpha_{01}$ . The solution of the form (9) is called normal-regular.

If the special curve  $(0,0)$  is regular, then the polynomial  $Q(x, y) \equiv 0$  and the solution of the system exists in the form of a generalized power series of two variables

$$Z(x, y) = x^{\rho_i} y^{\sigma_i} \sum_{\mu, \nu=0}^{\infty} A_{\mu, \nu}^{(i)} x^{\mu} y^{\nu}, A_{0,0} \neq 0, \quad (11)$$

where  $\rho_i, \sigma_i (i = \overline{1,4}), A_{\mu, \nu}^{(i)} (\mu, \nu = 0, 1, 2, \dots)$  - the unknown constants.

The highest degree of the polynomial  $Q(x, y)$  is determined by the rank  $p$ .

**Definition 2.** The values of number  $p$  determined by the equality (7) are called series order (9) and can be an integer or a fractional number (positive or negative).

## CONCLUSION OF THE RELATED SYSTEM OF THE LAGUERRE TYPE AND THE CONSTRUCTION OF ITS SOLUTION

### Formulation of the problem

From the system of Horn  $(\Psi_2)$  by means of converting

$$Z = \exp\left(\frac{x}{2} + \frac{y}{2}\right) \cdot x^{\frac{\alpha+1}{2}} y^{\frac{\beta+1}{2}} \cdot U(x, y) \quad (12)$$

a system of Laguerre type is installed

$$\left. \begin{aligned} x^2 U_{xx} - xy U_y + \left(-\frac{x^2}{4} - \frac{xy}{2} + kx + \frac{1}{4} - \alpha^2\right) \cdot U &= 0 \\ y^2 U_{yy} - xy U_x + \left(-\frac{y^2}{4} - \frac{xy}{2} + ky + \frac{1}{4} - \beta^2\right) \cdot U &= 0 \end{aligned} \right\} \quad (13)$$

where  $k = (\alpha + \beta + 2 - 2\lambda)/2$  is related with the basic Laguerre system (3).

Such systems belong to the Whitaker-type system [7]. By applying Frobenius-Latysheva method [13] we want to establish distinctive features of the system (12) and construct its normal-regular solution dependent on Laguerre's polynomials of two variables.

### MAIN RESULTS

**Theorem 1.** The system of second order partial differential equations (13) has four linearly independent partial solutions, which are expressed through the degenerate hypergeometric function of M.R. Humbert  $\Psi_2(\alpha, \gamma, \gamma'; x, y)$  in the form of normal-regular series

$$\begin{aligned} U(x, y) &= \exp\left(-\frac{x}{2} - \frac{y}{2}\right) \cdot x^{\frac{\alpha+1}{2}} y^{\frac{\beta+1}{2}} \cdot \Psi_2(-n, \alpha + 1, \beta + 1; x, y) = \\ &= \exp\left(-\frac{x}{2} - \frac{y}{2}\right) \cdot x^{\frac{\alpha+1}{2}} y^{\frac{\beta+1}{2}} \cdot L_{n,n}^{(\alpha, \beta)}(x, y) \end{aligned} \quad (14)$$

dependent on the Laguerre's polynomial of two variables

$$\begin{aligned}
 L_{n,n}^{(\alpha,\beta)}(x,y) = & 1 - \frac{n}{\Gamma(\alpha+1)}x - \frac{n}{\Gamma(\beta+1)}y + \frac{n(n-1)}{\Gamma(\alpha+1)\Gamma(\beta+1)}xy + \frac{n(n-1)}{\Gamma(\alpha+1)\Gamma(\alpha+2)}x^2 + \\
 & + \frac{n(n-1)}{\Gamma(\beta+1)\Gamma(\beta+2)}y^2 + \dots + (-1)^n \frac{n(n-1)\dots 1}{n!(\alpha+1)\dots(\alpha+n)}x^n + \\
 & + (-1)^n \frac{n(n-1)\dots 1}{n!(\alpha+1)\dots(\alpha+n-1)\Gamma(\beta+1)}x^{n-1}y + \dots + (-1)^n \frac{n(n-1)\dots 1}{n!(\beta+1)\dots(\beta+n)}y^n
 \end{aligned}
 \tag{15}$$

*Evidence.* For the proof Frobenius-Latysheva method is used. Like the degenerate system (2) the system (12) has a regular (0,0) singularity and an irregular (∞,∞) singularity. By highest degrees of independent variables  $x$  and  $y$  certain subranks:  $k'_1 = 0, k''_1 = 0$  and rank  $p = 1 + k = 1$ . Then according to the method of Frobenius-Latysheva for the construction of normal-regular solution of (9), in the system (13) the transformation is correct:

$$U = \exp(\alpha_{10}x + \alpha_{01}y)\Phi(x,y) \tag{16}$$

where  $\alpha_{10}$  and  $\alpha_{01}$  are uncertain coefficients, which need to be determined from the newly obtained support system.

$$\left. \begin{aligned}
 x^2\Phi_{xx} + 2\alpha_{10}^2x^2\Phi_x - xy\Phi_y + \left( \left( \alpha_{10}^2 - \frac{1}{4} \right)x^2 - \left( \alpha_{01}^2 - \frac{1}{2} \right)xy + kx + \frac{1}{4} - \frac{\alpha^2}{4} \right) \cdot \Phi &= 0 \\
 y^2\Phi_{yy} + 2\alpha_{01}^2y^2\Phi_y - xy\Phi_x + \left( \left( \alpha_{01}^2 - \frac{1}{4} \right)y^2 - \left( \alpha_{10}^2 - \frac{1}{2} \right)xy + ky + \frac{1}{4} - \frac{\beta^2}{4} \right) \cdot \Phi &= 0
 \end{aligned} \right\} \tag{17}$$

By equating coefficients to zero at the highest degrees of independent variables  $x^2$  and  $y^2$  at unknown  $\Phi(x,y)$ , we define a system of characteristic equations

$$\begin{aligned}
 b_{10}^{(1)}(\alpha_{10}, \alpha_{01}) &= \alpha_{10}^2 - \frac{1}{4} = 0, \\
 b_{01}^{(2)}(\alpha_{10}, \alpha_{01}) &= \alpha_{01}^2 - \frac{1}{4} = 0.
 \end{aligned}
 \tag{18}$$

This shows the fulfillment of the first necessary condition for the existence of a normal-regular solution (9) connected with the definition of the unknown coefficients of the  $Q(x,y)$  polynomial [13].

**Theorem 2.** Equality (18) is required for a supporting system to have at least one solution of the form (9).

The system (17) has four root pairs:

$$\begin{aligned}
 (\alpha_{10}^{(1)}, \alpha_{01}^{(1)}) &= \left( \frac{1}{2}, \frac{1}{2} \right), (\alpha_{10}^{(1)}, \alpha_{01}^{(2)}) = \left( \frac{1}{2}, -\frac{1}{2} \right), \\
 (\alpha_{10}^{(2)}, \alpha_{01}^{(1)}) &= \left( -\frac{1}{2}, \frac{1}{2} \right), (\alpha_{10}^{(2)}, \alpha_{01}^{(2)}) = \left( -\frac{1}{2}, -\frac{1}{2} \right),
 \end{aligned}
 \tag{19}$$

defining four polynomials of the first degree of the form (10), since the rank of the system is equal to one:

$$Q_i(x,y) = \alpha_{10}^{(i)}x + \alpha_{01}^{(i)}y, i = \overline{1,4}.$$

Four  $(\alpha_{10}^{(l)}, \alpha_{01}^{(l)})$ ,  $l = 1,2$  pairs in (18) define four systems from the auxiliary system (18). Each of them can have up to four linearly independent particular solutions. Thus, the initial system should have up to 16 private solutions. However, a detailed study shows that out of the four systems, only the system



$$\left. \begin{aligned} x^2 \Phi_{xx} + x^2 \Phi_x - xy \Phi_y + \left( kx + \frac{1}{4} - \frac{\alpha^2}{4} \right) \cdot \Phi &= 0 \\ y^2 \Phi_{yy} + y^2 \Phi_y - xy \Phi_x + \left( ky + \frac{1}{4} - \frac{\beta^2}{4} \right) \cdot \Phi &= 0 \end{aligned} \right\} \quad (20)$$

has four linearly independent particular solutions. All of them are expressed through degenerate hypergeometric function of Humbert  $\Psi_2(\alpha, \gamma, \gamma'; x, y)$ .

Indeed, by making up the system of characteristic functions of the system (20) we make sure that the system of defining equations with respect to the singularity  $(0,0)$

$$\left. \begin{aligned} f_{00}^{(1)}(\rho, \sigma) &= \rho(\rho - 1) + \frac{1}{4} - \frac{\alpha^2}{4} = 0, \\ f_{00}^{(2)}(\rho, \sigma) &= \sigma(\sigma - 1) + \frac{1}{4} - \frac{\beta^2}{4} = 0, \end{aligned} \right\} \quad (21)$$

has four pairs of roots:

$$\left. \begin{aligned} (\rho_1, \sigma_1) &= \left( \frac{1}{2} + \frac{\alpha}{2}, \frac{1}{2} + \frac{\beta}{2} \right), (\rho_1, \sigma_2) = \left( \frac{1}{2} + \frac{\alpha}{2}, \frac{1}{2} - \frac{\beta}{2} \right), \\ (\rho_2, \sigma_1) &= \left( \frac{1}{2} - \frac{\alpha}{2}, \frac{1}{2} + \frac{\beta}{2} \right), (\rho_2, \sigma_2) = \left( \frac{1}{2} - \frac{\alpha}{2}, \frac{1}{2} - \frac{\beta}{2} \right). \end{aligned} \right\} \quad (22)$$

This shows the fulfillment of the second necessary condition.

**Theorem 3.** In order for the system (13) to have a normal-regular solution of the form (9), it is necessary for the pair  $(\rho, \sigma)$  to be the root of the defining equations with respect to the  $(0,0)$  singularity of the form (20) obtained from the auxiliary system (17) by substituting instead of the unknown  $Z(x, y) = x^\rho \cdot y^\sigma$ .

The existence of four pairs of roots (22) ensures the existence of four linearly independent particular solutions of the system (20) in the form of generalized power series (12). Since, the values of  $\rho$  and  $\sigma$  are defined, it remains to find the unknown constants  $A_{\mu, \nu}(\mu, \nu = 0, 1, 2, \dots)$  with the help of a system of recurrent sequences

$$\sum A_{m-\mu, n-\nu}^{(i)} \cdot f_{\mu, \nu}^{(i)}(\rho + m - \mu, \sigma + n - \nu), (m, n = 0, 1, 2, \dots; i = 1, 2; \mu, \nu = 0, 1, 2, \dots).$$

Thus, the constructed particular solutions of (19) have the form of the

$$\begin{aligned} \Phi_1(x, y) &= x^{\frac{\alpha+1}{2}} y^{\frac{\beta+1}{2}} \cdot \Psi_2\left(\frac{\alpha+\beta}{2} + 1 - k, \alpha + 1, \beta + 1; x, y\right) \\ \Phi_2(x, y) &= x^{\frac{\alpha+1}{2}} y^{\frac{1-\beta}{2}} \cdot \Psi_2\left(\frac{\alpha-\beta}{2} + 1 - k, \alpha + 1, 1 - \beta; x, y\right) \\ \Phi_3(x, y) &= x^{\frac{1-\alpha}{2}} y^{\frac{\beta+1}{2}} \cdot \Psi_2\left(\frac{\beta-\alpha}{2} + 1 - k, 1 - \alpha, \beta + 1; x, y\right) \\ \Phi_4(x, y) &= x^{\frac{1-\alpha}{2}} y^{\frac{1-\beta}{2}} \cdot \Psi_2\left(\frac{-\alpha-\beta}{2} + 1 - k, 1 - \alpha, 1 - \beta; x, y\right) \end{aligned}$$

Considering  $k = (\alpha + \beta + 2 - 2\lambda)/2$ , these solutions are represented in the form

$$\Phi_1(x, y) = x^{\frac{\alpha+1}{2}} y^{\frac{\beta+1}{2}} \cdot \Psi_2(-n, \alpha + 1, \beta + 1; x, y)$$

$$\begin{aligned}\Phi_2(x, y) &= x^{\frac{\alpha+1}{2}} y^{\frac{1-\beta}{2}} \cdot \Psi_2(-n-\beta, \alpha+1, 1-\beta; x, y) \\ \Phi_3(x, y) &= x^{\frac{1-\alpha}{2}} y^{\frac{\beta+1}{2}} \cdot \Psi_2(-n-\alpha, 1-\alpha, \beta+1; x, y) \\ \Phi_4(x, y) &= x^{\frac{1-\alpha}{2}} y^{\frac{1-\beta}{2}} \cdot \Psi_2(-n-\alpha-\beta, 1-\alpha, 1-\beta; x, y)\end{aligned}\quad (23)$$

Hence, it is not difficult to notice that the system solution we are interested in (20):

$$\Phi_1(x, y) = x^{\frac{\alpha+1}{2}} y^{\frac{\beta+1}{2}} \cdot \Psi_2(-n, \alpha+1, \beta+1; x, y) = x^{\frac{\alpha+1}{2}} y^{\frac{\beta+1}{2}} \cdot L_{n,n}^{(\alpha,\beta)}(x, y) \quad (24)$$

and the remaining solutions will not be considered in the future.

The fulfillment of two necessary conditions ensures the existence of a normal-regular solution (14), dependent on the Laguerre's polynomial of two variables (15). The theorem is proved.

The General solution of the system (12) on the basis of (6), taking into account formulas (23), is presented as

$$\begin{aligned}U(x, y) &= C_1 U_1(x, y) + C_2 U_2(x, y) + U_3 Z_3(x, y) + C_4 U_4(x, y) = \\ &= C_1 \exp\left(-\frac{x}{2} - \frac{y}{2}\right) \Phi_1(x, y) + C_2 \exp\left(-\frac{x}{2} - \frac{y}{2}\right) \Phi_2(x, y) + \\ &+ C_3 \exp\left(-\frac{x}{2} - \frac{y}{2}\right) \Phi_3(x, y) + C_4 \exp\left(-\frac{x}{2} - \frac{y}{2}\right) \Phi_4(x, y),\end{aligned}$$

where  $C_i (i = \overline{1,4})$  - are arbitrary constants.

On the basis of the above reasoning, some statements can be made.

**Theorem 4.** The system (16) with respect to  $\Phi(x, y)$ , obtained from (13) by conversion

$$U(x, y) = \exp Q(x, y) \cdot \Phi(x, y) \quad (25)$$

has the same rank as the system (13).

Indeed, since the rank of the system (12) is equal to one, we present transformations (25) in the form of (15) and obtain a system with respect to  $\Phi(x, y)$ , where the rank is  $p = 1$ . The proof of theorem for the General case is given in the monograph [12].

**Theorem 5.** The system (13) for which  $p > 0$ ,  $m \leq 0$  has normally regular solution (14), which is expressed through the generalized Laguerre's polynomial of two variables and the right-hand side (14) converges near the singularity ( $x = 0, y = 0$ ).

The system (13) is said to be related the system of the Laguerre type. As we have seen, its solutions are also expressed through the degenerate hypergeometric function of M.R. Humbert  $\Psi_2(\alpha, \gamma, \gamma'; x, y)$  in the form of normal-regular series (14) dependent on the Laguerre's polynomial of two variables (15).

**Conclusion:** Thus, using the transformation (12), we have established the form of a system of Laguerre's type (3) related to the main system. The application of the Frobenius-Latysheva's method allowed us to construct normal-regular solutions of the derived related system (13) near the singularity (0,0). Generalized Laguerre polynomials also have representations through other hypergeometric functions of two variables [14, p.358].

The limit transition formula is fair [15]

$$\lim_{\alpha \rightarrow \infty} L_{m,n}^{(\alpha,\beta,\gamma)}\left(\frac{x}{\alpha}, \frac{y}{\alpha}\right) = L_m^{(\beta-1)}(x) \cdot L_n^{(\gamma-1)}(y).$$

Formula (24) can be similarly represented using the same limit transition as a product of Laguerre polynomials in variables  $x$  and  $y$ .

In the work [16, p. 6-17] the connection of considered systems with the overdetermined systems, studied in the works of Tajik Mathematical School [17] - [18] - [19] was indicated.

The research in this work can be extended to the case of three variables. The connection of the generalized Laguerre's polynomials of one and two variables with generalized hypergeometric functions [20], [21] of many variables was considered in [14] - [15], [22]. However, for this case the main theorem 1 and theorems 2-5 presented here haven't been proved yet. Also, the question of the computational application of special functions of several variables, as in the monograph [23] hasn't been touched upon. Following [24], it is necessary to develop a numerical method for calculating the values of the degenerate hypergeometric Humbert  $\Psi_2(\alpha, \gamma, \gamma'; x, y)$  functions through the products of Laguerre polynomials in variables and using our formula (24). The problem of the asymptotic expansion, given in [25], is also important when studying the properties of special functions of several variables. We have obtained an asymptotic expansion near the origin  $(0,0)$ .

Ж.Н.Тасмамбетов<sup>1</sup>, Н.Раджабов<sup>2</sup>, А.А.Исенова<sup>3</sup>

<sup>1</sup>Қ.Жұбанов атындағы АӨМУ, Ақтөбе, Қазақстан;

<sup>2</sup>Тәжік ұлттық университеті, Душанбе, Тәжікстан;

<sup>3</sup>Қ.Жұбанов атындағы АӨМУ, Ақтөбе, Қазақстан

### ЛАГЕРРА ТЕКТЕС ТУЫСТАС ЖҮЙЕНІҢ ШЕШІМДЕРІН ТҮРҒЫЗУ

**Аннотация.** Жұмыстың мақсаты – Горнның туындалған жүйесінен параметрлерді тікелей таңдау және экспоненциал түрлендіру көмегімен алынған Лагерра текті жүйені зерттеу. Мұндай екінші ретті дербес туындылы екі теңдеулерден тұратын дифференциалдық теңдеулер жүйесін біз, Лагерра текті негізгі жүйемен туыстас деп атадық. Аталған жүйелерді зерттеудің қиындығы мынада: егер жай дифференциалдық теңдеулер жағдайында Куммердің бір туындалған теңдеуі бар және оны қанағаттандыратын бір ғана гипергеометриялық туындалған функциясы бар болса, онда екі айнымалы жағдайында 20 туындалған жүйелер пайда болады және оларды қанағаттандыратын 20 туындалған гипергеометриялық функциялар белгілі. Өзірге, Лагерра тектес қанша жүйелер бар екендігі және олардың туындалған жүйелердің қайсысымен байланыста екендігі белгісіз. Жалпыға ортақ зерттеу әдісі жоқ. Ұсынылған жұмыста екі айнымалының Лагерра көпмүшелігіне тәуелді қалыпты-регуляр шешімдер тұрғызу үшін, екі айнымалы жағдайына Ж.Н.Тасмамбетов жалпылаған Фробениус-Латышева әдісі пайдаланылады. Ранг және антиранг түсініктерін пайдаланып, ерекше қисықтардың классификациясы жасалған және мұндай жүйелердің қалыпты-регуляр шешімдерін тұрғызуға қатысты негізгі түсініктер келтірілген. Төрт сызықты-тәуелсіз дербес шешімдердің бар болатындығы туралы негізгі теорема дәлелденген. Ол дербес шешімдер Лагерраның екі айнымалының көпмүшелігіне тәуелді М.П.Гумберттің туындалған гипергеометриялық функциясы арқылы өрнектелген қалыпты-регуляр қатар арқылы анықталады. Қорытындысында, зерттелген жүйенің артығымен анықталған жүйелермен байланысы және екі айнымалының Лагерра көпмүшелігінің кейбір басқаша өрнектелуі келтірілген.

**Түйін сөздер:** туыстас, жүйе, Лагерра текті жүйе, Горн жүйесі, қалыпты-регуляр шешім, ерекше қисықтар, ранг, антиранг, артығымен анықталған.

УДК 517.946:517.588

Ж.Н.Тасмамбетов<sup>1</sup>, Н.Раджабов<sup>2</sup>, А.А.Исенова<sup>3</sup>

<sup>1</sup> АРГУ им. К.Жубанова, Ақтөбе, Қазақстан;

<sup>2</sup> Таджикский национальный университет, Таджикистан, Душанбе;

<sup>3</sup> АРГУ им. К.Жубанова, Ақтөбе, Қазақстан

### ПОСТРОЕНИЯ РЕШЕНИЯ РОДСТВЕННОЙ СИСТЕМЫ ТИПА ЛАГЕРРА

**Аннотация.** Целью работы является изучение системы типа Лагерра, полученной из вырожденной системы Горна непосредственным подбором параметров, а также с помощью экспоненциального преобразования. Такая система, состоящая из двух дифференциальных уравнений в частных производных второго порядка, нами названа родственной с основной системой типа Лагерра. Трудности изучения состоят в том, что если в обыкновенном случае имеет место одно вырожденное уравнение Куммера и только одна вырожденная гипергеометрическая функция, удовлетворяющая ему, то в случае двух переменных появляются 20 вырожденных систем и 20 вырожденных гипергеометрических функций двух переменных удовлетворяющих им. Пока не известно, сколько существуют систем типа Лагерра, и с какими из 20-ти вырожденных систем они связаны. Отсутствует общий метод исследования. В данной работе для построения их нормально-регулярного решения, зависящего от

многочлена Лагерра двух переменных, применен обобщенный на этот случай Ж.Н.Тасмамбетовым метод Фробениуса-Латышевой. Приведена классификация особых кривых с помощью ранга и антиранга, а также основные сведения об особенностях построения нормально-регулярных решений таких систем. Доказана основная теорема о существовании четырех линейно-независимых частных решений, которые определяются через вырожденную гипергеометрическую функцию М.П.Гумберта в виде нормально-регулярных рядов зависящих от многочленов Лагерра двух переменных. В выводах указана связь таких систем с переопределенными системами и некоторыми представлениями многочлена Лагерра двух переменных.

**Ключевые слова:** Родственная, система, система типа Лагерра, система Горна, нормально-регулярное решение, особые кривые, ранг, антиранг, переопределенный

#### Information about authors:

Tasmambetov Zhaksylyk Nuradinovich – Aqtobe Regional Zhubanov State University, Doctor of physical and mathematical sciences, Professor Chair of Mathematics, [tasmam@rambler.ru](mailto:tasmam@rambler.ru);

Rajabov Nusrat – Tajik National University, Professor Chair Mathematical Analysis and Function Theory Tajik National University, Academician Academy of Science the Republic of Tajikistan, [nusrat38@mail.ru](mailto:nusrat38@mail.ru);

Issenova Akkenzhe Altmyshevna – Phd Student of Aqtobe Regional Zhubanov State University, [akkenje\\_ia@mail.ru](mailto:akkenje_ia@mail.ru).

#### REFERENCES

- [1] Witteker ET, Watson GN. A course of Modern Analysis. Part 2: Transcendental functions. Cambridge. At the university Press, **1927**. 515pp.
- [2] Kusnetsov DS. Special functions. Second Edition "High School", Moscow, **1965**. 423pp.
- [3] Jeffries G., Swirls B. Methods of Mathematical physics. Issue 3, Moscow: "Mir", **1970**. 344pp.
- [4] Campe de Feriet J., Campbell J., Pet'о R, Fohell' T. Functions of mathematical physics, Reference Guide GIFML: M.**1963**. 120 pp.
- [5] Komarov IV, Ponomarev LI, SlavyanovS.Yu. Spheroidal and Culon spheroidal functions. M.:Nauka, **1976**. 319pp.
- [6] BeytmenG.,Erdeyi A. Higher Transcendental Functions. Part I. Hypergeometric functions. The functions of the Lezhanders, M.: Science, **1965**. 294 pp.
- [7] Appel P, Kampe de Feriet MJ. Fonctions hypergeometriques et hypersperiques. Paris: Gauthier Villars. **1926**. 434 pp.
- [8] Krall HL, Seffer I.M. Orthogonal polynomials in two variables//№Ann.Mathem.pura ed appl.-**1967**. V,76, №4, p.325-376.
- [9] Engelis G.K. On some two-dimensional analogs of classical orthogonal polynomials//Lat.mat.Yearbook. **1974**. Realease 15, p. 169-202.
- [10] Suetin PK. Orthogonal polynomials in two variables. Gordon and breach Science publishers. Moscow, **1988**. 384pp.
- [11] TasmambetovZh.N. Confluent hypergeometric functions and two variables Laguerre polynomials as a solution of Wilczynski type system // AIP Conference Proceeding 1779, 020137(2016); doi.org /10.1063/1/4959751.
- [12] Wilczynski E.J. Projective differential geometry of Curves and Ruled surfaces. Leipzig: Leubner, **1906**. 120pp.
- [13] TasmambetovZh.N. Construction of normal and normally-regular solutions of special systems of partial equations of second order. IP Zhanadilov S.T., Aktobe.**2015**. 463pp.
- [14] Beniwal P.S. and Saran S. On a two variable analogue of generalized Laguerre polynomials, Proc.Nat.Acad.Sci.Seet. A **55(1985)**, p.358-365.
- [15] Rainville E.D. Special functions. Macmillan Co. New York, **1960**; reprinted by Chelsea Bronx, New York 1971.
- [16] TasmambetovZh.N. On the development of research on special systems of partial differential equations of second order// Materials of the international scientific conference of the "information Technology: Innovations science and education", Aktobe, February of 20-21 in 2015year .-p.6-17.
- [17] Radzhabov N.R, Elsaed Abdel Aal M. Over determined linear System of the second order with singular and super singular lines.
- [18] Mikhaylov L.G. Some Partial differential systems of equations and Partial division of two unknown functions. Dushanbe, Donisk Publ., **1986**,115p.
- [19] Shamsudinov F.M. On an over determined System of the second order differential equations with singular point. Trudy mat.centraimLobachevskogo NI –Kazan' **2014**. T.49-p.335-339.
- [20] Slater LJ. Generalized hypergeometric functions. Cambridge, at the university Press, 1966.274p.
- [21] Bailey W.N. Generalized hypergeometric series. Cambridge, Math. Tractk.32, Cambridge University Press, Cambridge.**1935**.
- [22] Subuhi Khan and Babita Agrawal. Some generating relations associated with multiple hypergeometric series //Riv. Mat. Univ. Parma (7) **5(2006)**.1-10.
- [23] Luke Yedell L. Mathematical functions and their approximations. Academic Press.Jue. New York – San Francisco – London.**1975**.608p.
- [24] Dzhumabaev D.S., Bakirova E.A., Kadirbayeva Zh.M. (**2018**) An algorithm for solving a control problem for a differential Equation with a parameter. News of the National Academy of Sciences of the Republic of Kazakhstan. Series of physic-mathematical sciences. Volume 5, Number 321(**2018**), pp 25-32. <https://doi.org/10.32014/2018.2518-1726.4> ISSN 1991-346X. ISSN 2518-1726 (Online), ISSN 1991-346X (Print).
- [25] Dauylbaev M.K., Atakhan N., Mirzakulova A.E. (**2018**) Asymptotic expansion of Solution of general bvp with initial jumps for higher-order singularly perturbed integro-differential equation. News of the National Academy of Sciences of the Republic of Kazakhstan. Series of physic-mathematical sciences. Volume 6, Number 322(2018), pp 28-36. <https://doi.org/10.32014/2018.2518-1726.14> ISSN 1991-346X.

**Publication Ethics and Publication Malpractice  
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www.nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

**ISSN 2518-1726 (Online), ISSN 1991-346X (Print)**

Редакторы *М. С. Ахметова, Т.А. Апендиев, Д.С. Аленов*  
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 05.02.2019.  
Формат 60x881/8. Бумага офсетная. Печать – ризограф.  
4,75 п.л. Тираж 300. Заказ 1.

---

*Национальная академия наук РК*  
*050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19*