

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Әль-фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
Әль-Фараби атындағы
Қазақ ұлттық университетінің

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-farabi kazakh
national university

SERIES
PHYSICO-MATHEMATICAL

3 (325)

MAY - JUNE 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р ы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі **Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев У.У. проф. корр.-мүшесі (Қазақстан)
Жүсіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошқаев К.А. PhD докторы (Қазақстан)
Сұраған Д. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қырғыстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорус)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.
Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жантаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф. чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

E d i t o r i n c h i e f
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d:

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskiy I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.27>

Volume 3, Number 325 (2019), 77 – 84

UDC 517.62/64, 517.968.7, 517.977.1/5

MRNTI 27.33.19; 27.37.17; 27.41.19

A.T.Assanova^{1,2}, E.A.Bakirova^{1,2,3}, Zh.M.Kadirbayeva^{1,2,4}

¹Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan;

²Institute of Information and Computational Technologies, Almaty, Kazakhstan;

³Kazakh National Women's Teacher Training University, Almaty, Kazakhstan;

⁴International Information Technology University, Almaty, Kazakhstan

assanova@math.kz, bakirova1974@mail.ru, zh.kadirbayeva@gmail.com

NUMERICAL IMPLEMENTATION OF SOLVING A BOUNDARY VALUE PROBLEM FOR A SYSTEM OF LOADED DIFFERENTIAL EQUATIONS WITH PARAMETER

Abstract. A linear two-point boundary value problem for the loaded differential equations with parameter is considered. This problem is investigated by parameterization method. We offer algorithm for solving to boundary value problem for the system of loaded differential equations with parameter. In first, original problem is reduced to equivalent problem consisting the Cauchy problems for system of ordinary differential equations with parameters in subintervals and functional relations with respect to introduced additional parameters. At fixed values of parameters the Cauchy problem for system of ordinary differential equations in subinterval has a unique solution. This solution is represented with fundamental matrix of system. Using these representations we compile a system of linear algebraic equations with respect to parameters. We proposed algorithm for finding of numerical solution to the equivalent problem. This algorithm includes of the numerical solving of the Cauchy problems for system of the ordinary differential equations and solving of the linear system of algebraic equations. For numerical solving of the Cauchy problem we apply the Runge–Kutta method of 4th order. The proposed numerical implementation is illustrated by example.

Key words: boundary value problem with parameter, loaded differential equation, numerical method, algorithm.

Funding: The work was supported by grants (№№ AP05132486, AP05131220, AP05132455) of Ministry of Education and Science of the Republic of Kazakhstan (2018-2020).

As well-known, the problem of constructing effective models finds its solution in many areas of science and technology. The active development of computer technology in recent decades, the emergence of new software tools designed to automate professional activity, has significantly affected the methods for solving the problems of identification of parameters. The application of software tools specialized in the field of scientific, technical and engineering calculations provides an opportunity for a deeper study of the investigated area, transferring the main burden of solving the problems from the development, debugging of algorithms and programs to the study of qualitative and numerical characteristics of the problem. Therefore, a modern approach in the theory of control and identification of parameters should be directed to the development of new constructive methods and modifications of known methods for solving boundary value problems for loaded differential equations with parameters. The theory of boundary value problems for the loaded differential equations with parameters is rapidly developing and is used in various fields of applied mathematics, biophysics, biomedicine, chemistry, etc. [1-10]. In spite of this, the questions of finding the effective criteria of unique solvability and constructing the numerical algorithms for finding the solutions of boundary value problems for the system of loaded differential equations with parameters still remain open. One of the constructive

methods for investigating and solving the boundary value problems with parameters for the system of ordinary differential equations is the parameterization method [11]. The parameterization method was developed for the investigating and solving the boundary value problems for the system of ordinary differential equations. On the basis of this method, coefficient criteria for the unique solvability of linear boundary value problems for the system of ordinary differential equations were obtained. Algorithms for finding the approximate solutions were also proposed and their convergence to the exact solution of the problem studied was established. Later, the parameterization method was developed for the two-point boundary value problems for the Fredholm integro-differential equations [12-16]. Necessary and sufficient conditions for the solvability and unique solvability are established, the algorithms for finding the approximate solutions of the problems considered are constructed. In [17], methods for investigating and solving the linear boundary value problems for the linear Fredholm integro-differential equation on the basis of new algorithms of parameterization method are constructed. In [18] this algorithm is used for solve boundary value problem for system of ordinary differential equations with parameter.

In present paper the proposed new algorithms of parameterization method are extended to boundary value problem for loaded differential equations with parameters. We offer the numerical implementation of these algorithms to solve boundary value problem for the loaded differential equations with parameters.

So, we consider the linear boundary value problem for the loaded differential equations with parameter

$$\frac{dx}{dt} = A(t)x + \sum_{j=0}^N K_j(t)x(\theta_j) + A_0(t)\mu + f(t), \quad x \in R^n, \quad \mu \in R^m, \quad t \in (0, T), \quad (1)$$

$$Bx(0) + Cx(T) + B_0\mu = d, \quad d \in R^{n+m}, \quad (2)$$

where the $(n \times n)$ -matrices $A(t)$, $K_j(t)$, $j = \overline{0, N}$, $(n \times m)$ -matrix $A_0(t)$, and n -vector-function $f(t)$ are continuous on $[0, T]$, the $((n + m) \times n)$ - matrices B , C , the $((n + m) \times m)$ - matrix B_0 are constants.

Let $C([0, T], R^n)$ denote the space of continuous functions $x: [0, T] \rightarrow R^n$ with the norm $\|x\|_1 = \max_{t \in [0, T]} \|x(t)\|$. A solution to problem (1), (2) is a pair $(\mu^*, x^*(t))$, with $x^*(t) \in C([0, T], R^n)$, $\mu^*(t) \in R^m$, where the function $x^*(t)$ is continuously differentiable on $(0, T)$ and satisfies the loaded differential equation (1) and boundary condition (2) with $\mu = \mu^*$.

Given the points: $\theta_0 = 0 < \theta_1 < \theta_2 < \dots < \theta_{N-1} < \theta_N = T$, and let $\Delta_N(\theta)$ be the partition of interval $[0, T]$ into N subintervals: $[0, T] = \bigcup_{r=1}^N [\theta_{r-1}, \theta_r)$.

By $C([0, T], \Delta_N, R^{nN})$ we denote the space of function systems $x[t] = (x_1(t), x_2(t), \dots, x_N(t))$, where $x_r: [\theta_{r-1}, \theta_r) \rightarrow R^n$ are continuous and have finite left-hand limits $\lim_{t \rightarrow \theta_r - 0} x_r(t)$ for all $r = \overline{1, N}$, with the norm $\|x\|_2 = \max_{r=1, N} \sup_{t \in [\theta_{r-1}, \theta_r)} \|x_r(t)\|$.

Denote by $x_r(t)$ the restriction of function $x(t)$ to the r -th interval $[\theta_{r-1}, \theta_r)$ and reduce problem (1), (2) to the equivalent multipoint boundary value problem with parameter for the loaded differential equations

$$\frac{dx_r}{dt} = A(t)x_r + \sum_{j=0}^{N-1} K_j(t)x_{j+1}(\theta_j) + K_N(t)x(\theta_N) + A_0(t)\mu + f(t), \quad (3)$$

$$t \in (\theta_{r-1}, \theta_r), r = \overline{1, N},$$

$$Bx_1(0) + Cx_N(T) + B_0\mu = d, \quad (4)$$

$$\lim_{t \rightarrow \theta_s - 0} x_s(t) = x_{s+1}(\theta_s), \quad s = \overline{1, N-1}, \quad (5)$$

$$\lim_{t \rightarrow \theta_N - 0} x_N(t) = x(\theta_N), \quad (6)$$

where (5), (6) are conditions for matching the solution at the interior points of partition $\Delta_N(\theta)$ and at the point $t = \theta_N$.

The solution of problem (3) - (6) is the triple $(\mu^*, x^*(\theta_N), x^*[t])$ with elements $\mu^* \in R^m$, $x^*(\theta_N) \in R^n$, $x^*[t] = (x_1^*(t), x_2^*(t), \dots, x_N^*(t)) \in C([0, T], \Delta_N, R^{nN})$, where functions $x_r^*(t)$, $r = \overline{1, N}$, are continuously differentiable on $[\theta_{r-1}, \theta_r)$, which satisfies system of loaded differential equations (3) and boundary condition (4) with $\mu = \mu^*$ and continuity conditions (5), (6).

We introduce additional parameters $\lambda_r = x_r(\theta_{r-1})$, $r = \overline{1, N}$, and $\lambda_{N+1} = x(\theta_N)$, $\lambda_{N+2} = \mu$. Making the substitution $u_r(t) = x_r(t) - \lambda_r$, on every r -th interval $[\theta_{r-1}, \theta_r)$, $r = \overline{1, N}$, we obtain multipoint boundary value problem with parameters

$$\frac{du_r}{dt} = A(t)(u_r + \lambda_r) + \sum_{j=0}^N K_j(t)\lambda_{j+1} + A_0(t)\lambda_{N+2} + f(t), \quad t \in [\theta_{r-1}, \theta_r), \quad p = \overline{1, N}, \quad (7)$$

$$u_r(\theta_{r-1}) = 0, \quad p = \overline{1, N}, \quad (8)$$

$$B\lambda_1 + C\lambda_{N+1} + B_0\lambda_{N+2} = d, \quad (9)$$

$$\lambda_s + \lim_{t \rightarrow \theta_s - 0} u_s(t) = \lambda_{s+1}, \quad s = \overline{1, N}. \quad (10)$$

A pair $(\lambda^*, u^*[t])$ with elements $\lambda^* = (\lambda_1^*, \lambda_2^*, \dots, \lambda_{N+1}^*, \lambda_{N+2}^*) \in R^{nN+m}$, $u^*[t] = (u_1^*(t), u_2^*(t), \dots, u_N^*(t)) \in C([0, T], \Delta_N, R^{nN})$, is said to be a solution to problem (7)-(10) if the functions $u_r^*(t)$, $r = \overline{1, N}$ are continuously differentiable on $[\theta_{r-1}, \theta_r)$, and satisfy (7) and additional conditions (9), (10) with $\lambda_j = \lambda_j^*$, $j = \overline{1, N+2}$, and initial conditions (8).

Problem (1), (2) is equivalent to problem (7)-(10) in the following sense. If a pair $(\lambda^*, u^*[t])$ is a solution to problem (7)-(10), then the pair $(x^*[t], \mu^*)$ with function $x^*(t)$ defined by the equalities $x^*(t) = \lambda_r^* + u_r^*(t)$, $t \in [\theta_{r-1}, \theta_r)$, $r = \overline{1, N}$, $x^*(T) = \lambda_{N+1}^*$, $\mu^* = \lambda_{N+2}^*$, is a solution to problem (1), (2). Conversely, if a pair $(\tilde{x}(t), \tilde{\mu})$ is a solution to problem (1), (2) and $\tilde{\lambda}_r = \tilde{x}(\theta_{r-1})$, $\tilde{\lambda}_{N+1} = \tilde{x}(\theta_N)$, $\tilde{\lambda}_{N+2} = \tilde{\mu}$, $\tilde{u}_r(t) = \tilde{x}(t) - \tilde{x}(\theta_{r-1})$, $t \in [\theta_{r-1}, \theta_r)$, $r = \overline{1, N}$, then the pair $(\tilde{\lambda}, \tilde{u}[t])$ with $\tilde{\lambda} = (\tilde{\lambda}_1, \tilde{\lambda}_2, \dots, \tilde{\lambda}_{N+1}, \tilde{\lambda}_{N+2}) \in R^{nN+m}$, and $\tilde{u}[t] = (\tilde{u}_1(t), \tilde{u}_2(t), \dots, \tilde{u}_N(t))$, is a solution to problem (7)-(10).

Let $X(t)$ be a fundamental matrix to the differential equation $\frac{dx}{dt} = A(t)x$

on $[\theta_{r-1}, \theta_r)$, $r = \overline{1, N}$.

Then the unique solution to the Cauchy problem for the system of ordinary differential equations (7), (8) at the fixed values $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_{N+1}, \lambda_{N+2})$ has the following form

$$u_r(t) = X(t) \int_{\theta_{r-1}}^t X^{-1}(\tau) A(\tau) d\tau \cdot \lambda_r + X(t) \int_{\theta_{r-1}}^t X^{-1}(\tau) \sum_{j=0}^N K_j(\tau) d\tau \lambda_{j+1} + \\ + X(t) \int_{\theta_{r-1}}^t X^{-1}(\tau) A_0(\tau) d\tau \lambda_{N+2} + X(t) \int_{\theta_{r-1}}^t X^{-1}(\tau) f(\tau) d\tau, \quad t \in (\theta_{r-1}, \theta_r), \quad r = \overline{1, N}. \quad (11)$$

Substituting (11) into continuity conditions (10) and taking into account (9), we get the system of algebraic equations with respect to unknown parameters $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_{N+1}, \lambda_{N+2}) \in R^{nN+m}$:

$$B\lambda_1 + C\lambda_{N+1} + B_0\lambda_{N+2} = d, \quad (12)$$

$$\lambda_s + X(\theta_s) \int_{\theta_{s-1}}^{\theta_s} X^{-1}(\tau) A(\tau) d\tau \cdot \lambda_r + X(\theta_s) \int_{\theta_{s-1}}^{\theta_s} X^{-1}(\tau) \sum_{j=0}^N K_j(\tau) d\tau \lambda_{j+1} - \lambda_{s+1} + \\ + X(\theta_s) \int_{\theta_{s-1}}^{\theta_s} X^{-1}(\tau) A_0(\tau) d\tau \cdot \lambda_{N+2} + X(\theta_s) \int_{\theta_{s-1}}^{\theta_s} X^{-1}(\tau) f(\tau) d\tau = 0, \quad s = \overline{1, N}. \quad (13)$$

Denoting by $Q_*(\Delta_N)$ the matrix corresponding to the left-hand side of system (12), (13) which is consist of the coefficients at the parameters $\lambda_r, \quad r = \overline{1, N+2}$, and then introducing the vector

$$F_*(\Delta_N) = \left(-d, X(\theta_1) \int_{\theta_0}^{\theta_1} X^{-1}(\tau) f(\tau) d\tau, \dots, X(\theta_N) \int_{\theta_{N-1}}^{\theta_N} X^{-1}(\tau) f(\tau) d\tau \right),$$

we write the system (12), (13) as:

$$Q_*(\Delta_N)\lambda = -F_*(\Delta_N), \quad \lambda \in R^{nN+m}. \quad (14)$$

The boundary value problem (1), (2) is solved by the following algorithm:

As can be seen from the equations (12), (13), the coefficients and right-hand side of the system (14) are composed of solutions to the Cauchy problems

$$\frac{dz}{dt} = A(t)z + A(t), \quad z(\theta_{r-1}) = 0, \quad r = \overline{1, N}, \quad (15)$$

$$\frac{dz}{dt} = A(t)z + K_j(t), \quad z(\theta_{r-1}) = 0, \quad r = \overline{1, N}, \quad (16)$$

$$\frac{dz}{dt} = A(t)z + A_0(t), \quad z(\theta_{r-1}) = 0, \quad r = \overline{1, N}, \quad (17)$$

$$\frac{dz}{dt} = A(t)z + f(t), \quad z(\theta_{r-1}) = 0, \quad r = \overline{1, N}. \quad (18)$$

Construct the following algorithm for the numerical solving of two-point boundary value problem for the systems of loaded differential equations with parameter by applying the Runge–Kutta method of 4th order for numerical solving of the Cauchy problem (15) – (18).

Suppose we have a partition $\theta_0 = 0 < \theta_1 < \dots < \theta_{N-1} < \theta_N = T$. Divide each subinterval $[\theta_{i-1}, \theta_i], \quad i = \overline{1, N}$, into N_i parts. Define the approximate values of coefficients and right-hand side of (14) via solutions to the Cauchy matrix and vector problems obtained using the Runge–Kutta method of 4th order with step $h_i = (\theta_i - \theta_{i-1}) / N_i, \quad i = \overline{1, N}$. Then we obtain the following approximate system of algebraic equations with respect to parameters λ :

$$Q_*^{\tilde{h}}(\Delta_N)\lambda = -F_*^{\tilde{h}}(\Delta_N), \quad \lambda \in R^{nN+m}. \quad (19)$$

Solving the system of linear algebraic equations (19) we find $\lambda^{\tilde{h}} \in R^{nN+m}$.

As noted above, $\lambda^{\tilde{h}} = (\lambda_1^{\tilde{h}}, \lambda_2^{\tilde{h}}, \dots, \lambda_{N+2}^{\tilde{h}}) \in R^{nN+m}$ components are the values of approximate solution to problem (1), (2) at the initial points of subintervals:

$$x^{\tilde{h}_r}(\theta_0) = \lambda_1^{\tilde{h}}, x^{\tilde{h}_r}(\theta_1) = \lambda_2^{\tilde{h}}, \dots, x^{\tilde{h}_r}(\theta_N) = \lambda_{N+1}^{\tilde{h}}, \mu^{\tilde{h}_r} = \lambda_{N+2}^{\tilde{h}}.$$

Applying the Runge–Kutta method of 4th order for numerical solving of Cauchy problem

$$\begin{aligned} \frac{dx}{dt} &= A(t)x + \sum_{j=0}^N K_j(t)\lambda_{j+1}^{\tilde{h}} + A_0\lambda_{N+2}^{\tilde{h}} + f(t), \\ x(\theta_{r-1}) &= \lambda^{\tilde{h}}, t \in [\theta_{r-1}, \theta_r), \quad r = \overline{1, N}, \end{aligned}$$

we determine the numerical solution to problem (1), (2). To illustrate the proposed approach of the numerical solving of two-point boundary value problem for systems of loaded differential equations with parameter (1), (2) based on the parametrization method, let us consider the following example

Example. Consider on $[0, T]$ the linear two-point boundary value problem for the systems of loaded differential equations with parameter:

$$\begin{aligned} \frac{dx}{dt} &= A(t)x + K_0(t)x(\theta_0) + K_1(t)x(\theta_1) + K_2(t)x(\theta_2) + A_0(t)\mu + f(t), \\ t &\in [0, T], \quad x \in R^2, \quad \mu \in R^3, \end{aligned} \tag{20}$$

$$Bx(0) + Cx(T) + B_0\mu = d, \quad d \in R^5, \tag{21}$$

where $A(t) = \begin{pmatrix} t^2 & t+1 \\ 2 & 3t \end{pmatrix}, K_0(t) = \begin{pmatrix} t & 2 \\ t^2 & t-4 \end{pmatrix}, K_1(t) = \begin{pmatrix} 8 & t^2+1 \\ t & 3t \end{pmatrix}, K_2(t) = \begin{pmatrix} t^3 & 0 \\ t & 4 \end{pmatrix},$

$$A_0(t) = \begin{pmatrix} 1 & t & t^2 \\ t+3 & 2 & t^3 \end{pmatrix}, f(t) = \begin{pmatrix} -2t^4 - 2t^3 - \frac{59}{8}t^2 + 7t + \frac{101}{8} \\ -3t^4 - 12t^3 + 2t^2 + \frac{117}{8}t - 5 \end{pmatrix},$$

$$B = \begin{pmatrix} 1 & 0 \\ 2 & 1 \\ 3 & -2 \\ 0 & 4 \\ 7 & 9 \end{pmatrix}, C = \begin{pmatrix} 5 & -2 \\ 0 & 4 \\ 5 & 3 \\ 8 & 11 \\ 7 & 8 \end{pmatrix}, B_0 = \begin{pmatrix} 1 & 3 & -5 \\ 0 & 2 & 4 \\ 4 & -2 & 3 \\ -1 & 3 & 8 \\ 4 & -2 & 0 \end{pmatrix}, d = \begin{pmatrix} -51 \\ 28 \\ 36 \\ 57 \\ -17 \end{pmatrix}.$$

In this example, the matrix of differential part is variable and the construction of fundamental matrix fails. We use a numerical implementation of algorithm of the parametrization method. Below we provide the results of numerical implementation of the algorithm by partitioning the subintervals $[0,0.5], [0.5,1]$ with step $h_1 = h_2 = 0.05$.

Solving the system of equations (19) we obtain the numerical values of the parameters:

$$\lambda_1^{\tilde{h}} = \begin{pmatrix} -0.99999562 \\ -2.00000517 \end{pmatrix}, \lambda_2^{\tilde{h}} = \begin{pmatrix} -0.74999836 \\ -1.62499829 \end{pmatrix}, \lambda_3^{\tilde{h}} = \begin{pmatrix} 0.00000017 \\ 0.00000511 \end{pmatrix}.$$

We find the numerical solutions at the other points of the subintervals applying the Runge-Kutta method of the 4th order to the following Cauchy problems:

$$\frac{dx_1}{dt} = A(t)x_1 + K_0(t)\lambda_1^{\tilde{h}} + K_1(t)\lambda_2^{\tilde{h}} + K_2(t)\lambda_3^{\tilde{h}} + B(t)\lambda_4^{\tilde{h}} + f(t), \quad t \in \left[0, \frac{1}{2}\right), \quad x_1(0) = \lambda_1^{\tilde{h}},$$

$$\frac{dx_2}{dt} = A(t)x_2 + K_0(t)\lambda_1^{\tilde{h}} + K_1(t)\lambda_2^{\tilde{h}} + K_2(t)\lambda_3^{\tilde{h}} + B(t)\lambda_4^{\tilde{h}} + f(t), \quad t \in \left[\frac{1}{2}, 1\right), \quad x_2\left(\frac{1}{2}\right) = \lambda_2^{\tilde{h}}.$$

The exact solution of the problem (20)-(21) is a pair $(\mu^*, x^*(t))$, where $\mu^* = \begin{pmatrix} 1 \\ -2 \\ 9 \end{pmatrix}$,

$$x^*(t) = \begin{pmatrix} t^2 - 1 \\ t^3 + t^2 - 2 \end{pmatrix}, \quad t \in [0, 1].$$

The results of calculations of numerical and exact solutions at the partition points are presented in the following table:

t	$\tilde{x}_1(t)$ (numerical solution)	$x_1^*(t)$	$\tilde{x}_2(t)$ (numerical solution)	$x_2^*(t)$
0	-0.99999562	-1	-2.00000517	-2
0.05	-0.99749604	-0.9975	-1.99737924	-1.997375
0.1	-0.98999643	-0.99	-1.98900339	-1.989
0.15	-0.97749678	-0.9775	-1.97412762	-1.974125
0.2	-0.9599971	-0.96	-1.9520019	-1.952
0.25	-0.93749739	-0.9375	-1.92187622	-1.921875
0.3	-0.90999764	-0.91	-1.88300059	-1.883
0.35	-0.87749786	-0.8775	-1.83462498	-1.834625
0.4	-0.83999805	-0.84	-1.7759994	-1.776
0.45	-0.79749822	-0.7975	-1.70637383	-1.706375
0.5	-0.74999836	-0.75	-1.62499829	-1.625
0.55	-0.69749848	-0.6975	-1.53112275	-1.531125
0.6	-0.63999858	-0.64	-1.42399724	-1.424
0.65	-0.57749867	-0.5775	-1.30287174	-1.302875
0.7	-0.50999875	-0.51	-1.16699627	-1.167
0.75	-0.43749884	-0.4375	-1.01562083	-1.015625
0.8	-0.35999893	-0.36	-0.84799544	-0.848
0.85	-0.27749906	-0.2775	-0.66337012	-0.663375
0.9	-0.18999923	-0.19	-0.46099489	-0.461
0.95	-0.09749948	-0.0975	-0.2401198	-0.240125
1	0.00000017	0	0.00000511	0

$\tilde{\mu}_1 = \lambda_{41}^{\tilde{h}}$ (numerical solution)	μ_1^*	$\tilde{\mu}_2 = \lambda_{42}^{\tilde{h}}$ (numerical solution)	μ_2^*	$\tilde{\mu}_3 = \lambda_{43}^{\tilde{h}}$ (numerical solution)	μ_3^*
0.9999919	1	-2.00000308	-2	8.99999553	9

For the difference of the corresponding values of the exact and constructed solutions of the problem the following estimate is true:

$$\max \|\mu^* - \tilde{\mu}\| < 0.0000081,$$

$$\max_{j=0,20} \|x^*(t_j) - \tilde{x}(t_j)\| < \varepsilon, \quad \varepsilon = 0.000005.$$

УДК 517.62/64, 517.968.7, 517.977.1/5
МРНТИ 27.33.19; 27.37.17; 27.41.19

А.Т. Асанова^{1,2}, Э.А. Бакирова^{1,2,3}, Ж.М. Кадирбаева^{1,2,4}

¹Математика және математикалық модельдеу институты, Алматы, Қазақстан

²Ақпараттық және есептеуіш технологиялар институты, Алматы, Қазақстан

³Қазақ Ұлттық қыздар педагогикалық университеті, Алматы, Қазақстан

⁴Халықаралық ақпараттық технологиялар университеті, Алматы, Қазақстан

ПАРАМЕТРІ БАР ЖҮКТЕЛГЕН ДИФФЕРЕНЦИАЛДЫҚ ТЕНДЕУЛЕР ЖҮЙЕСІ ҮШІН ШЕТТІК ЕСЕПТІ ШЕШУДІҢ САНДЫҚ ЖҮЗЕГЕ АСЫРЫЛУЫ

Аннотация. Параметрі бар жүктелген дифференциалдық тендеулер үшін сызықты екі нүктелі шеттік есеп қарастырылады. Аталған есеп параметрлеу әдісі арқылы зерттеледі. Параметрі бар жүктелген дифференциалдық тендеулер жүйесі үшін шеттік есептің шешімін табудың алгоритмі ұсынылады. Алдымен бастапқы есеп ішкіаралықтардағы параметрлері бар жәй дифференциалдық тендеулер жүйесі үшін Коши есебін және енгізілген параметрлерге қатысты функционалдық қатынастарды қамтитын пара-пар есепке келтіріледі. Параметрлердің бекітілген мәнінде ішкіаралықтағы жәй дифференциалдық тендеулер жүйесі үшін Коши есебінің жалғыз шешімі бар болады. Бұл шешім жүйенің фундаменталдық матрицасы арқылы кейіптеледі. Осы кейіптемелерді пайдалана отырып параметрлерге қатысты сызықты алгебралық тендеулер жүйесін құрамыз. Пара-пар есептің сандық шешімін табуға арналған алгоритм ұсынылады. Бұл алгоритм жәй дифференциалдық тендеулер жүйесі үшін Коши есептерін сандық шешуді және алгебралық тендеулер жүйесін шешуді қамтиды. Коши есептерін сандық түрде шешу үшін төртінші ретті Рунге-Куттаның әдісі қолданылады. Ұсынылып отырған сандық жүзеге асырылу мысалмен көрнектеледі.

Кілттік сөздер: параметрі бар шеттік есеп, жүктелген дифференциалдық тендеу, сандық әдіс, алгоритм.

УДК 517.62/64, 517.968.7, 517.977.1/5
МРНТИ 27.33.19; 27.37.17; 27.41.19

А.Т. Асанова^{1,2}, Э.А. Бакирова^{1,2,3}, Ж.М. Кадирбаева^{1,2,4}

¹Институт математики и математического моделирования, Алматы, Казахстан;

²Институт информационных и вычислительных технологий, Алматы, Казахстан;

³Казахский Национальный Женский Педагогический Университет, Алматы, Казахстан;

⁴Международный Университет Информационных Технологий, Алматы, Казахстан

ЧИСЛЕННАЯ РЕАЛИЗАЦИЯ РЕШЕНИЯ КРАЕВОЙ ЗАДАЧИ ДЛЯ СИСТЕМЫ НАГРУЖЕННЫХ ДИФФЕРЕНЦИАЛЬНЫХ УРАВНЕНИЙ С ПАРАМЕТРОМ

Аннотация. Рассматривается линейная двухточечная краевая задача для системы нагруженных дифференциальных уравнений с параметром. Данная задача исследуется методом параметризации. Предлагается алгоритм нахождения решения краевой задачи для системы нагруженных дифференциальных уравнений с параметром. Вначале исходная задача сводится к эквивалентной задаче, состоящей из задач Коши для системы обыкновенных дифференциальных уравнений с параметрами на подинтервалах и функциональных соотношений относительно введенных дополнительных параметров. При фиксированных значениях параметров задача Коши для системы обыкновенных дифференциальных уравнений на подинтервале имеет единственное решение. Это решение представляется через фундаментальную матрицу системы. Используя эти представления составляется система линейных алгебраических уравнений относительно параметров. Предлагается алгоритм нахождения численного решения эквивалентной задачи. Данный алгоритм включает численное решение задач Коши для системы обыкновенных дифференциальных уравнений и решение линейной системы алгебраических уравнений. Для численного решения задачи Коши применяется метод Рунге-Кутты четвертого порядка. Предлагаемая численная реализация иллюстрируется примером.

Ключевые слова: краевая задача с параметром, нагруженное дифференциальное уравнение, численный метод, алгоритм.

Information about authors:

Assanova Anar Turmaganbetkyzy, the member of Editorial Board of journal “News of the NAS RK. Physico-Mathematical Series”,

Institute of Mathematics and Mathematical Modeling, Institute of Information and Computational Technologies, chief scientific researcher, Doctor of Physical and Mathematical Sciences, professor, assanova@math.kz; <https://orcid.org/0000-0001-8697-8920>;

Bakirova Elmira Ainabekovna,

Institute of Mathematics and Mathematical Modeling, Institute of Information and Computational Technologies, leading scientific researcher, Candidate of Physical and Mathematical Sciences, ass. professor, Kazakh National Women's Teacher Training University, associated professor, bakirova1974@mail.ru; <https://orcid.org/0000-0002-3820-5373>;

Kadirbayeva Zhazira Muratbekovna,

Institute of Mathematics and Mathematical Modeling, Institute of Information and Computational Technologies, leading scientific researcher, Candidate of Physical and Mathematical Sciences, International Information Technology University, head teacher, zh.kadirbayeva@gmail.com; <https://orcid.org/0000-0001-8861-4100>

REFERENCES

- [1] Nakhushev A.M. Equations of mathematical biology. M.: Vyshaiya shkola, 1995. (in Russ.)
- [2] Dzhenaliev M.T. To the theory linear boundary value problems for loaded differential equations. Almaty: Computenryi centr ITPM, 1995. (in Russ.)
- [3] Nakhushev A.M. Problems with replacement for partial differential equations. M.: Nauka, 2006. (in Russ.)
- [4] Dzhenaliev M.T., Ramazanov M.I. Loaded equations as perturbations of differential equations. Almaty: Gylym, 2010. (in Russ.)
- [5] Nakhushev A.M. Loaded equations and their applications. M.: Nauka, 2012. (in Russ.)
- [6] Bakirova E.A. About of a criteria of unique solvability of two-point boundary value problem for system of loaded differential equations, *Izvestiya NAN RK. Ser.phyz.-matem.* 2005. No 1. P.95-102. (in Russ.)
- [7] Bakirova E.A. About of a necessary and sufficient conditions of unique solvability of two-point boundary value problem for loaded differential equations, *Mathematical journal.* 2005. Vol. 5. No 3. P.25-34. (in Russ.)
- [8] Kadirbayeva Zh.M. About of one algorithm for finding solution of linear two-point boundary value problem for loaded differential equations, *Mathematical journal.* 2009. Vol.9. No 2(32). P.64-70. (in Russ.)
- [9] Kadirbayeva Zh.M. About of a necessary and sufficient conditions of unique solvability of two-point boundary value problem for loaded differential equations, *Mathematical journal.* 2009. Vol.9. No 4(34). P.63-71. (in Russ.)
- [10] Assanova A.T., Kadirbayeva Zh.M. On the numerical algorithms of parametrization method for solving a two-point boundary-value problem for impulsive systems of loaded differential equations, *Computational and Applied Mathematics.* 2018. - Vol. 37, No. 4. -P. 4966-4976. DOI: <https://doi.org/10.1007/s40314-018-0611-9>
- [11] Dzhumabaev D.S. Criteria for the unique solvability of a linear boundary-value problem for an ordinary differential equation, *USSR Comp. Math. & Math. Phys.* 1989. V.29, No. 1. P. 34-46.
- [12] Dzhumabaev D.S. A method for solving the linear boundary value problem for an integro-differential equation, *Comp. Math. & Math. Phys.* 2010. V.50, No. 7. P. 1150-1161. DOI: <https://doi.org/10.1134/S0965542510070043>
- [13] Dzhumabaev D.S. An algorithm for solving a linear two-point boundary value problem for an integrodifferential equation, *Comp. Math. & Math. Phys.* 2013. V.53, No. 6. P. 736-758. DOI: <https://doi.org/10.1134/S0965542513060067>
- [14] Dzhumabaev D.S., Bakirova E.A. Criteria for the unique solvability of a linear two-point boundary value problem for systems of integro-differential equations, *Differ. Equ.* 2013. V.49, No. 9. P. 1087-1102. DOI: <https://doi.org/10.1134/S0012266113090048>
- [15] Dzhumabaev D.S. Necessary and sufficient conditions for the solvability of linear boundary-value problems for the Fredholm integrodifferential equations, *Ukrainian Math. J.* 2015. V.66, No. 8. P. 1200-1219. DOI: <https://doi.org/10.1007/s11253-015-1003-6>
- [16] Dzhumabaev D.S. Solvability of a linear boundary value problem for a Fredholm integro-differential equation with impulsive inputs, *Differ. Equ.* 2015. V. 51, No. 9. P. 1180-1196. DOI: <https://doi.org/10.1134/S0012266115090086>
- [17] Dzhumabaev D.S. On one approach to solve the linear boundary value problems for Fredholm integro-differential equations, *J. Comp. Appl. Math.* 2016. V. 294, P. 342-357. DOI: <https://doi.org/10.1016/j.cam.2015.08.023>
- [18] Dzhumabaev D.S., Bakirova E.A., Kadirbayeva Zh.M. An algorithm for solving a control problem for a differential equation with a parameter, *News of the NAS RK. Phys.-Math. Series.* 2018. Volume 5, Number 321. P.25-32. DOI: <https://doi.org/10.32014/2018.2518-1726.4>

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы *М. С. Ахметова, Т.А. Апендиев, Д.С. Аленов*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 10.06.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
8,3 п.л. Тираж 300. Заказ 3.

Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19