

**ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)**

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Әль-фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Казахский национальный университет
имени Аль-фараби

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-farabi kazakh
national university

SERIES
PHYSICO-MATHEMATICAL

3 (325)

MAY - JUNE 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Бас редакторы
ф.-м.ғ.д., проф., КР ҮФА академигі **F.M. Мұтанов**

Редакция алқасы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев Ү.Ү. проф. корр.-мүшесі (Қазақстан)
Жусіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Ә. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорус)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«КР ҮФА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Үлттық ғылым академиясы» РКБ (Алматы қ.)
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы қуәлік

Мерзімділігі: жылдана 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Үлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Р е д а к ц и о н на я кол л е г и я:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жантаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф. чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

E d i t o r i n c h i e f
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d:

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskyi I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

N E W S

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.28>

Volume 3, Number 325 (2019), 85 – 96

UDK 517.43

A.Sh. Shaldanbayev¹, A.B. Imanbayeva², A.Zh. Beisebayeva³, A.A. Shaldanbayeva⁴¹“Silkway” International University, Shymkent;^{2,3}M.Auezov South Kazakhstan State University, Shymkent;⁴“Regional Social-Innovative University”, Shymkent.shaldanbaev51@mail.ru, aigul_baratovna@bk.ru, akbope_a@mail.ru, altima_a@mail.ru**ON THE SQUARE ROOT OF THE OPERATOR
OF STURM-LIOUVILLE FOURTH-ORDER**

Abstract. In the present work found the root of the positive operator of the Sturm - Liouville problem of the fourth order, which is invertible composition operator of the Sturm - Liouville problem and its adjoint. The found root does not possess the property of positivity, but is a self-adjoint operator in the essential. One theorem of Putnam of algebraic character is used as a leading idea. It is hoped that the results will find applications in spectral operator theory and theoretical physics.

Keywords. Kato conjecture, dissipative operator, square root of operator, Putnam theorem, deviating argument, fractional powers of an operator, inverse problem, spectrum, unitary operator, self-adjoint operator, positive operator, functional differential operator, spectral theory.

1. Introduction.

Definition. Let H be a Hilbert space. A linear bounded operator B is called positive if $(Bx, x) \geq 0$ for all $x \in H$. We write $B \geq 0$ if B is positive.

Lemma 1.1 (On the square root). Let A be a linear bounded operator and $A \geq 0$. Then there is a single operator $B \gg 0$ and $B^2 = A$. Moreover, B commutes with any bounded operator commuting with A . [1. p.219].

When extending the concept of root to dissipative operators, the hypothesis of Kato [3] arose, consisting in the fact that the domain of the root definition from the operator always coincides with the domain of the root definition from the conjugate operator. But in 1972 A. Macintosh [4] built a counterexample, since then the hypothesis has been slightly reformulated: to find the largest class of operators that satisfies this condition, and is now given very active research in this direction [5-57].

Many operators of theoretical physics have square roots [57–64]; in particular, the square root of an operator in a Banach space was found in [65, pp.169-176]. We give an excerpt from this work.

Consider a generating operator A in a Banach space \mathcal{B} with the following properties:

- 1) The operator $(I + \gamma^2 A)^{-1}$ exists, is defined everywhere in \mathcal{B} and is bounded by one;
- 2) The operator A^{-1} exists;
- 3) $\|e^{iAt}\| \leq M$, $-\infty < t < +\infty$.

Under these conditions, the following lemmas are valid.

Lemma 1.1. Operator

$$T = \frac{2e^{-i\pi/4}}{\sqrt{\pi}} A \int_0^\infty e^{iAx^2} dx$$

exists as an operator in \mathcal{B} on the area $D(A)$.

Lemma 1.2. For any $y \in D(A)$ the equality is true

$$T^2 g = Ag.$$

In connection with these results, the following problem arises.

Formulation of the problem. Let a reversible Sturm-Liouville operator L

$$Ly = -y''(x), \quad x \in (0,1)$$

$$\begin{cases} a_{11}y(0) + a_{12}y'(0) + a_{13}y(1) + a_{14}y'(1) = 0, \\ a_{21}y(0) + a_{22}y'(0) + a_{23}y(1) + a_{24}y'(1) = 0, \end{cases}$$

where a_{ij} ($i = 1, 2; j = 1, 2, 3, 4$) – are complex numbers. Then it takes the following form:

$$Ly = -y''(x), \quad x \in (0,1) \quad (1)$$

$$\begin{cases} \Delta_{13}y(0) - (\Delta_{12} + \Delta_{32})y'(0) - \Delta_{13}y(1) - \\ \quad - (\Delta_{14} + \Delta_{34})y'(1) = 0; \\ (\Delta_{12} + \Delta_{13} + \Delta_{14})y(0) - (\Delta_{32} + \Delta_{42})y'(0) + \\ \quad + (\Delta_{32} + \Delta_{34})y(1) - (\Delta_{34} + \Delta_{24})y'(1) = 0; \end{cases} \quad (2)$$

conjugate which has the form:

$$L^+ z = -z''(x), \quad x \in (0,1), \quad (1)^+$$

$$\begin{cases} \bar{\Delta}_{13}z(0) - (\bar{\Delta}_{34} + \bar{\Delta}_{32})z'(0) - \bar{\Delta}_{13}z(1) - \\ \quad - (\bar{\Delta}_{14} + \bar{\Delta}_{12})z'(1) = 0; \\ (\Delta_{12} + \Delta_{13} + \Delta_{14})y(0) - (\Delta_{32} + \Delta_{42})y'(0) + \\ \quad + (\Delta_{32} + \Delta_{34})y(1) - (\Delta_{34} + \Delta_{24})y'(1) = 0. \end{cases} \quad (2)^+$$

The question is whether there is a unitary operator.

$$T = i \cos \varphi I + \sin \varphi \cdot S,$$

such that the formula takes place

$$TL = L^+ T^*, \quad (3)$$

where I – is an identical operator and S is:

$$Su(x) = u(1-x). \quad (4)$$

2. Research methods.

As a suggestive idea, we take the following theorem of Putnam.

Theorem [2. p.337]. $M, N, T \in \mathcal{B}(H)$ Suppose that the operators M, N are normal and the operator T is invertible. Suppose that

$$M = TNT^{-1}. \quad (*)$$

If $T = UP$ – the polar decomposition of the operator T , then

$$M = UNU^{-1}.$$

Two operators connected by a relation (*) are called similar. If U – is a unitary operator and the relation (1.9) is satisfied, then the operators M and N are called unitary equivalent. Thus, this theorem establishes that such normal operators are unitary equivalent.

Our operators A, B are Hermitian (i.e, symmetric), and such operators belong to the class of normal operators; so there is a unitary operator T such that

$$AT = TB.$$

We believe that this particular operator is the solution of the equations:

$$(Tl)^2 = l^+l = A, \quad (lT)^2 = ll^+ = B.$$

Perhaps the operator T we need to impose additional conditions?!

From formula

$$(Tl)^2 = Tl \cdot Tl = l^*l,$$

we see that we need to require $Tl = l^*T^*$, then

$$Tl \cdot Tl = l^* \underbrace{T^*T}_I l = l^*l = A,$$

Next, from $Tl = l^*T^*$, we have

$$TlT = l^*, \quad lT = T^{-1}l^* = T^*l^*,$$

then

$$(lT)^2 = lT \cdot lT = |lT = T^*l^*| = lT \cdot T^*l^* = ll^* = B.$$

In addition,

$$TB = Tll^* = l^*T^*l^* = l^*lT = AT.$$

We proved the following Lemma.

Lemma 2.1. If T – is a unitary operator satisfying the condition

$$(Tl)^* = Tl = l^*T^*,$$

then we have the formula

a) $(Tl)^2 = l^*l = A$,

b) $(lT)^2 = ll^* = B$,

c) $AT = TB$.

Thus, the problem was reduced to finding a unitary operator T , with the property, $Tl = l^*T^*$. This Lemma forms the basis of our method.

3. Research Results.

Lety(x) $\in D(L)$, then

$$z(x) = T^*y(x) = -i \cos \varphi y(x) + \sin \varphi \cdot y(1-x) \in D(L^+),$$

therefore, there are formulas

$$\begin{aligned} z(x) &= -i \cos \varphi y(x) + \sin \varphi \cdot y(1-x), \\ z'(x) &= -i \cos \varphi y'(x) - \sin \varphi \cdot y'(1-x), \\ z(0) &= -i \cos \varphi y(0) + \sin \varphi \cdot y(1), \\ z'(0) &= -i \cos \varphi y'(0) - \sin \varphi \cdot y'(1), \\ z(1) &= -i \cos \varphi y(1) + \sin \varphi \cdot y(1), \\ z'(1) &= -i \cos \varphi y'(1) - \sin \varphi \cdot y'(0), \\ z(0) - z(1) &= (i \cos \varphi + \sin \varphi)y(1) - (i \cos \varphi + \sin \varphi)y(0) = \\ &= (i \cos \varphi + \sin \varphi)[y(1) - y(0)], \\ \overline{\Delta_{13}}(i \cos \varphi + \sin \varphi)[y(1) - y(0)] + (\overline{\Delta_{32}} + \overline{\Delta_{34}}) & \\ [i \cos \varphi y'(0) + \sin \varphi \cdot y'(1)] + & \\ + (\overline{\Delta_{12}} + \overline{\Delta_{14}})[i \cos \varphi y'(1) + \sin \varphi \cdot y'(0)] &= 0; \\ \overline{\Delta_{13}}(i \cos \varphi + \sin \varphi)[y(1) - y(0)] + & \\ + [i \cos \varphi (\overline{\Delta_{32}} + \overline{\Delta_{34}}) + \sin \varphi (\overline{\Delta_{12}} + \overline{\Delta_{14}})]y'(0) + & \\ + [\sin \varphi (\overline{\Delta_{32}} + \overline{\Delta_{34}}) + i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{14}})]y'(1) &= 0; \end{aligned}$$

$$\begin{aligned}
 & (\overline{\Delta_{12}} + \overline{\Delta_{14}} + \overline{\Delta_{34}})[-i \cos \varphi y(0) + \sin \varphi \cdot y(1)] + \\
 & + (\overline{\Delta_{32}} + \overline{\Delta_{34}})[i \cos \varphi y'(0) + \sin \varphi \cdot y'(1)] + \\
 & + (\overline{\Delta_{12}} + \overline{\Delta_{32}})[-i \cos \varphi y(1) + \sin \varphi \cdot y(0)] + (\overline{\Delta_{12}} + \overline{\Delta_{24}}) \cdot \\
 & \quad \cdot [i \cos \varphi y'(1) + \sin \varphi \cdot y'(0)] = \\
 & = [-i \cos \varphi (\overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) + (\overline{\Delta_{12}} + \overline{\Delta_{32}}) \sin \varphi]y(0) + \\
 & + [\sin \varphi (\overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) - i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{32}})]y(1) + \\
 & + [(\overline{\Delta_{32}} + \overline{\Delta_{42}}) \sin \varphi + i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{24}})]y'(1) = 0.
 \end{aligned}$$

This boundary condition corresponds to the matrix

$$\begin{pmatrix}
 -\overline{\Delta_{13}}(\sin \varphi + i \cos \varphi) \\
 -i \cos \varphi (\overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) + (\overline{\Delta_{12}} + \overline{\Delta_{32}}) \sin \varphi \\
 i \cos \varphi (\overline{\Delta_{32}} + \overline{\Delta_{42}}) + \sin \varphi (\overline{\Delta_{12}} + \overline{\Delta_{24}}) \\
 \overline{\Delta_{13}}(\sin \varphi + i \cos \varphi) \\
 \sin \varphi (\overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) - i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{32}}) \\
 \sin \varphi (\overline{\Delta_{32}} + \overline{\Delta_{34}}) + i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{14}}) \\
 (\overline{\Delta_{32}} + \overline{\Delta_{42}}) \sin \varphi + i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{24}})
 \end{pmatrix}.$$

Combining the obtained system of equations with respect to the unknowns: $y(0), y'(0), y(1), y'(1)$ with the previously known boundary conditions (2), we obtain a system of homogeneous algebraic equations. Since this resulting system of equations obviously has nontrivial solutions, its determinant is zero.

Taking advantage of this fact, we obtain one equation to determine the unknown value φ .

$$\left\{
 \begin{array}{l}
 1) \Delta_{13}y(0) - (\Delta_{12} + \Delta_{32})y'(0) - \Delta_{13}y(1) - \\
 \quad (\Delta_{14} + \Delta_{34})y'(1) = 0; \\
 2) (\Delta_{12} + \Delta_{13} + \Delta_{14})y(0) - (\Delta_{32} + \Delta_{42})y'(0) + \\
 \quad (\Delta_{32} + \Delta_{34})y(1) - (\Delta_{34} + \Delta_{24})y'(1) = 0; \\
 3) \overline{\Delta_{13}}(\sin \varphi + i \cos \varphi)y(0) + [i \cos \varphi (\overline{\Delta_{32}} + \overline{\Delta_{34}}) + (\overline{\Delta_{12}} + \overline{\Delta_{14}}) \sin \varphi] \cdot \\
 \quad \cdot y'(0) - \Delta_{13}(\sin \varphi + i \cos \varphi)y(1) + \\
 \quad + [(\overline{\Delta_{32}} + \overline{\Delta_{34}}) \sin \varphi + i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{14}})]y'(1) = 0; \\
 4) [-i \cos \varphi (\overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) + \sin \varphi (\overline{\Delta_{12}} + \overline{\Delta_{32}})]y(0) + \\
 \quad + [i \cos \varphi (\overline{\Delta_{32}} + \overline{\Delta_{42}}) + \sin \varphi (\overline{\Delta_{12}} + \overline{\Delta_{24}})]y'(0) + \\
 \quad [(\overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) \cdot \sin \varphi - i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{32}})]y(1) + \\
 \quad + [(\overline{\Delta_{32}} + \overline{\Delta_{42}}) \sin \varphi + i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{24}})]y'(1) = 0.
 \end{array}
 \right.$$

This system of homogeneous algebraic equations is denoted by (5).

The determinant of the system of equations (5) has the following form:

$$\begin{bmatrix}
 \Delta_{13} \\
 \Delta_{12} + \Delta_{13} + \Delta_{14} \\
 \overline{\Delta_{13}}(\sin \varphi + i \cos \varphi) \\
 -i \cos \varphi (\overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) + \sin \varphi (\overline{\Delta_{12}} + \overline{\Delta_{32}})
 \end{bmatrix}$$

$$\begin{aligned}
& -(\Delta_{12} + \Delta_{32}) \\
& -(\Delta_{32} + \Delta_{42}) \\
& i \cos \varphi (\overline{\Delta_{32}} + \overline{\Delta_{34}}) + (\overline{\Delta_{12}} + \overline{\Delta_{14}}) \sin \varphi \\
& [i \cos \varphi (\overline{\Delta_{32}} + \overline{\Delta_{42}}) + \sin \varphi (\overline{\Delta_{12}} + \overline{\Delta_{24}})] \\
& \quad \frac{-\Delta_{13}}{\Delta_{32} + \Delta_{34}} \\
& \quad -\overline{\Delta_{13}} (\sin \varphi + i \cos \varphi) \\
& (\overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) \sin \varphi - i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{32}}) \\
& \quad \left. \begin{array}{l} -(\Delta_{14} + \Delta_{34}) \\ -(\Delta_{34} + \Delta_{24}) \\ (\overline{\Delta_{32}} + \overline{\Delta_{34}}) \sin \varphi + i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{14}}) \\ (\overline{\Delta_{32}} + \overline{\Delta_{42}}) \sin \varphi + i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{24}}) \end{array} \right] \\
& \quad \left. \begin{array}{l} -(\Delta_{12} + \Delta_{32}) \\ -(\Delta_{32} + \Delta_{42}) \\ (\overline{\Delta_{12}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) \sin \varphi \\ (\overline{\Delta_{12}} + \overline{\Delta_{24}}) \sin \varphi \end{array} \right]
\end{aligned}$$

This determinant splits into the sum of two determinants:

$$\begin{aligned}
& \left[\begin{array}{cc} \Delta_{13} & -(\Delta_{12} + \Delta_{32}) \\ \Delta_{12} + \Delta_{13} + \Delta_{14} & -(\Delta_{32} + \Delta_{42}) \\ -\overline{\Delta_{13}} \sin \varphi & (\overline{\Delta_{12}} + \overline{\Delta_{14}}) \sin \varphi \\ (\overline{\Delta_{12}} + \overline{\Delta_{32}}) \sin \varphi & (\overline{\Delta_{12}} + \overline{\Delta_{24}}) \sin \varphi \end{array} \right] \\
& + \left[\begin{array}{cc} \Delta_{13} & -(\Delta_{14} + \Delta_{34}) \\ \Delta_{12} + \Delta_{13} + \Delta_{14} & -(\Delta_{34} + \Delta_{24}) \\ -\overline{\Delta_{13}} i \cos \varphi & (\overline{\Delta_{12}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) \sin \varphi \\ -i \cos \varphi (\overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) & (\overline{\Delta_{13}} + \overline{\Delta_{42}}) \sin \varphi \end{array} \right] \\
& \left. \begin{array}{ccc} -\Delta_{13} & -(\Delta_{14} + \Delta_{34}) & -(\Delta_{12} + \Delta_{32}) \\ \Delta_{32} + \Delta_{34} & -(\Delta_{34} + \Delta_{24}) & -(\Delta_{32} + \Delta_{42}) \\ \overline{\Delta_{13}} \sin \varphi & (\overline{\Delta_{32}} + \overline{\Delta_{34}}) \sin \varphi & (\overline{\Delta_{32}} + \overline{\Delta_{42}}) \sin \varphi \\ (\overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) \sin \varphi & (\overline{\Delta_{32}} + \overline{\Delta_{42}}) \sin \varphi & (\overline{\Delta_{12}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) \sin \varphi \end{array} \right] \\
& \left. \begin{array}{ccc} -(\Delta_{12} + \Delta_{32}) & -\Delta_{13} & -(\Delta_{14} + \Delta_{34}) \\ -(\Delta_{32} + \Delta_{42}) & \Delta_{32} + \Delta_{34} & -(\Delta_{34} + \Delta_{24}) \\ i \cos \varphi (\overline{\Delta_{32}} + \overline{\Delta_{34}}) & i \cdot \overline{\Delta_{13}} \cos \varphi & i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{14}}) \\ i \cos \varphi (\overline{\Delta_{32}} + \overline{\Delta_{42}}) & -i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{32}}) & i \cos \varphi (\overline{\Delta_{12}} + \overline{\Delta_{24}}) \end{array} \right]
\end{aligned}$$

This matrix is denoted by (6).

(a) Calculate the first determinant:

$$\begin{aligned}
& \sin^2 \varphi \left[\begin{array}{cccc} \Delta_{13} & -(\Delta_{12} + \Delta_{32}) & -\Delta_{13} & -(\Delta_{14} + \Delta_{34}) \\ \Delta_{12} + \Delta_{13} + \Delta_{14} & -(\Delta_{32} + \Delta_{42}) & \Delta_{32} + \Delta_{34} & -(\Delta_{34} + \Delta_{24}) \\ -\overline{\Delta_{13}} & \overline{\Delta_{12}} + \overline{\Delta_{14}} & \overline{\Delta_{13}} & \overline{\Delta_{32}} + \overline{\Delta_{34}} \\ \overline{\Delta_{12}} + \overline{\Delta_{32}} & \overline{\Delta_{12}} + \overline{\Delta_{24}} & \overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}} & \overline{\Delta_{32}} + \overline{\Delta_{42}} \end{array} \right] = \\
& = \sin^2 \varphi \left[\begin{array}{cccc} 0 & -\Delta & -\Delta_{13} & -(\Delta_{14} + \Delta_{34}) \\ \Delta & 0 & \Delta_{32} + \Delta_{34} & -(\Delta_{34} + \Delta_{24}) \\ 0 & \bar{\Delta} & \overline{\Delta_{13}} & \overline{\Delta_{32}} + \overline{\Delta_{34}} \\ \bar{\Delta} & \bar{\Delta} & \overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}} & \overline{\Delta_{32}} + \overline{\Delta_{42}} \end{array} \right] = \\
& = \sin^2 \varphi \left[\begin{array}{cccc} 0 & -\Delta & -\Delta_{13} & -(\Delta_{14} + \Delta_{34}) \\ \Delta & 0 & \Delta_{32} + \Delta_{34} & -(\Delta_{34} + \Delta_{24}) \\ 0 & \bar{\Delta} & \overline{\Delta_{13}} & \overline{\Delta_{32}} + \overline{\Delta_{34}} \\ \bar{\Delta} & 0 & \overline{\Delta_{14}} + \overline{\Delta_{34}} & -(\Delta_{24} + \Delta_{34}) \end{array} \right] = \\
& \quad \text{—————} 89 \text{ —————}
\end{aligned}$$

$$\begin{aligned}
 &= \sin^2 \varphi \left\{ -\Delta \begin{vmatrix} -\Delta & -\Delta_{13} & -(\Delta_{14} + \Delta_{34}) \\ \bar{\Delta} & \frac{\Delta_{13}}{\Delta_{13}} & \frac{\Delta_{32} + \Delta_{34}}{\Delta_{32} + \Delta_{34}} \\ 0 & \frac{\Delta_{14} + \Delta_{34}}{\Delta_{14} + \Delta_{34}} & -(\Delta_{24} + \Delta_{34}) \end{vmatrix} - \bar{\Delta} \begin{vmatrix} -\Delta & -\Delta_{13} & -(\Delta_{14} + \Delta_{34}) \\ 0 & \Delta_{32} + \Delta_{34} & -(\Delta_{34} + \Delta_{24}) \\ \bar{\Delta} & \frac{\Delta_{13}}{\Delta_{13}} & \frac{\Delta_{32} + \Delta_{34}}{\Delta_{32} + \Delta_{34}} \end{vmatrix} \right\} = \\
 &= \sin^2 \varphi \left\{ \Delta \begin{vmatrix} \Delta & \frac{\Delta_{13}}{\Delta_{13}} & \frac{\Delta_{14} + \Delta_{34}}{\Delta_{32} + \Delta_{34}} \\ \bar{\Delta} & \frac{\Delta_{13}}{\Delta_{13}} & \frac{\Delta_{32} + \Delta_{34}}{\Delta_{32} + \Delta_{34}} \\ 0 & \frac{\Delta_{14} + \Delta_{34}}{\Delta_{14} + \Delta_{34}} & -(\Delta_{24} + \Delta_{34}) \end{vmatrix} - \bar{\Delta} \begin{vmatrix} \Delta & \frac{\Delta_{13}}{\Delta_{13}} & \frac{\Delta_{14} + \Delta_{34}}{\Delta_{32} + \Delta_{34}} \\ 0 & \Delta_{32} + \Delta_{34} & -(\Delta_{34} + \Delta_{24}) \\ \bar{\Delta} & \frac{\Delta_{13}}{\Delta_{13}} & \frac{\Delta_{32} + \Delta_{34}}{\Delta_{32} + \Delta_{34}} \end{vmatrix} \right\} = \\
 &= \sin^2 \varphi \left\{ \Delta \cdot \left[-(\Delta_{14} + \Delta_{34}) \begin{vmatrix} \Delta & \frac{\Delta_{14} + \Delta_{34}}{\Delta_{32} + \Delta_{34}} \\ \bar{\Delta} & \frac{\Delta_{14} + \Delta_{34}}{\Delta_{32} + \Delta_{34}} \end{vmatrix} - (\Delta_{24} + \Delta_{34}) \begin{vmatrix} \Delta & \frac{\Delta_{13}}{\Delta_{13}} \\ \bar{\Delta} & \frac{\Delta_{13}}{\Delta_{13}} \end{vmatrix} \right] - \right. \\
 &\quad \left. - \Delta \left[-(\Delta_{32} + \Delta_{34}) \begin{vmatrix} \Delta & \frac{\Delta_{14} + \Delta_{34}}{\Delta_{32} + \Delta_{34}} \\ \bar{\Delta} & \frac{\Delta_{14} + \Delta_{34}}{\Delta_{32} + \Delta_{34}} \end{vmatrix} - (\Delta_{34} + \Delta_{24}) \begin{vmatrix} \Delta & \frac{\Delta_{13}}{\Delta_{13}} \\ \bar{\Delta} & \frac{\Delta_{13}}{\Delta_{13}} \end{vmatrix} \right] \right\} = \\
 &= \sin^2 \varphi \cdot \left\{ \left| \Delta \frac{\Delta_{14} + \Delta_{34}}{\Delta_{32} + \Delta_{34}} \right| \cdot [\bar{\Delta}(\Delta_{32} + \Delta_{34}) - \Delta(\Delta_{14} + \Delta_{34})] + \right. \\
 &\quad \left. + \left| \Delta \frac{\Delta_{13}}{\Delta_{13}} \right| \cdot [\bar{\Delta}(\Delta_{24} + \Delta_{34}) - \Delta(\Delta_{24} + \Delta_{34})] \right\} = \\
 &= \sin^2 \varphi \cdot \{ [\Delta(\Delta_{32} + \Delta_{34}) - \bar{\Delta}(\Delta_{14} + \Delta_{34})] \cdot [\bar{\Delta}(\Delta_{32} + \Delta_{34}) - \Delta(\Delta_{14} + \Delta_{34})] + \\
 &\quad + (\Delta \cdot \Delta_{13} - \bar{\Delta} \cdot \Delta_{13}) [\bar{\Delta}(\Delta_{24} + \Delta_{34}) - \Delta(\Delta_{24} + \Delta_{34})] \} = \\
 &= \sin^2 \varphi \cdot [|\Delta|^2 \cdot |\Delta_{32} + \Delta_{34}|^2 - \Delta^2(\Delta_{32} + \Delta_{34})(\Delta_{14} + \Delta_{34}) - \bar{\Delta}^2(\Delta_{14} + \Delta_{34})(\Delta_{32} + \Delta_{34}) + \\
 &\quad + |\Delta|^2 \cdot |\Delta_{14} + \Delta_{34}|^2 + |\Delta|^2 \cdot \Delta_{13}(\Delta_{24} + \Delta_{34}) - \Delta^2 \Delta_{13}(\Delta_{24} + \Delta_{34}) - \\
 &\quad - \bar{\Delta}^2 \Delta_{13}(\Delta_{24} + \Delta_{34}) + |\Delta|^2 \Delta_{13}(\Delta_{24} + \Delta_{34})] = \\
 &= \sin^2 \varphi \cdot \{ |\Delta|^2 \cdot [|\Delta_{32} + \Delta_{34}|^2 + |\Delta_{14} + \Delta_{34}|^2] - \\
 &\quad - \Delta^2 \cdot [(\Delta_{32} + \Delta_{34})(\Delta_{14} + \Delta_{34}) + \Delta_{13}(\Delta_{24} + \Delta_{34})] - \\
 &\quad - \bar{\Delta}^2 [(\Delta_{14} + \Delta_{34})(\Delta_{32} + \Delta_{34}) + \Delta_{13}(\Delta_{24} + \Delta_{34})] + \\
 &\quad + |\Delta|^2 [\Delta_{13}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\Delta_{24} + \Delta_{34})] \};
 \end{aligned}$$

Using the obvious formula $\Delta_{13}\Delta_{24} + \Delta_{14}\Delta_{32} = \Delta_{12}\Delta_{34}$ transform the coefficients of the resulting expression

$$\begin{aligned}
 &(\Delta_{14} + \Delta_{34})(\Delta_{32} + \Delta_{34}) + \Delta_{13}(\Delta_{24} + \Delta_{34}) = \Delta_{14}\Delta_{32} + \Delta_{14}\Delta_{34} + \Delta_{34}\Delta_{32} + \\
 &+ \Delta_{34}^2 + \Delta_{13}\Delta_{24} + \Delta_{13}\Delta_{34} = \Delta_{12}\Delta_{34} + \Delta_{14}\Delta_{34} + \Delta_{34}\Delta_{32} + \Delta_{34}^2 + \Delta_{13}\Delta_{34} = \\
 &= \Delta_{34}(\Delta_{12} + \Delta_{13} + \Delta_{14} + \Delta_{32} + \Delta_{34}) = \Delta_{34} \cdot \Delta.
 \end{aligned}$$

Thus, the value of the first determinant will be

$$\begin{aligned}
 &\sin^2 \varphi \cdot \{ |\Delta|^2 \cdot [|\Delta_{32} + \Delta_{34}|^2 + |\Delta_{14} + \Delta_{34}|^2] - \\
 &\quad - \Delta^2 \cdot \Delta_{34} \cdot \bar{\Delta} - \bar{\Delta}^2 \cdot \Delta_{34} \cdot \Delta + |\Delta|^2 [\Delta_{13}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\Delta_{24} + \Delta_{34})] \} = \\
 &\sin^2 \varphi \cdot \{ |\Delta|^2 \cdot [|\Delta_{32} + \Delta_{34}|^2 + |\Delta_{14} + \Delta_{34}|^2] - [|\Delta|^2 \Delta \cdot \Delta_{34} + \bar{\Delta} \cdot \Delta_{34} |\Delta|^2] +
 \end{aligned}$$

$$\begin{aligned}
& + |\Delta|^2 [\overline{\Delta_{13}}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\overline{\Delta_{24}} + \overline{\Delta_{34}})] \} \\
& = |\Delta|^2 \cdot \sin^2 \varphi \cdot \{ [|\Delta_{32} + \Delta_{34}|^2 + |\Delta_{14} + \Delta_{34}|^2] - \\
& - [\Delta \cdot \overline{\Delta_{34}} + \bar{\Delta} \cdot \Delta_{34}] + [\overline{\Delta_{13}}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\overline{\Delta_{24}} + \overline{\Delta_{34}})] \}.
\end{aligned}$$

Thus, the coefficient at $\sin^2 \varphi$ turned out to be a real value.

Now we calculate the second determinant from formula (6).

b)

$$\begin{aligned}
& -\cos^2 \varphi \begin{bmatrix} \Delta_{13} & -(\Delta_{12} + \Delta_{32}) & -\Delta_{13} & -(\Delta_{14} + \Delta_{34}) \\ \Delta_{12} + \Delta_{13} + \Delta_{14} & -(\Delta_{32} + \Delta_{42}) & \Delta_{32} + \Delta_{34} & -(\Delta_{34} + \Delta_{24}) \\ -\overline{\Delta_{13}} & \overline{\Delta_{32}} + \overline{\Delta_{34}} & \overline{\Delta_{13}} & \overline{\Delta_{12}} + \overline{\Delta_{14}} \\ -(\overline{\Delta_{13}} + \overline{\Delta_{14}} + \overline{\Delta_{34}}) & \overline{\Delta_{32}} + \overline{\Delta_{42}} & -(\overline{\Delta_{12}} + \overline{\Delta_{32}}) & \overline{\Delta_{12}} + \overline{\Delta_{24}} \end{bmatrix} = \\
& = -\cos^2 \varphi \begin{bmatrix} 0 & -\Delta & -\Delta_{13} & -(\Delta_{14} + \Delta_{34}) \\ \Delta & 0 & \Delta_{32} + \Delta_{34} & -(\Delta_{34} + \Delta_{24}) \\ 0 & \bar{\Delta} & \overline{\Delta_{13}} & \overline{\Delta_{12}} + \overline{\Delta_{14}} \\ -\bar{\Delta} & 0 & -(\overline{\Delta_{12}} + \overline{\Delta_{32}}) & \overline{\Delta_{12}} + \overline{\Delta_{24}} \end{bmatrix} = \\
& = -\cos^2 \varphi \left\{ -\Delta \begin{vmatrix} -\Delta & -\Delta_{13} & -(\Delta_{14} + \Delta_{34}) \\ \bar{\Delta} & \overline{\Delta_{13}} & \overline{\Delta_{12}} + \overline{\Delta_{14}} \\ 0 & -(\overline{\Delta_{12}} + \overline{\Delta_{32}}) & \overline{\Delta_{12}} + \overline{\Delta_{24}} \end{vmatrix} + \bar{\Delta} \begin{vmatrix} -\Delta & -\Delta_{13} & -(\Delta_{14} + \Delta_{34}) \\ 0 & \Delta_{32} + \Delta_{34} & -(\Delta_{34} + \Delta_{24}) \\ \bar{\Delta} & \overline{\Delta_{13}} & \overline{\Delta_{12}} + \overline{\Delta_{14}} \end{vmatrix} \right\} = \\
& = -\cos^2 \varphi \left\{ \Delta \cdot \begin{vmatrix} \Delta & \Delta_{13} & \Delta_{14} + \Delta_{34} \\ \bar{\Delta} & \overline{\Delta_{13}} & \overline{\Delta_{12}} + \overline{\Delta_{14}} \\ 0 & -(\overline{\Delta_{12}} + \overline{\Delta_{32}}) & \overline{\Delta_{12}} + \overline{\Delta_{24}} \end{vmatrix} + \bar{\Delta} \cdot \begin{vmatrix} \Delta & \Delta_{13} & \Delta_{14} + \Delta_{34} \\ \bar{\Delta} & \overline{\Delta_{13}} & \overline{\Delta_{12}} + \overline{\Delta_{14}} \\ 0 & \Delta_{32} + \Delta_{34} & -(\Delta_{24} + \Delta_{34}) \end{vmatrix} \right\} = \\
& = -\cos^2 \varphi \left\{ \Delta \cdot \left[(\overline{\Delta_{12}} + \overline{\Delta_{32}}) \begin{vmatrix} \Delta & \Delta_{14} + \Delta_{34} \\ \bar{\Delta} & \overline{\Delta_{12}} + \overline{\Delta_{14}} \end{vmatrix} + (\overline{\Delta_{12}} + \overline{\Delta_{24}}) \begin{vmatrix} \Delta & \Delta_{13} \\ \bar{\Delta} & \overline{\Delta_{13}} \end{vmatrix} \right] + \right. \\
& \quad \left. + \bar{\Delta} \left[-(\Delta_{32} + \Delta_{34}) \begin{vmatrix} \Delta & \Delta_{14} + \Delta_{34} \\ \bar{\Delta} & \overline{\Delta_{12}} + \overline{\Delta_{14}} \end{vmatrix} - (\Delta_{24} + \Delta_{34}) \begin{vmatrix} \Delta & \Delta_{13} \\ \bar{\Delta} & \overline{\Delta_{13}} \end{vmatrix} \right] \right\} = \\
& = \cos^2 \varphi \cdot \left\{ \left| \frac{\Delta}{\bar{\Delta}} \frac{\Delta_{14} + \Delta_{34}}{\Delta_{12} + \Delta_{14}} \right| \cdot [\Delta(\overline{\Delta_{12}} + \overline{\Delta_{32}}) - \bar{\Delta}(\Delta_{32} + \Delta_{34})] + \right. \\
& \quad \left. + \left| \frac{\Delta}{\bar{\Delta}} \frac{\Delta_{13}}{\Delta_{13}} \right| \cdot [\Delta(\overline{\Delta_{12}} + \overline{\Delta_{24}}) - \bar{\Delta}(\Delta_{24} + \Delta_{34})] \right\} = \\
& = \cos^2 \varphi \cdot \{ [\Delta(\overline{\Delta_{12}} + \overline{\Delta_{14}}) - \bar{\Delta}(\Delta_{14} + \Delta_{34})] \cdot [\Delta(\overline{\Delta_{12}} + \overline{\Delta_{32}}) - \bar{\Delta}(\Delta_{32} + \Delta_{34})] + \\
& \quad + (\Delta \cdot \overline{\Delta_{13}} - \Delta_{13} \cdot \bar{\Delta}) [\Delta(\overline{\Delta_{12}} + \overline{\Delta_{24}}) - \bar{\Delta}(\Delta_{24} + \Delta_{34})] \}.
\end{aligned}$$

Using the obvious formula $\Delta_{13}\Delta_{24} + \Delta_{14}\Delta_{32} = \Delta_{12}\Delta_{34}$, we transform the expression under the curly bracket.

$$\begin{aligned}
& \Delta^2 \cdot (\overline{\Delta_{12}} + \overline{\Delta_{14}})(\overline{\Delta_{12}} + \overline{\Delta_{32}}) - |\Delta|^2 (\overline{\Delta_{12}} + \overline{\Delta_{14}})(\Delta_{32} + \Delta_{34}) - |\Delta|^2 (\Delta_{14} + \Delta_{34})(\overline{\Delta_{12}} + \overline{\Delta_{32}}) + \\
& + \bar{\Delta}^2 (\Delta_{14} + \Delta_{34})(\Delta_{32} + \Delta_{34}); \tag{7}
\end{aligned}$$

$$\begin{aligned}
& \Delta^2 \cdot \overline{\Delta_{13}}(\overline{\Delta_{12}} + \overline{\Delta_{24}}) - |\Delta|^2 \overline{\Delta_{13}}(\Delta_{24} + \Delta_{34}) - \\
& - |\Delta|^2 \Delta_{13}(\overline{\Delta_{12}} + \overline{\Delta_{24}}) + \bar{\Delta}^2 \Delta_{13}(\Delta_{24} + \Delta_{34}); \tag{8}
\end{aligned}$$

$$\begin{aligned} (\overline{\Delta_{12}} + \overline{\Delta_{14}})(\overline{\Delta_{12}} + \overline{\Delta_{32}}) + \overline{\Delta_{13}}(\overline{\Delta_{12}} + \overline{\Delta_{24}}) &= \overline{\Delta_{12}^2} + \overline{\Delta_{12}} \cdot \overline{\Delta_{32}} + \overline{\Delta_{14}} \cdot \overline{\Delta_{12}} + \\ \overline{\Delta_{14}} \cdot \overline{\Delta_{32}} + \overline{\Delta_{13}} \cdot \overline{\Delta_{12}} + \overline{\Delta_{13}} \cdot \overline{\Delta_{24}} &= \\ \overline{\Delta_{12}^2} + \overline{\Delta_{12}} \cdot \overline{\Delta_{32}} + \overline{\Delta_{14}} \cdot \overline{\Delta_{12}} + \overline{\Delta_{12}} \cdot \overline{\Delta_{34}} + \overline{\Delta_{13}} \cdot \overline{\Delta_{12}} &= \overline{\Delta_{12}} \cdot \bar{\Delta}. \end{aligned} \quad (9)$$

Similarly, we have

$$\begin{aligned} (\Delta_{14} + \Delta_{34})(\Delta_{32} + \Delta_{34}) + \Delta_{13}(\Delta_{24} + \Delta_{34}) &= \Delta_{14}\Delta_{32} + \Delta_{14}\Delta_{34} + \\ \Delta_{34}\Delta_{32} + \Delta_{34}^2 + \Delta_{13}\Delta_{24} + \Delta_{13}\Delta_{34} &= \Delta_{14}\Delta_{34} + \Delta_{34}\Delta_{32} + \Delta_{34}^2 + \\ + \Delta_{12}\Delta_{34} + \Delta_{13}\Delta_{34} &= \Delta_{34} \cdot \Delta. \end{aligned} \quad (10)$$

Adding formulas (7), (8) and considering (9) and (10), we have

$$\begin{aligned} |\Delta|^2 \cdot \Delta \cdot \bar{\Delta}_{12} + |\Delta|^2 \bar{\Delta} \cdot \Delta_{34} - \\ - |\Delta|^2 [(\overline{\Delta_{12}} + \overline{\Delta_{14}})(\overline{\Delta_{32}} + \Delta_{34}) + (\Delta_{14} + \Delta_{34})(\overline{\Delta_{12}} + \overline{\Delta_{32}}) + \\ + \overline{\Delta_{13}}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\overline{\Delta_{12}} + \overline{\Delta_{24}})]. \end{aligned} \quad (11)$$

Now convert the expression under the square bracket.

$$\begin{aligned} (\overline{\Delta_{12}} + \overline{\Delta_{14}})(\overline{\Delta_{32}} + \Delta_{34}) + (\Delta_{14} + \Delta_{34})(\overline{\Delta_{12}} + \overline{\Delta_{32}}) + \overline{\Delta_{13}}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\overline{\Delta_{12}} + \overline{\Delta_{24}}) &= \\ = (\bar{\Delta} - \bar{\Delta}_{13} - \bar{\Delta}_{32} - \bar{\Delta}_{34})(\Delta_{32} + \Delta_{34}) + (\Delta - \Delta_{12} - \Delta_{32} - \Delta_{13})(\overline{\Delta_{12}} + \overline{\Delta_{32}}) + \\ + \overline{\Delta_{13}}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\overline{\Delta_{12}} + \overline{\Delta_{24}}) &= \bar{\Delta}(\Delta_{32} + \Delta_{34}) - |\Delta_{32} + \Delta_{34}|^2 - \overline{\Delta_{13}}(\Delta_{32} + \Delta_{34}) + \\ + \Delta(\overline{\Delta_{12}} + \overline{\Delta_{32}}) - |\overline{\Delta_{12}} + \overline{\Delta_{32}}|^2 - \Delta_{13}(\overline{\Delta_{12}} + \overline{\Delta_{32}}) + \overline{\Delta_{13}}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\overline{\Delta_{12}} + \overline{\Delta_{24}}) &= \\ - |\Delta_{32} + \Delta_{34}|^2 - |\overline{\Delta_{12}} + \overline{\Delta_{32}}|^2 + \overline{\Delta_{13}}(\Delta_{24} - \Delta_{32}) + \Delta_{13}(\overline{\Delta_{24}} - \overline{\Delta_{32}}) + \bar{\Delta}(\Delta_{32} + \Delta_{34}) + \\ \Delta(\overline{\Delta_{12}} + \overline{\Delta_{32}}) &= \bar{\Delta}\Delta_{32} + \Delta\bar{\Delta}_{32} + \bar{\Delta}\Delta_{34} + \Delta\bar{\Delta}_{12} + \\ + \bar{\Delta}_{13}(\Delta_{24} - \Delta_{32}) + \Delta_{13}(\overline{\Delta_{24}} - \overline{\Delta_{32}}) - |\Delta_{32} + \Delta_{34}|^2 - |\overline{\Delta_{12}} + \overline{\Delta_{32}}|^2; \end{aligned} \quad (12)$$

Substituting (12) into (1), we obtain

$$\begin{aligned} |\Delta|^2 \cdot \{|\Delta_{32} + \Delta_{34}|^2 + |\overline{\Delta_{12}} + \overline{\Delta_{32}}|^2 - \\ - [\bar{\Delta}\Delta_{32} + \Delta\bar{\Delta}_{32} + \bar{\Delta}_{13}(\Delta_{24} - \Delta_{32}) + \Delta_{13}(\overline{\Delta_{24}} - \overline{\Delta_{32}})]\}. \end{aligned}$$

Thus, the second determinant has the form:

$$\begin{aligned} -\cos^2 \varphi \{|\Delta_{32} + \Delta_{34}|^2 + |\overline{\Delta_{12}} + \overline{\Delta_{32}}|^2 - \\ - [\bar{\Delta}\Delta_{32} + \Delta\bar{\Delta}_{32} + \bar{\Delta}_{13}(\Delta_{24} - \Delta_{32}) + \Delta_{13}(\overline{\Delta_{24}} - \overline{\Delta_{32}})]\} |\Delta|^2. \end{aligned}$$

Adding this formula to the first determinant, we obtain the equations for determining φ .

$$\begin{aligned} |\Delta|^2 \sin^2 \varphi \{[|\Delta_{32} + \Delta_{34}|^2 + |\Delta_{14} + \Delta_{34}|^2] - \\ - [\Delta \cdot \overline{\Delta_{34}} + \bar{\Delta} \cdot \Delta_{34}] + [\overline{\Delta_{13}}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\overline{\Delta_{24}} + \overline{\Delta_{34}})]\} - |\Delta|^2 \cdot \cos^2 \varphi \{|\Delta_{32} + \Delta_{34}|^2 + \\ + |\overline{\Delta_{12}} + \overline{\Delta_{32}}|^2 - [\bar{\Delta}\Delta_{32} + \Delta\bar{\Delta}_{32} + \bar{\Delta}_{13}(\Delta_{24} - \Delta_{32}) + \Delta_{13}(\overline{\Delta_{24}} - \overline{\Delta_{32}})]\} = 0. \end{aligned}$$

According to our assumption $\Delta \neq 0$, so we can reduce this value, and as a result we have

$$\begin{aligned} \sin^2 \varphi - \{|\Delta_{32} + \Delta_{34}|^2 + |\overline{\Delta_{12}} + \overline{\Delta_{32}}|^2 - \\ - [\bar{\Delta}\Delta_{32} + \Delta\bar{\Delta}_{32} + \bar{\Delta}_{13}(\Delta_{24} - \Delta_{32}) + \Delta_{13}(\overline{\Delta_{24}} - \overline{\Delta_{32}})]\} \cdot \cos^2 \varphi = 0. \end{aligned}$$

Theorem 3.1. If φ is the solution to the equation

$$\begin{aligned} \{[|\Delta_{32} + \Delta_{34}|^2 + |\Delta_{14} + \Delta_{34}|^2] - [\Delta \cdot \overline{\Delta_{34}} + \bar{\Delta} \cdot \Delta_{34}] + \\ + [\overline{\Delta_{13}}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\overline{\Delta_{24}} + \overline{\Delta_{34}})]\} \cdot \sin^2 \varphi - \\ \{|\Delta_{32} + \Delta_{34}|^2 + |\overline{\Delta_{12}} + \overline{\Delta_{32}}|^2 + \\ + [\bar{\Delta}\Delta_{32} + \Delta\bar{\Delta}_{32} + \bar{\Delta}_{13}(\Delta_{24} - \Delta_{32}) + \Delta_{13}(\overline{\Delta_{24}} - \overline{\Delta_{32}})]\} \cdot \cos^2 \varphi = 0, \end{aligned} \quad (13)$$

that is the formula

$$TL = L^+ T^*,$$

————— 92 —————

where

$$T = i \cos \varphi I + \sin \varphi S, \quad Sy(x) = y(1-x), \quad (14)$$

and I – is the identity operator.

Theorem 3.2.

If φ – is the solution of equation (13), then there are formulas

1) $(TL)^2 = L^*L$;

2) $(LT)^2 = LL^*$;

where L – is the Sturm-Liouville operator,

$$T = i \cos \varphi I + \sin \varphi S, \quad Sy(x) = y(1-x), \quad (14)$$

and I – is the identity operator.

4. Discussions.

If

a) $\{[|\Delta_{32} + \Delta_{34}|^2 + |\Delta_{14} + \Delta_{34}|^2] [\Delta \cdot \overline{\Delta_{34}} + \bar{\Delta} \cdot \Delta_{34}] + [\overline{\Delta_{13}}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\overline{\Delta_{24}} + \overline{\Delta_{34}})]\} = 0,$
 $\{|\Delta_{32} + \Delta_{34}|^2 + |\overline{\Delta_{12}} + \overline{\Delta_{32}}|^2 -$
 $-[\bar{\Delta}\Delta_{32} + \Delta\bar{\Delta}_{32} + \bar{\Delta}_{13}(\Delta_{24} - \Delta_{32}) + \Delta_{13}(\overline{\Delta_{24}} - \overline{\Delta_{32}})]\} \neq 0,$

that $\cos \varphi = 0$ and the operator T takes the form $T = S$, where

$$Sy(x) = y(1-x);$$

b) $\{[|\Delta_{32} + \Delta_{34}|^2 + |\Delta_{14} + \Delta_{34}|^2] [\Delta \cdot \overline{\Delta_{34}} + \bar{\Delta} \cdot \Delta_{34}] + [\overline{\Delta_{13}}(\Delta_{24} + \Delta_{34}) + \Delta_{13}(\overline{\Delta_{24}} + \overline{\Delta_{34}})]\} \neq 0 \cdot$
 $\{|\Delta_{32} + \Delta_{34}|^2 + |\overline{\Delta_{12}} + \overline{\Delta_{32}}|^2 - [\bar{\Delta}\Delta_{32} + \Delta\bar{\Delta}_{32} + \bar{\Delta}_{13}(\Delta_{24} - \Delta_{32}) + \Delta_{13}(\overline{\Delta_{24}} - \overline{\Delta_{32}})]\} = 0, \quad \text{to}$
 $\sin \varphi = 0$, and the operator T takes the form $T = il$, where I – is the unit operator.

In the first case, the closure of the SL operator is self-adjoint, and in the second case, the operator L is skew-symmetric, therefore its closure is the normal operator.

5. The authors are grateful to Anton Selitsky for his advice on the Kato hypothesis.

УДК 517.43

А.Ш. Шалданбаев¹, А.Б. Иманбаева²,
 А.Ж. Бейсебаева³, А.А. Шалданбаева⁴

¹“Silkway” Халықаралық Университеті, Шымкент;

^{2,3}М.Әуезов атындағы Оңтүстік Қазақстан Мемлекеттік Университеті, Шымкент;

⁴Аймақтық әлеуметтік - инновациялық университеті, Шымкент.

ТӨРТИНШІ РЕТТІ ШТУРМ-ЛИУВИЛЛ ОПЕРАТОРЫНЫҢ КВАДРАТ ТҮБІРІ ТУРАЛЫ

Аннотация. Бұл еңбекте Штурм-Лиувиллдің оң төртінші ретті операторынан квадрат түбір табылды. Ол қайтымды Штурм-Лиувилл операторы мен оның сынарының композициясы немесе көбейтіндісі. Табылған түбір оң емес, бірақ тегі жалқы операторлар санатына жатады. Бағдарлаушы ретінде Путнамның бір алгебралық теоремасы қолданылды. Зерттеу нәтижелері операторлардың спектрлі теориясы мен теориялық физикада қолданыс табады деп қүтілуде.

Түйін сөздер. Катоның гипотезасы, диссипативті оператор, оператордың квадрат түбірі, Путнамның теоремасы, ауытқыған аргумент, оператордаң бөлшек түбірлері, кері есептер, спектр, жалқы оператор, оң операторлар, функционел-дифференциал оператор, спектрлі теория.

**А.Ш. Шалданбаев¹, А.Б. Иманбаева²,
А.Ж. Бейсебаева³, А.А. Шалданбаева⁴**

¹Международный университет “Silkway”, Шымкент;

^{2,3}Южно-Казахстанский государственный университет имени М.Ауезова, Шымкент;

⁴Региональный социально-инновационный университет, Шымкент.

О КВАДРАТНОМ КОРНЕ ИЗ ОПЕРАТОРА ШТУРМА-ЛИУВИЛЯ ЧЕТВЁРТОГО ПОРЯДКА

Аннотация. В настоящей работе найден корень из положительного оператора Штурма - Лиувилля четвертого порядка, являющегося композицией обратимого оператора Штурма - Лиувилля и его сопряженного. Найденный корень не обладает свойством положительности, но является самосопряженным оператором в существенном. В качестве наводящей идеи использована одна теорема Путнама алгебраического характера. Можно надеяться, что результаты работы найдут приложения в спектральной теории операторов и теоретической физике.

Ключевые слова: гипотеза Като, диссипативный оператор, квадратный корень из оператора, теорема Путнама, отклоняющиеся аргумент, дробные степени оператора, обратная задача, спектр, унитарный оператор, самосопряженный оператор, положительный оператор, функционально-дифференциальный оператор, спектральная теория.

Information about authors:

Shaldanbayev A.Sh. – doctor of physico-mathematical Sciences, Professor, head of the center for mathematical modeling «Silkway» International University, Shymkent; <http://orcid.org/0000-0002-7577-8402>;

Imanbayeva A.B. - M.Auezov South Kazakhstan State University, Shymkent, Candidate of physical and mathematical Sciences, associate Professor; <https://orcid.org/0000-0002-8203-7637>;

Beisebayeva A.Zh. - M.Auezov South Kazakhstan State University, Shymkent; <https://orcid.org/0000-0003-4839-9156>;

Shaldanbayeva A.A. - Regional Social-Innovative University, Shymkent; <https://orcid.org/0000-0003-2667-3097>

REFERENCES

- [1] U. Rudin, Functional Analysis, M.: Mir, **1975**.
- [2] C.R. Putnam, Amer.Math., 73 (**1951**), 357-362.
- [3] T. Kato, Fractional powers of dissipative operators, II, J. Math. Soc. Japan, 14 (**1962**), pp. 242–248.
- [4] A. McIntosh, On the compatibility of $A^{1/2}$ and $A^{*1/2}$, Proc. Amer. Math. Soc., 32:2 (**1972**), pp. 430–434.
- [5] A.M. Selitskiy, Initial data space of the second boundary value problem for a parabolic differential-difference equation, Scientific Bulletin of BelsSU, Series: Mathematics. Physics. (**2011**). № 23(118). Issue 25, pp. 102-112.
- [6] M.S. Agranovich and A.M. Selitskiy, Fractional degrees of operators corresponding to coercive problems in Lipschitz domains, Funct. Analysis and its Appl., (**2013**), volume 47, issue 2, pp.2–17.
- [7] V. Balakrishnan (**1960**), Fractional powers of closed operators and the semi-groups generated by them. Pacific J. Math., 10:419–437.
- [8] Andreas Axelsson, Stephen Keith, and Alan McIntosh (**2006**), The Kato square root problem for mixed boundary value problems. J. London Math. Soc. (2), 74(1):113–130.
- [9] Andrea Bonito and Joseph E. Pasciak (**2015**), Numerical approximation of fractional powers of elliptic operators. Math. Comp., 84(295):2083–2110.
- [10] Luis Caffarelli and Luis Silvestre (**2007**), An extension problem related to the fractional Laplacian. Comm. Partial Differential Equations, 32(7-9):1245–1260.
- [11] Alan McIntosh (**1989**), The square root problem for elliptic operators: a survey. In Functional analytic methods for partial differential equations, Tokyo, volume 1450 of Lecture Notes in Math., pages 122–140. Springer, Berlin, 1990.
- [12] M. S. Agranovich (**2012**), Remarks on strongly elliptic second-order systems in Lipschitz domains, Russian J. Math. Phys., 20:4, 405–416.
- [13] M.S. Agranovich (**2013**), Sobolev spaces, their generalization and elliptic problems in domains with smooth and Lipschitz bound, M., Izd. MSNMO.
- [14] Mihoko Matsuki and Teruo Ushijima (**1993**), A note on the fractional powers of operators approximating a positive definite selfadjoint operator. J. Fac. Sci. Univ. Tokyo Sect. IA Math., 40(2):517–528.

- [15] Y. Berg and Y. Lefstrem (1980), Interpolation Spaces, Mir, M.
- [16] T. Kato (1972), Perturbation Theory of Linear Operators, Mir, M.
- [17] M.A. Krasnoselsky, P.P. Zabreiko, E.I. Pumulnik and P.E. Sobolevsky (1966), Integral operators in spaces of summable functions, Nauka, M.
- [18] J.-L. Lyons and E. Madjenes (1971), Nonhomogeneous Boundary Value Problems and their Applications, Mir, M.
- [19] A. M. Selitsky, Space of initial data of the 3rd boundary value problem for parabolic differential-difference equation in one-dimensional case, Math.Notes, 92: 4 (2012), 636–640.
- [20] A.M. Selitsky, Modeling of some optical systems on the basis of a parabolic differential-difference equation, Math. Modeling, 24:12 (2012), 38–42.
- [21] A. L. Skubachevsky and R.V. Shamin, Second-order parabolic differential-difference equations, Dokl. RAN, 379:5 (2001), 595–598.
- [22] H. Triebel, Theory of Functional Spaces, Mir, M., 1986.
- [23] I. Ya. Shneiberg, Spectral properties of linear operators in interpolational families of Banach spaces, Math. investig. 9:2 (1974), 214–229.
- [24] Sh. Agmon, On the eigenfunctions and on the eigenvalues of general elliptic boundary value problems, Comm. Pure Appl. Math., 15:2 (1962), 119–147.
- [25] W. Arendt, Semigroups and evolution equations: functional calculus, regularity and kernel estimates, in: Handbook of Differential Equations, Evolutionary Differential Equations, vol. 1, Elsevier/North-Holland, Amsterdam, 2004, 1–85.
- [26] P. Auscher, N. Badr, R. Haller-Dintelmann, J. Rehberg, The square root problem for second order, divergence form operators with mixed boundary conditions on L_p , <http://arxiv.org/abs/1210.0780v1>.
- [27] P. Auscher, S. Hofmann, M. Lacey, J. Lewis, A. McIntosh, P. Tchamitchian, The solution of Kato's conjectures, C. R. Acad. Sci. Paris, Sér. 1, 332:7 (2001), 601–606.
- [28] P. Auscher, A. McIntosh, A. Nahmod, Holomorphic functional calculi of operators, quadratic estimates and interpolation, Indiana Univ. Math. J., 46:2 (1997), 375–403.
- [29] P. Auscher, S. Hofmann, A. McIntosch, P. Tchamitchian, The Kato square root problem for higher order elliptic operators and systems on R^n , J. Evol. Equ., 1:4 (2001), 361–385.
- [30] P. Auscher, P. Tchamitchian, Square root problem for divergence operators and related topics, Astérisque, 249 (1998), 1–171.
- [31] P. Auscher, P. Tchamitchian, Square roots of elliptic second order divergence operators on strongly Lipschitz domains: L_2 theory, J. Anal. Math., 90 (2003), 1–12.
- [32] P. Auscher, P. Tchamitchian, Square roots of elliptic second order divergence operators on strongly Lipschitz domains: L_p theory, Math. Ann., 320:3 (2001), 577–623.
- [33] A. Axelsson, S. Keith, A. McIntosh, The Kato square root problem for mixed boundary value problems, J. London Math. Soc., 74:1 (2006), 113–130.
- [34] S. Blunck, P. Kunstmann, Calderron–Zygmund theory for non-integral operators and the H^∞ functional calculus, Rev. Mat. Iberoamericana, 19:3 (2003), 919–942.
- [35] A.F. M. ter Elst, D.W. Robinson, On Kato's square root problem, Hokkaido Math. J., 26:2 (1997), 365–376.
- [36] J. Griepentrog, K. Gröger, H.-Ch. Kaiser, J. Rehberg, Interpolation for function spaces related to mixed boundary value problems, Math. Nachr., 241 (2002), 110–120.
- [37] D. Grisvard, Caractérisation de quelques espaces d'interpolation, Arc. Rational Mech. Anal., 25 (1967), 40–63.
- [38] M. Haase, (2006), The Functional Calculus for Sectorial Operators, Birkhäuser, Basel.
- [39] S. Hoffmann, A short course on the Kato problem, Contemp. Math., 289 (2001), 61–67.
- [40] T. Hytönen, A. McIntosh and P. Portal, Kato's square root problem in Banach spaces, J. Funct. Anal., 254:3 (2008), 675–726.
- [41] S. Janson, P. Nilsson, J. Peetre, Notes on Wolff's note on interpolation spaces, Proc. London Math. Soc. (3), 48:2 (1984), 283–299.
- [42] H. Komatsu, Fractional powers of operators, Pacif. J. Math., 19 (1966), 285–346.
- [43] J. L. Lions (1961), Equations différentielles opérationnelles et problèmes aux limites, Springer-Verlag, Berlin etc.
- [44] J. L. Lions, Espaces d'interpolation et domaines de puissances fractionnaires d'opérateurs, J. Math. Soc. Japan, 14 (1962), 233–241.
- [45] A. McIntosh, Square roots of elliptic operators, J. Funct. Anal., 61:3 (1985), 307–327.
- [46] A. McIntosh, Operators which have an H^∞ functional calculus, in: Miniconference on Operator Theory and Partial Differential Equations, Proc. Centre Math. Anal.Austral. Nat. Univ., vol. 14, Austral. Nat. Univ., Canberra, (1986), 210–231.
- [47] A. McIntosh (1990), Square Root Problem for Elliptic Operators: a Survey, Lecture Notes in Math., vol. 1450, Springer-Verlag, Berlin.
- [48] W. McLean (2000), Strongly Elliptic Systems and Boundary Integral Equations, Cambridge Univ. Press, Cambridge, UK.

- [49] J. Nečas (2012), *Les méthodes directes en théorie des équations elliptiques*, Masson, Paris, 1967; *Direct Methods in the Theory of Elliptic Equations*, Springer-Verlag, Berlin–Heidelberg.
- [50] L. Nirenberg, Remarks on strongly elliptic partial differential equations, *Comm. Pure Appl. Math.*, 8 (1965), 649–675.
- [51] R.T. Seeley, Norms and domains of the complex powers $A^z B$, *Amer. J. Math.*, 93:2(1971), 299–309.
- [52] R.T. Seeley, Interpolation in L_p with boundary conditions, *Studia Math.*, 44 (1972), 47–60.
- [53] A.L. Skubachevskii and R. V. Shamin, Mixed boundary value problem for parabolic differential-difference equation, *Funct. Differ. Eq.*, 8:3–4 (2001), 407–424.
- [54] T.W. Wolff, A note on interpolation spaces, in: *Lecture Notes in Math.*, vol. 918, Springer-Verlag, Berlin–New York, (1982), 199–204.
- [55] A. Yagi, Coincidence entre des espaces d’interpolation et des domaines de puissances fractionnaires d’opérateurs, *C. R. Acad. Sci. Paris, Sér. I*, 299:6 (1984), 173–176.
- [56] L. Hörmander (1965), *Linear Partial Differential Operators*, Mir, M.
- [57] M.I. Vishik, On strongly elliptic systems of differential equations, *Math. Sb.*, 29(71), 3(1951), 615–676.
- [58] R.V. Shamin, On spaces of initial data for differential equations in a Hilbert space, *Math. Sb.*, 194:9 (2003), 141–156.
- [59] H. Namsrai, Square Klein-Gordon operator and physical interpretation, *International Journal of Theoretical Physics*, (1998). T. 37. №5. pp.1531-1540.
- [60] R. Putsio (1994), On square root of Laplace – Beltrami operator as Hamiltonian, *Classical and Quantum Gravity*. Vol. 11. No.3. Pp. 609-620.
- [61] T.L. Gill and V.V. Zakhari, Analytical representation of square operator, *Physical Journal A: Mathematics and general.* (2005). Vol. 38. №11. Pp. 2479-2496.
- [62] P.N. Vabishchevich, Numerical solution of non-stationary spatial-fractional problems with square root of an elliptic operator, *Mathematical Modeling and Analysis*. (2016). Vol. 21. No.2. Pp. 220-238.
- [63] A. Bzdak and L. Hadash, Square root of a dirac operator on superspace and Maxwell equations, *Physical Letters. Section B: Nuclear Physics, Elementary Particle and High Energy Physics.* (2004). vol. 582. №1-2. Pp. 113-116.
- [64] H.T. Ito, Resonances of square root of the Pauli operator, *Publications of Research Institute of Mathematical Sciences.* (2017). Vol. 53. No.4. Pp. 517-549.
- [65] H. Namsray and H.V. Von Geramb, Quantization and nonlocality of square body: Review of *International Journal of Theoretical Physics*. (2001). vol. 40. № 11. pp. 1929-2010.
- [66] V.P. Maslov (1988), *Asymptotic Methods and Perturbation Theory*, M.: Mir.
- [67] V.A. Marchenko (1977), *Sturm-Liouville Operators and Their Applications*, Kiev, Naukova Dumka.
- [68] T.Sh. Kalmenov, S.T. Akhmetova and A.Sh. Shaldanbaev, To spectral theory of equation with deviating arguments, *Mathematical Journal, Almaty*, (2004), vol.4., No.3, pp.41-48.
- [69] M.I. Akylbayev, A.Zh. Beysebayeva and A.Sh. Shaldanbayev, News of the National Academy of Sciences of the Republic of Kazakhstan, Physico-mathematical Series, Volume 1, Number 317 (2018), 34 – 50.
- [70] T.Sh. Kal'menov and A.Sh. Shaldanbaev, On a criterion of solvability of the inverse problem of heat conduction, *Journal of Inverse and Ill-Posed Problems* 18, 352-369 (2010).
- [71] I. Orazov, A. Shaldanbayev and M. Shomanbayeva, About the Nature of the Spectrum of the Periodic Problem for the Heat Equation with a Deviating Argument, *Abstract and Applied Analysis* №:128363 DOI: 10.1155/2013/128363 Published: (2013), WOS:000325557100001.

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

www.nauka-nanrk.kz

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы *M. С. Ахметова, Т.А. Апендиев, Д.С. Аленов*
Верстка на компьютере *A.M. Кульгинбаевой*

Подписано в печать 10.06.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
8,3 п.л. Тираж 300. Заказ 3.

*Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19*