

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Әль-фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
Әль-Фараби атындағы
Қазақ ұлттық университетінің

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-farabi kazakh
national university

SERIES
PHYSICO-MATHEMATICAL

3 (325)

MAY - JUNE 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р ы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі **Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев У.У. проф. корр.-мүшесі (Қазақстан)
Жүсіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошқаев К.А. PhD докторы (Қазақстан)
Сұраған Д. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қырғыстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорус)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.
Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жантаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф. чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

E d i t o r i n c h i e f
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d:

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskiy I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.26>

Volume 3, Number 325 (2019), 71 – 76

UDC 539.14

**A.D. Duisenbay¹, N.ZH. Takibayev¹,
 V.S. Vasilevsky², V.O. Kurmangaliyeva¹, E.M. Akzhigitova¹**

¹Kazakh National University named after al-Farabi, Almaty, Kazakhstan;
 Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine
aknurka__93@mail.ru, venera_baggi@mail.ru, kzo1994@mail.ru

FORM FACTORS AND DENSITY DISTRIBUTIONS OF PROTONS AND NEUTRONS IN ⁷Li AND ⁷Be

Abstract. Form factors and density distribution of protons and neutrons in ⁷Li and ⁷Be are investigated within a microscopic two-cluster model. The model correctly treat the Pauli principle and make use of the oscillator basis to expand wave function of two-cluster system. Dynamics of two-cluster system is totally governed by a semirealistic nucleon-nucleon potential. We demonstrate that the model used correctly reproduces form factors for the ground state of ⁷Li and ⁷Be.

Key words: two-cluster system, form factors, density distribution, nucleon-nucleon potential etc.

Introduction

In this paper we consider the form factor of the elastic scattering of electron and density distribution of protons and neutrons in light nuclei ⁷Li and ⁷Be. In previous paper [1] we have investigated properties of light nuclei ⁵He, ⁵Li, ⁶Li, ⁷Li, ⁷Be and ⁸Be within a microscopic two-cluster model. This model is the algebraic version of the resonating group method. In Ref. [1] main attention was paid to the spectrum and wave functions of bound and resonance states in these nuclei, and also to phase shifts of the elastic two-cluster scattering. It was shown in Ref. [1] that the two-cluster model used correctly reproduces the main properties of light nuclei ⁵He, ⁵Li, ⁶Li, ⁷Li, ⁷Be and ⁸Be. The aim of the present investigations is to obtain complementary information on structure of bounds states in light nuclei ⁷Li and ⁷Be. This

Table 1 - Main input parameters of calculations and energy of the ground state in ⁷Li and ⁷Be.

Nucleus	Potential	b, fm	m/u	f_{LS}	E_{GS}, MeV
⁷ Li	MP	1.3451	0.969	1.00	-2.468
	MHNP	1.362	0.0002	1.00	-2.467
⁷ Be	MP	1.3451	0.969	1.00	-1.630
	MHNP	1.362	0.0002	1.00	-1.588

will be done within the same two-cluster model. It is well known that the form factor provides information on distribution of charge in a nucleus. Values of the form factor in the region of small transferred momenta allows one to determine the proton root-mean-square radius.

Details of calculations and results

As in previous paper [1], we represent ⁷Li and ⁷Be as two-cluster system $\alpha+t$ and $\alpha+^3\text{He}$, respectively. We use a common oscillator length b to describe distribution of nucleons in each cluster.

Two semirealistic nucleon-nucleon interactions: the Minnesota potential (MP) [2], [3] and the modified Hasegawa-Nagata potential (MHNP) [4, 5, 6] are employed in our calculations. As in Ref. [1] we select the oscillator length b to minimize the two-cluster threshold. We also slightly modified the Majorana parameter m of MHNP and the parameter u of MP to reproduce the energy of the ground state accounted from the two-

cluster threshold. In Table 1 we display all input parameters of our calculations. In all our calculations we make of 200 oscillator functions. This number of functions guarantees a good precision for the energy and wave functions of bound states, even for weakly bound states. We have also checked that this number of oscillator functions provides form factors and densities distributions with acceptable precision.

Since we consider light nuclei ${}^7\text{Li}$ and ${}^7\text{Be}$ within two-cluster microscopic model, wave function of bound and continuous spectrum states is represented in the following form:

$$\Psi_{EJ\pi} = \hat{A} \left\{ \left[\Phi_1(A_1, s_1) \Phi_2(A_2, s_2) \right]_S \psi_{ELS}^{J\pi}(q) Y_L(\hat{q}) \right\}_J \quad (1)$$

Notations which we are using here are identical to those of Ref. [1].

Since wave functions $\Phi_1(A_1, s_1)$ and $\Phi_2(A_2, s_2)$, describing internal motion of nucleons inside each cluster are known and fixed, thus we have to find the inter-cluster function $\psi_{ELS}^{J\pi}(q)$ by solving the dynamic equations of the resonating group method (RGM). In the standard version of the RGM, they are the integro-differential equation. In the algebraic version of RGM, which we employ, the dynamic equations are transformed into a set of linear algebraic equations. This is achieved by using a full set of the radial part of oscillator functions $\{\Phi_{nL}(q, b)\}$. By expanding the inter-cluster function $\psi_{ELS}^{J\pi}(q)$ over oscillator functions

$$\psi_{ELS}^{J\pi}(q) = \sum_{n=0}^{\infty} C_{nL;SJ} \Phi_{nL}(q, b) \quad (2)$$

or the total two-cluster function Ψ_{EJ} over cluster oscillator functions $\{nL; SJ\}$

$$\Psi_{EJ} = \sum_{n=0}^{\infty} C_{nL;SJ} |nL; SJ\rangle, \quad (3)$$

we arrive to a system of linear algebraic equations

$$\sum_{n=0}^{\infty} \{ \langle nL | \hat{H} | \tilde{n}L \rangle - E \langle nL | \tilde{n}L \rangle \} C_{nL;SJ} = 0. \quad (4)$$

In real calculations, the infinite set of equations (4) can be reduced to a finite set of equations. And thus to find energies and wave functions of bound state, we need to obtain eigenvalues and corresponding eigenfunctions of the $N \times N$ matrix

$$\| \langle nL | \hat{H} | \tilde{n}L \rangle \|_N.$$

It is worthwhile noticing that, by assuming that the wave function $\Psi_{EJ\pi}$ of a bound state is obtained, one has to calculate the matrix element

$$F_p(q) = \langle \Psi_{EJ\pi} | \hat{F}_p | \Psi_{EJ\pi} \rangle \quad (5)$$

$$F_n(q) = \langle \Psi_{EJ\pi} | \hat{F}_n | \Psi_{EJ\pi} \rangle \quad (6)$$

in order to determine the proton and neutron form factors, and one has also to calculate matrix elements

$$D_p(\mathbf{r}) = \left\langle \Psi_{EJ\pi} \left| \frac{1}{2} \sum_{i=1}^A (1 + \hat{t}_{iz}) \delta(\mathbf{r} - \mathbf{r}_i) \right| \Psi_{EJ\pi} \right\rangle, \quad (7)$$

$$D_n(\mathbf{r}) = \left\langle \Psi_{EJ\pi} \left| \frac{1}{2} \sum_{i=1}^A (1 - \hat{t}_{iz}) \delta(\mathbf{r} - \mathbf{r}_i) \right| \Psi_{EJ\pi} \right\rangle, \quad (8)$$

to obtain the density distribution of protons $D_p(\mathbf{r})$ and neutrons $D_n(\mathbf{r})$. Here the operators \hat{F}_p and \hat{F}_n are

$$\hat{F}_p = \frac{1}{2} \sum_{i=1}^A (1 + \hat{t}_{iz}) \exp \{ i(\mathbf{q}\mathbf{r}_i) \} \quad (9)$$

$$\hat{F}_n = \frac{1}{2} \sum_{i=1}^A (1 - \hat{t}_{iz}) \exp \{ i(\mathbf{q}\mathbf{r}_i) \} \quad (10)$$

and the operators $\frac{1}{2}(1 + \hat{t}_{iz})$ and $\frac{1}{2}(1 - \hat{t}_{iz})$ are projection operators on proton and neutron state, respectively. In equations (5), (6), (7) and (8), integration is carried out over all spatial, spin and isospin coordinates.

It is worth noticing that by calculating both proton and neutron form factor or densities we can easily construct the matter form factors or densities from the relations

$$F_m(q) = F_p(q) + F_n(q)$$

$$D_m(\mathbf{r}) = D_p(\mathbf{r}) + D_n(\mathbf{r})$$

Recall that the nucleon densities are normalized by the conditions

$$\int d\mathbf{r}D_p(\mathbf{r}) = Z, \quad \int d\mathbf{r}D_n(\mathbf{r}) = N, \quad \int d\mathbf{r}D_m(\mathbf{r}) = A,$$

where Z is the total number of protons, N is the total number of neutrons, and the mass number $A=Z+N$.

Form factor

In section we consider form factor of elastic scattering of electrons.

The elastic form factors for ground $3/2^-$ states of ${}^7\text{Li}$ and ${}^7\text{Be}$ are presented in Fig. 1. The form factors are calculated with MHNPF.




Figure 1 - Form factors of the elastic electron scattering from the ground states of ${}^7\text{Li}$ and ${}^7\text{Be}$.




Figure 2 - Form factor $F(q)$ of elastic electron scattering on the first excited $1/2^-$ states in ${}^7\text{Li}$ and ${}^7\text{Be}$.

In Fig. 2 we display form factor of electron elastic scattering from the excited $1/2^-$ states in ${}^7\text{Li}$ and ${}^7\text{Be}$. These results are also obtained with MHNP.

On both figures 1 and 2 we use the logarithmic scale and indicate only an exponent of $\log(F(q))$. We can see that the form factors for the ground and first excited states rapidly turn to the zero with increasing of the momentum transfer q . The form factor for the ground state of ${}^7\text{Li}$ has a node at $q=2.49 \text{ fm}^{-1}$, while for ${}^7\text{Be}$ the first node appear at $q=2.03 \text{ fm}^{-1}$. For the first excited $1/2^-$ states, the position of a node slightly shifted to small values of the momentum transfer q and equals to $q=2.26 \text{ fm}^{-1}$ for ${}^7\text{Li}$ and equals to $q=1.86 \text{ fm}^{-1}$ for ${}^7\text{Be}$.

The present results are in agreement with other microscopic model. For example, the form factors for the ground state of ${}^7\text{Li}$ and ${}^7\text{Be}$, presented above, are very close to those obtained in Refs. [7], [8], [9] within the other realization of the resonating group method. There is also fairly good agreement with the experimental data Ref. [10].

Density distributions

Now we turn our attention to the density distribution of proton and neutrons. The proton and neutron density distributions for the ${}^7\text{Li}$ ground states are shown in Fig. 3. They are calculated with MHNP. As we see the protons are mainly concentrated at small distances, while the neutron density distribution is more dispersed in the space.

In Figure 4 we display the proton and neutron density distribution for the ground state of ${}^7\text{Be}$. This nucleus is mirror to the nucleus ${}^7\text{Li}$, and thus we have inverse picture for proton and neutron density distributions with respect to Fig. 3. Despite that the energy of the ${}^7\text{Be}$ is smaller than the ground state energy of ${}^7\text{Li}$, both nuclei have similar density distributions of proton and neutron.

Conclusion

We have applied a two-cluster microscopic model to study density distribution of protons and neutrons in light atomic nuclei ${}^7\text{Li}$ and ${}^7\text{Be}$. We also have studied the form factors of elastic electron scattering from these nuclei. The microscopic model which has been employed is the algebraic version of the resonating group method. The later is using the full set of oscillator functions to expand wave functions of bound and scattering states. Nucleon-nucleon interaction being a key ingredient of any microscopic model was represented by two semirealistic potentials which are often used in two- and three-cluster microscopic models.

It was demonstrated that the present two-cluster model reproduces fairly well the elastic form factors of ${}^7\text{Li}$ and ${}^7\text{Be}$ as a function of the momentum transfer. It was also demonstrated that our results are compatible with results of other theoretical models.




Figure 3 - The proton and neutron density distribution in the ground state of ${}^7\text{Li}$.




Figure 4 - The proton and neutron density distributions in the ground state of ${}^7\text{Be}$

Acknowledgment

One of the authors (V.V.) is grateful to the members of the subdepartment of theoretical and nuclear physics from the Physical and Technical Department, al-Farabi Kazakh National University, Almaty, Republic of Kazakhstan, for hospitality and stimulating discussion during his stay at al-Farabi Kazakh National University.

This work was supported in part by the Program of Fundamental Research of the Physics and Astronomy Department of the National Academy of Sciences of Ukraine (Project No. 0117U000239) and by the Ministry of Education and Science of the Republic of Kazakhstan, Research Grant IRN: AP 05132476.

А.Д. Дүйсенбай¹, Н.Ж. Такибаев¹, В.С. Василевский²,
В.О. Құрманғалиева¹, Е.М. Ақжігітова¹

¹Әл-Фараби атындағы Қазақ Ұлттық Университеті, Алматы, Қазақстан;

²Боголюбов атындағы Теориялық Физика Институты, Киев, Украина

${}^7\text{Li}$ ЖӘНЕ ${}^7\text{Be}$ ЯДРОЛАРЫНДАҒЫ ПРОТОНДАР МЕН НЕЙТРОНДАР ТЫҒЫЗДЫҒЫНЫҢ ТАРАЛУЫ МЕН ФОРМ ФАКТОРЛАРЫ

Аннотация. ${}^7\text{Li}$ және ${}^7\text{Be}$ ядроларындағы нейтрондар мен протондар тығыздығының таралуы мен форм факторлары микроскопиялық екі кластерлік модель ретінде зерттелінді. Бұл модель Паули принципін ескере отырып, екі кластерлік жүйенің толқындық функциясын жіктеу үшін осцилляторлық негізді қолданады. Екі кластерлік жүйенің динамикасы толығымен жартылай реалистік нуклон-нуклондық потенциалымен анықталады. Қолданылып отырған модель ${}^7\text{Li}$ және ${}^7\text{Be}$ ядроларының негізгі күйлері үшін форм-факторды дұрыс беретіндігін көрсетеміз.

Түйін сөздер: екі кластерлік жүйе, форм факторлар, тығыздықтың таралуы, нуклон-нуклондық потенциал және т.б.

А.Д. Дуйсенбай¹, Н.Ж. Такибаев¹, В.С. Василевский²,
В.О. Курмангалиева¹, Э.М. Акжигитова¹

¹Казахский национальный университет имени аль-Фараби, Алматы, Казахстан;

²Институт Теоретической Физики имени Боголюбова, Киев, Украина

ФОРМ ФАКТОРЫ И РАСПРЕДЕЛЕНИЕ ПЛОТНОСТИ ПРОТОНОВ И НЕЙТРОНОВ В ЯДРАХ ⁷Li И ⁷Be

Аннотация. Форм факторы и распределение плотности протонов и нейтронов в ядрах ⁷Li и ⁷Be исследованы в рамках микроскопической двухкластерной модели. Модель правильно учитывает принцип Паули и использует осцилляторную основу для разложения волновой функции двухкластерной системы. Динамика двухкластерной системы полностью определяется полуреалистическим нуклон-нуклонным потенциалом. Мы показываем, что используемая модель правильно воспроизводит форм-факторы для основного состояния ⁷Li и ⁷Be.

Ключевые слова: двухкластерная система, форм факторы, распределение плотности, нуклон-нуклонный потенциал и т.д.

Information about authors:

Duisenbay A.D. - al-Farabi Kazakh National University, Almaty, Kazakhstan, 1st course PhD student, aknurka_93@mail.ru, ORCID: <https://orcid.org/0000-0003-3881-3958>;

Takibayev N.Zh. - al-Farabi Kazakh National University, Almaty, Kazakhstan, d.ph.-m.sc., professor, academic of NAS RK, takibayev@gmail.com, ORCID: <https://orcid.org/0000-0002-2604-6838>;

Vasilevsky V.S. - Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine, professor, vsvasilevsky@gmail.com, ORCID: <https://orcid.org/0000-0003-0417-5978>;

Kurmangaliyeva V.O. - al-Farabi Kazakh National University, Almaty, Kazakhstan, c.ph.-m.sc., docent, venera_bagg@mail.ru, ORCID: <https://orcid.org/0000-0001-8046-8508>;

Akzhigitova E.M. - al-Farabi Kazakh National University, Almaty, Kazakhstan, 1st course PhD student, kzo1994@mail.ru, ORCID: <https://orcid.org/0000-0002-8544-1248>

REFERENCES

- [1] V.S. Vasilevsky, K. Kato, V. Kumangaliyeva, A.D. Duisenbay, N. Kalzhigitov, N. Takibayev, Investigation of discrete and continuous spectrum states in two-cluster system. Sapporo, Japan: Hokkaido University, 2017
- [2] D.R. Thompson, M. LeMere, and Y.C. Tang, Systematic investigation of scattering problems with the resonating-group method, Nucl. Phys., vol. A286, no. 1, pp. 53-66, 1977.
- [3] I. Reichstein and Y.C. Tang, Study of N+ α system with the resonating group method Nucl. Phys. A, vol. 158, pp. 529-545, Dec. 1970.
- [4] A. Hasegawa and S. Nagata, Ground state of ⁶Li, Prog. Theor. Phys., vol. 45, pp. 1786-1807, 1971.
- [5] F. Tanabe, A. Tohsaki and R. Tamagaki, $\alpha\alpha$ scattering at intermediate energies, Prog. Theor. Phys., vol. 53, pp. 677-691, 1975.
- [6] M. Odsuren, A.T. Sarsembayeva, G. Khuukhenkhuu, S. Davaa, K. Kato, B. Usukhbayar, Higher excited states of $\alpha+\alpha$ system // News of the National Academy of Sciences of the Republic of Kazakhstan, series of physical-mathematical series. Vol. 2, №324. – 2019. – P.5-8, ISSN 2518-1726, <https://doi.org/10.32014/2019.2518-1726.6>
- [7] T. Kaneko, M. Shirata, H. Kanada, and Y.C. Tang, Neutron and proton form factors of ⁷Li and ⁷Be, Phys. Lett. B, vol. 192, pp. 259-262, July 1987.
- [8] H. Kanada, Q.K.K. Liu and Y.C. Tang, Charge form factor of ⁷Li with resonating-group wave function, Phys. Rev. C, vol. 22, pp. 813-822, Aug. 1980.
- [9] T. Kajino, T. Matsuse and A. Arima, Electromagnetic properties of ⁷Li and ⁷Be in a cluster model, Nucl. Phys. A, vol. 413, p. 323, Jan. 1984.
- [10] L.R. Suelzle, M.R. Yearian, and H. Crannell, Elastic Electron Scattering from Li⁶ and Li⁷, Phys. Rev., vol. 162, pp. 992-1004, Oct. 1967.

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы *М. С. Ахметова, Т.А. Апендиев, Д.С. Аленов*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 10.06.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
8,3 п.л. Тираж 300. Заказ 3.

Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19