

ISSN 2518-1726 (Online),  
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ  
Әль-фараби атындағы Қазақ ұлттық университетінің

# Х А Б А Р Л А Р Ы

---

---

## ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН  
Қазақстан Республикасының  
Ғылым Академиясының  
Әль-Фараби атындағы  
Қазақ ұлттық университетінің

## NEWS

OF THE NATIONAL ACADEMY OF SCIENCES  
OF THE REPUBLIC OF KAZAKHSTAN  
Al-farabi kazakh  
national university

**SERIES**  
**PHYSICO-MATHEMATICAL**

**4 (326)**

**JULY-AUGUST 2019**

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р ы  
ф.-м.ғ.д., проф., ҚР ҰҒА академигі **Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

**Жұмаділдаев А.С.** проф., академик (Қазақстан)  
**Кальменов Т.Ш.** проф., академик (Қазақстан)  
**Жантаев Ж.Ш.** проф., корр.-мүшесі (Қазақстан)  
**Өмірбаев У.У.** проф. корр.-мүшесі (Қазақстан)  
**Жүсіпов М.А.** проф. (Қазақстан)  
**Жұмабаев Д.С.** проф. (Қазақстан)  
**Асанова А.Т.** проф. (Қазақстан)  
**Бошқаев К.А.** PhD докторы (Қазақстан)  
**Сұраған Д.** корр.-мүшесі (Қазақстан)  
**Quevedo Hernando** проф. (Мексика),  
**Джунушалиев В.Д.** проф. (Қырғыстан)  
**Вишневский И.Н.** проф., академик (Украина)  
**Ковалев А.М.** проф., академик (Украина)  
**Михалевич А.А.** проф., академик (Белорус)  
**Пашаев А.** проф., академик (Әзірбайжан)  
**Такибаев Н.Ж.** проф., академик (Қазақстан), бас ред. орынбасары  
**Тигиняну И.** проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)  
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде  
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.  
Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор  
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Редакционная коллегия:

**Джумадильдаев А.С.** проф., академик (Казахстан)  
**Кальменов Т.Ш.** проф., академик (Казахстан)  
**Жантаев Ж.Ш.** проф., чл.-корр. (Казахстан)  
**Умирбаев У.У.** проф. чл.-корр. (Казахстан)  
**Жусупов М.А.** проф. (Казахстан)  
**Джумабаев Д.С.** проф. (Казахстан)  
**Асанова А.Т.** проф. (Казахстан)  
**Бошкаев К.А.** доктор PhD (Казахстан)  
**Сураган Д.** чл.-корр. (Казахстан)  
**Quevedo Hernando** проф. (Мексика),  
**Джунушалиев В.Д.** проф. (Кыргызстан)  
**Вишневский И.Н.** проф., академик (Украина)  
**Ковалев А.М.** проф., академик (Украина)  
**Михалевич А.А.** проф., академик (Беларусь)  
**Пашаев А.** проф., академик (Азербайджан)  
**Такибаев Н.Ж.** проф., академик (Казахстан), зам. гл. ред.  
**Тигиняну И.** проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов  
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

E d i t o r i n c h i e f  
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d:

**Dzhumadildayev A.S.** prof., academician (Kazakhstan)  
**Kalmenov T.Sh.** prof., academician (Kazakhstan)  
**Zhantayev Zh.Sh.** prof., corr. member. (Kazakhstan)  
**Umirbayev U.U.** prof. corr. member. (Kazakhstan)  
**Zhusupov M.A.** prof. (Kazakhstan)  
**Dzhumabayev D.S.** prof. (Kazakhstan)  
**Asanova A.T.** prof. (Kazakhstan)  
**Boshkayev K.A.** PhD (Kazakhstan)  
**Suragan D.** corr. member. (Kazakhstan)  
**Quevedo Hernando** prof. (Mexico),  
**Dzhunushaliyev V.D.** prof. (Kyrgyzstan)  
**Vishnevskiy I.N.** prof., academician (Ukraine)  
**Kovalev A.M.** prof., academician (Ukraine)  
**Mikhalevich A.A.** prof., academician (Belarus)  
**Pashayev A.** prof., academician (Azerbaijan)  
**Takibayev N.Zh.** prof., academician (Kazakhstan), deputy editor in chief.  
**Tiginyanu I.** prof., academician (Moldova)

**News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.**

**ISSN 2518-1726 (Online), ISSN 1991-346X (Print)**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

## NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN  
 PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.47>

Volume 4, Number 326 (2019), 83 – 91

UDC 517.946:517.588

IRSTI 27.29.21; 27.23.25

Zh.N. Tasmambetov<sup>1</sup>, N. Rajabov<sup>2</sup>, Zh.K.Ubayeva<sup>3</sup><sup>1</sup>Aqtobe Regional Zhybanov State University;<sup>2</sup>Tajic National University;<sup>3</sup>Aqtobe Regional Zhybanov State University[tasmam@rambler.ru](mailto:tasmam@rambler.ru), [nusrat38@mail.ru](mailto:nusrat38@mail.ru), [zhanar\\_ubaeva@mail.ru](mailto:zhanar_ubaeva@mail.ru)

## THE COORDINATED SOLUTION OF TWO DIFFERENTIAL EQUATIONS IN PRIVATE DERIVATIVES OF THE THIRD ORDER

**Abstract.** In the study the possibility of constructing a solution near a variety of special features of the system consisting of two differential equations in partial derivatives of the third order was investigated. There definitive aspects of such systems in comparison with systems consisting of two differential equations in partial derivatives of second order were established. The classification of regular and irregular singularities was carried out, the relevant solutions are constructed. A specific example shows that the application of the Frobenius - Latysheva method to the construction of a solution of a degenerate hypergeometric system consisting of two third-order partial differential equations. A variety of properties of system solution were considered, which shows that these solutions have the properties of generalized hypergeometric functions.

**Key words:** solution, constructed, system, features, regular, irregular, generalized, third order.

**Introduction.** The solutions of particular cases of systems of differential equations of second order type

$$\left. \begin{aligned} G^{(0)} \cdot Z_{xx} + G^{(1)} \cdot Z_{xy} + G^{(2)} \cdot Z_x + G^{(3)} \cdot Z_y + G^{(4)} \cdot Z &= 0 \\ Q^{(0)} \cdot Z_{yy} + Q^{(1)} \cdot Z_{xy} + Q^{(2)} \cdot Z_x + Q^{(3)} \cdot Z_y + Q^{(4)} \cdot Z &= 0 \end{aligned} \right\} \quad (1)$$

where  $G^{(i)} = G^{(i)}(x, y)$ ,  $Q^{(i)} = Q^{(i)}(x, y)$ ,  $(i = \overline{0,4})$  analytic functions or polynomials of two variables,  $Z = Z(x, y)$  – total unknown, are the generalized hypergeometric functions of two variables. The general theory of systems type (1) was established by American mathematician E. Wilczynski [1] - [2] using it to support special cases of projective differential geometry. He proved compatibility and integrability conditions

$$1 - \frac{G^{(1)}}{G^{(0)}} \cdot \frac{Q^{(1)}}{Q^{(0)}} \neq 0 \quad (2)$$

of system type (1) [1].

In carrying out these important conditions the following statement is fair.

**Theorem 1.** Suppose that the conditions of compatibility and integrability are fulfilled (2). Then, system (1) has four linearly independent particular solutions of the form

$$Z_j(x, y) = \sum_{m,n=0}^{\infty} A_{m,n} \cdot x^m \cdot y^n, \quad (A_{0,0} \neq 0), \quad (j = 1,2,3,4) \quad (3)$$

The overall solution of system (1) is represented as the sum

$$Z_k(x, y) = \sum_{k=1}^4 G_j \cdot Z_k(x, y), \quad k = 1, 2, 3, 4 \quad (4)$$

where  $G_j$ - is arbitrary constant, that is, the general solution of a system of partial differential equations depends on arbitrary constants, and not on an arbitrary constant.

**Theorem 2.** If the compatibility conditions are satisfied, and the integrability condition is not satisfied, that is,

$$1 - \frac{G^{(1)}}{G^{(0)}} \cdot \frac{Q^{(1)}}{Q^{(0)}} = 0 \quad (5)$$

then system (1) has no more than three linearly independent solutions, or the number of them will be countless.

Further development of the study was obtained in the works of J. Horn [3] - [4], P. Appell, J. Kampe de Feriet [5], W. Sternberg [6], A. Erdelyi [7] - [8], P. Humbert [9], E. Ince [10], L. Borgnesser [11] and others, in connection with the study of the theory of generalized hypergeometric series of two variables [12] - [13] - [14]. Using the concepts of rank  $p = 1 + k$  ( $k$  – subrank) and antiranga  $m = -1 - \lambda$  ( $\lambda$  – antipodrang). Zh.N. Tasmambetov developed [15] efficient algorithms for constructing normal, subnormal series, and normal-regular and final solutions of systems of the form (1) consisting of two second-order partial differential equations.

The aim of this work is to study a special system consisting of two differential equations in partial derivatives of the third order; to establish the distinctive features of such systems; to classify regular and irregular features and construct the corresponding solutions; to show the application of the Frobenius-Latysheva method to constructing a solution of a degenerate hypergeometric system consisting of two third-order partial differential equations by using a specific example.

### THE COORDINATED SOLUTION OF TWO DIFFERENTIAL EQUATIONS IN PRIVATE DERIVATIVES OF THE THIRD ORDER

**Statement.** Studying the possibility of coordinated solving a homogeneous system of third-order partial differential equations consisting of two equations of the form

$$\begin{aligned} x^3 g^{(0)} p_{30} + x^2 y g^{(1)} p_{21} + x^2 g^{(2)} p_{20} + x y g^{(3)} p_{11} + x g^{(4)} p_{10} + y g^{(5)} p_{01} + g^{(6)} p_{0,0} &= 0, \\ y^3 g^{(0)} p_{03} + x y^2 g^{(1)} p_{12} + y^2 g^{(2)} p_{02} + x y g^{(3)} p_{11} + x g^{(4)} p_{10} + y g^{(5)} p_{01} + g^{(6)} p_{0,0} &= 0, \end{aligned} \quad (6)$$

where

$$\begin{aligned} p_{30} = Z_{xxx}, p_{03} = Z_{yyy}, p_{21} = Z_{xxy}, p_{12} = Z_{xyy}, p_{20} = Z_{xx}, p_{02} = Z_{yy}, \\ p_{11} = Z_{xy}, p_{10} = Z_x, p_{01} = Z_y, p_{0,0} = Z(x, y) \end{aligned} \quad (7)$$

– total unknown, and the coefficients

$$\begin{aligned} g^{(i)} = g^{(i)}(x, y) = a_{00}^{(i)} + a_{10}^{(i)} \cdot x, \\ q^{(i)} = q^{(i)}(x, y) = b_{00}^{(i)} + b_{01}^{(i)} \cdot y, (i = \overline{0,6}). \end{aligned} \quad (8)$$

It is required to classify regular and irregular singularities and establish the type of solution near each singularity; to determine the number of linearly independent solutions near these features; to examine the relationship of the original system with systems solutions that are generalized hypergeometric functions of two variables and properties of such functions.

#### Solution of a system of differential equations.

To construct solutions of the system of equations (6), it is required to carry out a number of conditions, let us discuss the major of them

1. It must be carried out the compatibility conditions. However, the general compatibility condition, as for system (1), is difficult to control. Usually, such checking is carried out for specific systems solutions which are generalized hypergeometric functions of two variables. The Kampe de Feriet method can be applied [5, p.155], [16, p.21] - [17].

2. Integrability condition

$$1 - \frac{g^{(1)}}{g^{(0)}} \cdot \frac{q^{(1)}}{q^{(0)}} \neq 0. \quad (9)$$

3. Unknown function  $p_{0,0} = Z(x, y)$  depends on two variables, therefore, as before, depending on the regularity and irregularity of the singular curves, solutions should be sought in the form of generalized, normal and normal-regular series of two variables [15, p.159].

4. If in (6) the coefficients at higher derivatives  $p_{30}$  and  $p_{03} : x^3 \cdot g^{(0)}(x, y) \equiv 1$  and  $y^3 \cdot g^{(0)}(x, y) \equiv 1$  then solution can be sought as a simple series of two variables (3) since there are no singularities. Then, in order to formulate Theorems 1 and 2 for this case, it is necessary to establish the number of linearly independent particular solutions of system (6), which we will do next.

5. The special curves of the system (6) are established by equating to zero the coefficients of the highest derivatives.  $p_{30} = Z_{xxx}$  и  $p_{03} = Z_{yyy} :$

$$x^3 \cdot (a_{00}^{(0)} + a_{10}^{(0)} \cdot x) = 0, \quad x_1 = 0, \quad x_2 = -\frac{a_{00}^{(0)}}{a_{10}^{(0)}},$$

$$y^3 \cdot (b_{00}^{(0)} + b_{01}^{(0)} \cdot y) = 0, \quad y_1 = 0, \quad y_2 = -\frac{b_{00}^{(0)}}{b_{01}^{(0)}}.$$

Composed from them pairs  $(0,0), (0, -\frac{b_{00}^{(0)}}{b_{01}^{(0)}}), (-\frac{a_{00}^{(0)}}{a_{01}^{(0)}}, 0), (-\frac{a_{00}^{(0)}}{a_{01}^{(0)}}, -\frac{b_{00}^{(0)}}{b_{01}^{(0)}})$  - represent the final features,  $(0, \infty), (\infty, 0), (\infty, -\frac{b_{00}^{(0)}}{b_{01}^{(0)}}), (-\frac{a_{00}^{(0)}}{a_{01}^{(0)}}, \infty)$  and  $(\infty, \infty)$  - features at infinity. Normally, when building solutions two pairs of features identified  $(0,0)$  and  $(\infty, \infty)$ . Classification for regularity and irregularity will be carried out using simple rules [15].

**Rule I.** If in (6) coefficient  $a_{00}^{(0)} \neq 0, b_{00}^{(0)} \neq 0$ , then the feature  $(0,0)$  is special regular. When  $a_{00}^{(0)} = 0, b_{00}^{(0)} = 0$ , then  $(0,0)$  is a special irregular.

**Rule II.** If in (6) coefficient  $a_{10}^{(0)} \neq 0$  and  $b_{01}^{(0)} \neq 0$ , then the feature  $(\infty, \infty)$  - special regular, and when  $a_{10}^{(0)} = 0, b_{01}^{(0)} = 0$  - special irregular.

#### Features of constructing a solution by the Frobenius-Latyshev method.

The Frobenius-Latyshev method allows determining directly the structure and construct solutions of the system near particular curves. So, when the feature  $(0,0)$  is regular, the desired solution is represented as a generalized power series of two variables

$$Z(x, y) = x^\rho \cdot y^\sigma \cdot \sum_{m,n=0}^{\infty} A_{m,n} \cdot x^m \cdot y^n, \quad (10)$$

where  $\rho, \sigma$  and  $A_{m,n}$  ( $m, n = 0, 1, 2, 3, \dots$ ) - unknown constants to be determined. If the feature  $(0,0)$  is irregular, then the desired solution is represented as

$$Z(x, y) = \exp Q(x, y) \cdot x^\rho \cdot y^\sigma \cdot \sum_{m,n=0}^{\infty} A_{m,n} \cdot x^m \cdot y^n, \quad A_{0,0} \neq 0 \quad (11)$$

$Q(x, y)$  - polynomial of two variables

$$Q(x, y) = \frac{\alpha_{p1} \cdot x^p}{p} + \frac{y^p \cdot \alpha_{0p}}{p} + \dots + \alpha_{11} \cdot x \cdot y + \alpha_{10} \cdot x + \alpha_{01} \cdot y, \quad (12)$$

with uncertain parameters  $\alpha_{p,0}, \alpha_{0,p}, \dots, \alpha_{11}, \alpha_{10}, \alpha_{01}$ .

Similarly, near the regular singularity  $(\infty, \infty)$  the solution is constructed in the form

$$Z(x, y) = x^\rho \cdot y^\sigma \cdot \sum_{m,n=0}^{\infty} B_{m,n} \cdot x^{-m} \cdot y^{-n}, \quad B_{0,0} \neq 0 \quad (13)$$

and near irregular features in the form

$$Z(x, y) = \exp(x, y) \cdot x^\rho \cdot y^\sigma \cdot \sum_{m,n=0}^{\infty} B_{m,n} \cdot x^{-m} \cdot y^{-n}, \quad B_{0,0} \neq 0 \quad (14)$$

where  $\rho, \sigma$  and  $B_{m,n}$  ( $m, n = 0, 1, 2, 3, \dots$ ) – unknown constant.

Polynomial  $Q(x, y)$  is common to the type (11) and (14). Its degree is determined by the rank value

$$p = 1 + k, \quad k = \max_{(1 \leq s \leq n)} \frac{\beta_s - \beta_0}{s} \quad (15)$$

introduced by A. Poincare to study ordinary differential equations, and  $\beta_0, \beta_s$  - the greatest degrees of the coefficients of the equation [18].

### MAIN RESULTS

We will start the application of the Frobenius-Latysheva method with drawing up a system of Frobenius characteristic functions [15].

**Definition 1.** The system of characteristic functions of Frobenius is called the system

$$\begin{aligned} L_1[x^\rho \cdot y^\sigma] &\equiv x^\rho \cdot y^\sigma \cdot [f_{00}^{(1)}(\rho, \sigma) + f_{11}^{(1)}(\rho, \sigma) \cdot x] \\ L_2[x^\rho \cdot y^\sigma] &\equiv x^\rho \cdot y^\sigma \cdot [f_{00}^{(2)}(\rho, \sigma) + f_{11}^{(1)}(\rho, \sigma) \cdot y] \end{aligned} \quad (16)$$

where

$$f_{00}^{(1)}(\rho, \sigma) = a_{00}^{(0)} \rho(\rho-1)(\rho-2) + a_{00}^{(1)} \rho(\rho-1)\sigma + a_{00}^{(2)} \rho(\rho-1) + a_{00}^{(3)} \rho\sigma + a_{00}^{(4)} \rho + a_{00}^{(5)} \sigma + a_{00}^{(6)} \quad (16.1)$$

$$f_{00}^{(2)}(\rho, \sigma) = b_{00}^{(0)} \sigma(\sigma-1)(\sigma-2) + b_{00}^{(1)} \rho\sigma(\sigma-1) + b_{00}^{(2)} \sigma(\sigma-1) + b_{00}^{(3)} \rho\sigma + b_{00}^{(4)} \rho + b_{00}^{(5)} \sigma + b_{00}^{(6)} \quad (16.2)$$

$$f_{10}^{(1)}(\rho, \sigma) = a_{10}^{(0)} \rho(\rho-1)(\rho-2) + a_{10}^{(1)} \rho(\rho-1)\sigma + a_{10}^{(2)} \rho(\rho-1) + a_{10}^{(3)} \rho\sigma + a_{10}^{(4)} \rho + a_{10}^{(5)} \sigma + a_{10}^{(6)} \quad (16.3)$$

$$f_{01}^{(2)}(\rho, \sigma) = b_{01}^{(0)} \sigma(\sigma-1)(\sigma-2) + b_{01}^{(1)} \rho\sigma(\sigma-1) + b_{01}^{(2)} \sigma(\sigma-1) + b_{01}^{(3)} \rho\sigma + b_{01}^{(4)} \rho + b_{01}^{(5)} \sigma + b_{01}^{(6)} \quad (16.4)$$

obtained by substituting into system (6) with coefficients of the form (8) instead of  $Z = x^\rho \cdot y^\sigma$ . Define the system of determining equations for features  $(0,0)$  and  $(\infty, \infty)$  from (16).

**Definition 2.** The system of defining equations for the feature  $(0,0)$  is called the system

$$f_{00}^j(\rho, \sigma) = 0 \quad (j = 1, 2) \quad (17)$$

from which the indicators of series (10) and (11) are determined in the form of pairs  $(\rho_t, \sigma_t)$ .

**Definition 3.** A system of defining equations for a feature  $(\infty, \infty)$  is called a system

$$\begin{aligned} f_{10}(\rho, \sigma) &= 0, \\ f_{01}(\rho, \sigma) &= 0 \end{aligned} \quad (18)$$

from which the indicators of series (13) and (14) are determined in the form of pairs  $(\rho_t, \sigma_t)$ .

In this pairs  $(\rho_t, \sigma_t)$  it is important to determine the index  $t$ , since the number of such pairs allows determining the number of linearly independent solutions of the original system (6) near the specified features.



**Theorem 3.** For the existence the solutions of the form (10) in the system (6) with coefficients (8) near the singularity (0,0) requires the equality (17). **Theorem 4.** For the existence of solutions of the form (13) in the system (6) with coefficients (8) near the singularity  $(\infty, \infty)$  equality is necessary (18).

Let's determine how many roots have the systems of constitutive equations (17) and (18). In view of this, the system (17) can be written in expanded form, using (16.1) and (16.2). From  $f_{0,0}^{(1)}(\rho, \sigma) = 0$  we will find out

$$\sigma = \frac{a_{00}^{(0)} \rho(\rho - 1)(\rho - 2) + a_{00}^{(2)} \rho(\rho - 1) + a_{00}^4 \rho + a_{10}^{(6)}}{a_{00}^{(1)} \rho(\rho - 1) + a_{00}^{(3)} \rho + a_{00}^{(5)}}$$

and substituting in the second equation  $f_{00}^{(2)}(\rho, \sigma) = 0$  of the system (17), after exclusion  $\rho$ , we obtain the ninth degree equation for  $\sigma$ . If there are only simple roots, then from here we define nine roots  $\sigma_t$  ( $t = \overline{1,9}$ ). Similarly, you can define nine simple roots  $\rho_t$  ( $t = \overline{1,9}$ ). Of these, make nine pairs of roots  $(\rho_t, \sigma_t)$  ( $t = \overline{1,9}$ ) of system (17), which allows us to construct nine linearly independent particular solutions of system (17) near the singularity (0,0).

**Theorem 5.** Suppose the system (6) with coefficients of the form (8), where  $a_{00}^{(0)} \neq 0, b_{00}^{(0)} \neq 0$  conditions of compatibility and integrability are satisfied (9). Then system (6) has nine linearly independent regular particular solutions of the form

$$Z_t(x, y) = x^{\rho_t} \cdot y^{\sigma_t} \cdot \sum_{m,n=0}^{\infty} A_{m,n}^{(t)} \cdot x^m \cdot y^n, \quad A_{0,0}^{(t)} \neq 0, \quad (t = \overline{1,9}), \quad (19)$$

near feature (0,0), where  $\rho_t, \sigma_t$  ( $t = \overline{1,9}$ ),  $A_{m,n}^{(t)}$  ( $m, n = 0, 1, 2, \dots$ ).

A similar argument can be sure that the system (6) also has nine linear - independent particular solutions near the singularity  $(\infty, \infty)$ .

**Theorem 6.** Suppose the system (6) with coefficients of the form (8), where  $a_{10}^{(0)} \neq 0, b_{01}^{(0)} \neq 0$  conditions of compatibility and integrability are satisfied (9). Then the system (6) has nine linearly independent regular particular solutions of the form

$$Z_t(\rho, \sigma) = x^{\rho_t} \cdot y^{\sigma_t} \cdot \sum_{m,n=0}^{\infty} B_{m,n}^{(t)} \cdot x^m \cdot y^n, \quad B_{0,0}^{(t)} \neq 0, \quad (t = \overline{1,9}), \quad (20)$$

near feature  $(\infty, \infty)$ , where  $\rho_t, \sigma_t$ , ( $t = \overline{1,9}$ ),  $B_{m,n}^{(t)}$  ( $m, n = 0, 1, 2, \dots$ ) - unknown constant.

In the theorem 5 and 6, conditions  $a_{00}^{(0)} \neq 0, b_{00}^{(0)} \neq 0$  and  $a_{10}^{(0)} \neq 0, b_{01}^{(0)} \neq 0$  essential since, ninth degree equations are relatively  $\rho$  and  $\sigma$  it turns out only when they are non-zero. Therefore, near regular singular curves (0,0) and  $(\infty, \infty)$  there are nine regular linearly independent particular solutions. Then, the overall solution is represented as the following sum

$$Z(x, y) = \sum_{t=1}^9 C_t \cdot Z_t(x, y), \quad (t = \overline{1,9}). \quad (21)$$

Unknown row coefficients (19)  $A_{m,n}^{(t)}$  ( $m, n = 0, 1, 2, \dots$ ) are determined from systems of recurrent sequences

$$\sum_{m,n=0}^{\infty} A_{\mu-m, \nu-n}^{(t)} \cdot f_{m,n}^{(j)}(\rho + \mu - m, \sigma + \nu - n) = 0 \quad (22)$$

$(\mu, \nu = 0, 1, 2, \dots, ; j = 1, 2; t = \overline{1,9})$  (obtained by substituting series (19) into system (6) with coefficients (8).

**Application to the construction of a specific system solutions.**

All hypergeometric functions of two variables satisfy a system consisting of two partial differential equations of second order. The coefficient of such systems are polynomials  $x$  and  $y$ . These coefficients can be calculated if in the hypergeometric functions defined by double series

$$F(x, y) = \sum_{m,n} a_{m,n} \cdot x^m \cdot y^n, \tag{23}$$

the coefficients satisfy the following relations

$$\frac{a_{m+1,n}}{a_{m,n}} = \frac{P(m,n)}{R(m,n)}, \quad \frac{a_{m,n+1}}{a_{m,n}} = \frac{Q(m,n)}{S(m,n)}, \tag{24}$$

where P, R, Q, S are known polynomials.

The use of such an approach enabled the intensification of the study of relations between systems consisting of two second-order equations and generalized hypergeometric series of two variables.

However, the relationship between systems consisting of two equations of the third and fourth orders and generalized hypergeometric series has not been studied at the proper level. Here, it should be noted the work of Kampe de Feriet [5], [16] - [17], which provides a method for constructing a series of third and fourth order systems, while ensuring the compatibility of such systems. In this paper we will also proceed from the studies of J. Kampe de Feriet [5, p.155-169].

**Theorem 7.** The system of partial differential equations consisting of two equations of third order

$$\begin{aligned} x^2 \cdot p_{30} + (\delta_1 + \delta_2 + 1) \cdot x \cdot p_{20} + (\delta_1 \cdot \delta_2 - x) \cdot p_{10} - y \cdot p_{01} - \alpha \cdot p_{0,0} &= 0, \\ y^2 \cdot p_{30} + (\delta_1' + \delta_2' + 1) \cdot y \cdot p_{02} + (\delta_1 \cdot \delta_2 - y) \cdot p_{01} - x \cdot p_{10} - \alpha \cdot p_{0,0} &= 0, \end{aligned} \tag{25}$$

where  $p_{30} = Z_{xxx}, p_{03} = Z_{yyy}, p_{20} = Z_{xx}, p_{02} = Z_{yy}, p_{10} = Z_x, p_{01} = Z_y, p_{0,0} = Z(x, y)$  – common unknown, has nine linearly independent regular particular solutions, and the general solution is represented as the sum (21).

**Argument.** We will carry out the proof by the Frobenius-Latyshev method, while revealing the additional properties of system (25) can be obtained from the original system (6) with coefficients of the form (8) using the Kampe de Feriet method [5, p.155-164]. As we noted above, this ensures the compatibility of the two equations of the system (25). The integrability condition (9) is also satisfied, since  $g^{(1)} = q^{(1)} \equiv 0$ .

Coefficients  $a_{00}^{(0)} \neq 0$  и  $b_{00}^{(0)} \neq 0$ , so the feature (0,0) is regular based on the Rule I, a  $a_{10}^{(0)} = 0, b_{01}^{(0)} = 0$ , therefore, on the basis of the Rule II feature  $(\infty, \infty)$  – irregular. Let us build regular solutions in the form (10) near feature (0,0).

To this end, we compose a system of Frobenius characteristic functions (15) - (16) and define systems of defining equations (17) with respect to the singularity (0,0), in a transformed expanded form

$$\left. \begin{aligned} f_{00}^{(1)}(\rho, \sigma) &= \rho \cdot (\rho - 1 + \delta_1) \cdot (\rho - 1 + \delta_2) = 0, \\ f_{00}^{(2)}(\rho, \sigma) &= \sigma \cdot (\sigma - 1 + \delta_1) \cdot (\sigma - 1 + \delta_2) = 0. \end{aligned} \right\} \tag{26}$$

It has nine pairs of roots  $(\rho_t, \sigma_t, t = \overline{1,9})$ :

1.  $(\rho_1 = 0, \sigma_1 = 0)$ , 2.  $(\rho_2 = 1 - \delta_1, \sigma_1 = 0)$ , 3.  $(\rho_1 = 0, \sigma_2 = 1 - \delta_1')$ , 4.  $(\rho_3 = 1 - \delta_2, \sigma_1 = 0)$ ,
5.  $(\rho_1 = 0, \sigma_3 = 1 - \delta_2')$ , 6.  $(\rho_2 = 1 - \delta_1, \sigma_2 = 1 - \delta_1')$ , 7.  $(\rho_2 = 1 - \delta_1, \sigma_3 = 1 - \delta_2')$ ,
8.  $(\rho_3 = 1 - \delta_2, \sigma_2 = 1 - \delta_1')$ , 9.  $(\rho_3 = 1 - \delta_2, \sigma_3 = 1 - \delta_2')$ .

Then, on the basis of Theorem 5, the system (25) near the singularity (0,0) has nine linearly - independent regular partial solutions, corresponding to these indicators.:

$$\begin{aligned}
 Z_1 &= \Phi(\alpha, \delta_1, \delta_1', \delta_2, \delta_2'; x, y), \\
 Z_2 &= x^{1-\delta_1} \cdot \Phi(\alpha + 1 - \delta_1; 2 - \delta_1', \delta_1', \delta_2 + 1 - \delta_1, \delta_2'; x, y), \\
 Z_3 &= y^{1-\delta_1'} \cdot \Phi(\alpha + 1 - \delta_1', \delta_1; 2 - \delta_1', \delta_2, \delta_2' + 1 - \delta_1'; x, y), \\
 Z_4 &= x^{1-\delta_2} \cdot \Phi(\alpha + 1 - \delta_2; \delta_1 + 1 - \delta_2, \delta_1', 2 - \delta_2'; x, y), \\
 Z_5 &= y^{1-\delta_2'} \cdot \Phi(\alpha + 1 - \delta_2', \delta_1; \delta_1' + 1 - \delta_2', 2 - \delta_2'; x, y), \\
 Z_6 &= x^{1-\delta_1} \cdot y^{1-\delta_1'} \cdot \Phi(\alpha + 2 - \delta_1, \delta_1'; 2 - \delta_1, 2 - \delta_1', \delta_2 + 1 - \delta_1, \delta_2' + 1 - \delta_1'; x, y), \\
 Z_7 &= x^{1-\delta_1} \cdot y^{1-\delta_2'} \cdot \Phi(\alpha + 2 - \delta_1 - \delta_2'; 2 - \delta_1, 2 - \delta_1 \cdot \delta_1', \delta_1' + 1 - \delta_2', \delta_2 + 1 - \delta_1, 2 - \delta_1'; x, y), \\
 Z_8 &= x^{1-\delta_2} \cdot y^{1-\delta_2'} \cdot \Phi(\alpha + 2 - \delta_2 - \delta_1, \delta_1 + 1 - \delta_2, 2 - \delta_1' 2 - \delta_2, \delta_2' + 1 - \delta_1'; x, y), \\
 Z_9 &= x^{1-\delta_2} \cdot y^{1-\delta_2'} \cdot \Phi(\alpha + 2 - \delta_2 - \delta_2', \delta_1 + 1 - \delta_2, \delta_1 + 1 - \delta_2 \cdot \delta_1' + 1 - \delta_2', 2 - \delta_2, 2 - \delta_2'; x, y).
 \end{aligned}
 \tag{27}$$

All the solutions obtained coincide exactly with the solutions in the monograph. [5, p.166].

**Conclusions:** We note some properties of the studied system and its solution.

**Properties 1.** The monograph [5] does not classify the regular and irregular features of the system (25). In system (25)  $a_{10}^{(0)} = 0$  and  $b_{01}^{(0)} = 0$ . Therefore, on the basis of Rule II, we conclude that the peculiarity  $(\infty, \infty)$  irregular and system (25) must have a solution in the form of a normal Tome series (14). However, in this case, the system of defining equations will be written as

$$\begin{aligned}
 f_{10}(\rho, \sigma) &= -(\rho + \sigma + 2) = 0, \\
 f_{01}(\rho, \sigma) &= -(\rho + \sigma + 2) = 0
 \end{aligned}$$

that is, it boils down to one equation and defines an infinite number of solutions. Then, due to the uncertainty of the indicators  $(\rho_i, \sigma_i)$  there is no solution in the form of a normal Tome series (14)

**Properties 2.** It is also not difficult to make sure that the solutions of system (25) presented in the form of (27) possess many properties of generalized hypergeometric series [5],[14]. Thus, the first derivatives (27) are represented as

$$\begin{aligned}
 \frac{\partial Z_1}{\partial x} &= \frac{\partial \Phi[\alpha, \delta_1, \delta_1', \delta_2, \delta_2'; x, y]}{\partial x} = \frac{\alpha}{\delta_1 \cdot \delta_2} \cdot \Phi[\alpha + 1, \delta_1 + 1, \delta_1'; 1 + \delta_2, \delta_2'; x, y] \\
 \frac{\partial Z_1}{\partial y} &= \frac{\partial \Phi[\alpha, \delta_1, \delta_1', \delta_2, \delta_2'; x, y]}{\partial y} = \frac{\alpha}{\delta_1' \cdot \delta_2'} \cdot \Phi[\alpha + 1, \delta_1, \delta_1' + 1, \delta_2, \delta_2' + 1; x, y] \\
 \frac{\partial Z_1}{\partial x \partial y} &= \frac{\partial \Phi[\alpha, \delta_1, \delta_1', \delta_2, \delta_2'; x, y]}{\partial x \partial y} = \frac{\alpha(\alpha + 1)}{\delta_1 \cdot \delta_2 \cdot \delta_1' \cdot \delta_2'} \cdot \Phi[\alpha + 2, \delta_1 + 1, \delta_1' + 1, \delta_2 + 1, \delta_2' + 1; x, y]
 \end{aligned}
 \tag{28}$$

**Properties 3.** The first solution  $Z_1(x, y)$  represents a double row depending on five parameters  $\alpha, \delta_1, \delta_2, \delta_1', \delta_2'$  [14]:

$$Z_1(x, y) = {}_1\Phi_4 \left[ \begin{matrix} \alpha \\ \delta_1, \delta_1', \delta_2, \delta_2' \end{matrix}; x, y \right] = \sum_{m,n=0}^{\infty} \frac{(\alpha)_{m+n}}{(\delta_1)_m (\delta_1')_n (\delta_2)_m (\delta_2')_n} \cdot \frac{x^m}{m!} \cdot \frac{y^n}{n!}
 \tag{29}$$

In the same way, it can be verified that the remaining solutions (27) have similar properties, that is, all the functions listed in (27) are generalized hypergeometric functions [14], [19]. The question of the computational application of special functions of several variables, as in monographs, remains topical [20]. The development [21] of a numerical method for calculating the values of the degenerate

hypergeometric Humbert functions is also required  $\Psi_2(\alpha, \gamma, \gamma'; x, y)$  through the products of Laguerre polynomials [22].

**Ж.Н. Тасмамбетов<sup>1</sup>, Н. Раджабов<sup>2</sup>, Ж.К. Убаева<sup>3</sup>**

<sup>1</sup>Қ.Жұбанов атындағы АӨМУ, Ақтөбе, Қазақстан;

<sup>2</sup>Тәжік ұлттық университеті, Душанбе, Тәжікстан;

<sup>3</sup>Қ.Жұбанов атындағы АӨМУ, Ақтөбе, Қазақстан

### **ҮШІНШІ РЕТТІ ДЕРБЕС ТУЫНДЫЛЫ ЕКІ ДИФФЕРЕНЦИАЛДЫҚ ТЕНДЕУЛЕРДІҢ БІРЛЕСКЕН ШЕШІМІ**

**Аннотация.** Мақалада үшінші ретті дербес туындылы екі дифференциалдық тендеулерден тұратын арнайы жүйенің ерекше нүктелерінің маңайындағы шешімдерін тұрғызу мүмкіндіктері зерттелді. Мұндай жүйелердің айрықша ерекше нүктелері екінші ретті дербес туындылы екі дифференциалдық тендеулерден тұратын жүйелермен салыстыру арқылы алынды. Регулярлы және иррегулярлы ерекшеліктеріне жіктеу жүргізіліп, олардың сәйкес шешімдері тұрғызылды. Нақты мысал арқылы үшінші ретті дербес туындылы екі дифференциалдық тендеулерден тұратын туындалған гипергеометриялық жүйенің шешімін табуда Фробениус-Латышева әдісінің қолдануы келтірілген. Жүйе шешімдерінің бірқатар қасиеттері келтіріліп, бұл шешімдердің өзі жалпыланған гипергеометриялық функциялардың да қасиеттеріне ие екені көрсетілген.

**Түйін сөздер:** шешім, тұрғызылған, жүйе, ерекше нүктелер, регулярлы, иррегулярлы, жалпыланған, үшінші ретті.

УДК 517.946:517.588

**Ж.Н. Тасмамбетов<sup>1</sup>, Н. Раджабов<sup>2</sup>, Ж.К. Убаева<sup>3</sup>**

<sup>1</sup>АРГУ им. К.Жубанова, Ақтөбе, Қазақстан;

<sup>2</sup>Таджикский национальный университет, Таджикистан, Душанбе;

<sup>3</sup>АРГУ им. К.Жубанова, Ақтөбе, Қазақстан

### **СОВМЕСТНОЕ РЕШЕНИЕ ДВУХ ДИФФЕРЕНЦИАЛЬНЫХ УРАВНЕНИЙ В ЧАСТНЫХ ПРОИЗВОДНЫХ ТРЕТЬЕГО ПОРЯДКА**

**Аннотация.** В работе изучены возможности построения решения вблизи различных особенностей специальной системы состоящим из двух дифференциальных уравнений в частных производных третьего порядка. Установлены отличительные особенности таких систем по сравнению с системами состоящих из двух дифференциальных уравнений в частных производных второго порядка. Проведена классификация регулярных и иррегулярных особенностей, построены соответствующие им решения. На конкретном примере показано применение метода Фробениуса-Латышевой к построению решения вырожденной гипергеометрической системы состоящих из двух дифференциальных уравнений в частных производных третьего порядка. Рассмотрены ряд свойств решения системы, которое показывает, что эти решения обладают свойствами обобщенных гипергеометрических функций.

**Ключевые слова:** решение, построены, система, особенности, регулярные, иррегулярные, обобщенная, третьего порядка.

#### **Information about authors:**

**Tasmambetov Zhaksylyk Nuradinovich** – Aqtobe Regional Zhubanov State University, Doctor of physical and mathematical sciences, Professor Chair of Mathematics, [tasmam@rambler.ru](mailto:tasmam@rambler.ru);

**Rajabov Nusrat** – Tajik National University, Professor Chair Mathematical Analysis and Function Theory Tajik National University, Academician Academy of Science the Republic of Tajikistan, [nusrat38@mail.ru](mailto:nusrat38@mail.ru);

**Ubayeva Zhanar Kartbaevna** – Phd Student of Aqtobe Regional Zhubanov State University, [Zhanar\\_ubayeva@mail.ru](mailto:Zhanar_ubayeva@mail.ru).

#### **REFERENCES**

- [1] Wilczynski E.J. Projective differential geometry of Curves and Ruled surfaces. Leipzig: Leubner, 1966-120pp.
- [2] Wilczynski E.J. On a Certain Completerly Integrable system of Linear. Partial Differential Equations // Amer. Journal of Math., 36, №3, 1994, p.180-194.

- [3] Horn J. Uber die Convergenz der hypergeometrischen Reihen Zweierund dreier Veranderlichen //Math. Ann, 1889,v.34, p.544-600.
- [4] Horn J. Uber das verhalten der Integrale Lineere Differenzen und Differentialgleichungen fur gobe der veranderlichen // Journal fur Math, 1910, v.138, p. 159-191.
- [5] Appell P., Kampe de Feriet M.J. Fonctions hypergeometriques et hypersperiques. Paris: Gauthier Villars. 1926. -434 pp.
- [6] Sternberg W. Uber dis asymptotische Integration von Differential gleichungen. Marh. Ann., 1920, Bd.81, 119-186 pp.
- [7] Erdelyi A. Same confluent Hypergeometric Functions of two variables. Proc. Roy. Sos Edinburgh, 1940, v.60, p.344-361.
- [8] Erdelyi A. Same confluent Hypergeometric Functions of two variables. Acta. Math, 1950, v.83, p.130-163.
- [9] Humbert P. The confluent Hypergeometric Functions of two variables. Proc. Roy. Sos. Edin., 1920, v.10, p.73-96.
- [10] Ince E.L. Proc. Roy. Sos. Edinburgh, A61,1942, p.195-209.
- [11] Borngusser L. Uber hypergeometrische Functionen zweier Verander Lichen, Dissertation. Darmstsd , 1933, 236 p.
- [12] Bateman G., Erdelyi A. Higher Transcendental Functions. Part I. Hypergeometric functions. The functions of the Lezhanders. M.: Science, 1965, -294 pp. (In Russian).
- [13] Slater L.J., Lit D., PH. D. Confluent hypergeometric series. Cambridge At the university Press, 1960.-249 pp.
- [14] Slater L.J., Lit D., PH. D. Generalized Hypergeometric functions. Cambridge At the university Press, 1966.-292 pp.
- [15] Tasmambetov Zh.N. Construction of normal and normally-regular solutions of special systems of partial equations of second order. IP Zhanadilov S.T., Actobe, 2015, 463pp. (In Russian).
- [16] Kampe de Feriet J. Fonctions de la physique Mathematique. Fascicule IX, M.: 1963, 102 pp. (In Russian).
- [17] Kampe de Feriet J. Quelques proprietis des fonctions hypergeometriques dordre superieur a deux variables. Comptes rendus, t. CLXXIII, 1921, p.489
- [18] Latysheva K.Ya., Tereschenko N.I., Orel N.S. Normal-regular solutions and their applications. Kiev, v.sh., 1974, p.136. (In Russian).
- [19] Bailey WN. Generalized hypergeometric series. Cambridge Mathematical Tract. 1935, №32 (Second edition 1964). Cambridge University Press.
- [20] Luke Yedell L. Mathematical funktions and their approximations. Academic Rpress.Jue. New York- San Francisco - London. 1975.608 p.
- [21] Dzhumabaev D.S., Bakirova EA., Kadirbayeva ZhM. (2018) An algorithm for solving a control problem for a differential Equation with a parameter. News of the National Academy of Sciences of the Republic of Kazakhstan. Series of physic-mathematical sciences. Volume 5, Number 321(2018), pp. 25-32. <https://doi.org/10.32014/2018.2518-1726.4> ISSN 1991-346X. ISSN 2518-1726 (Online), ISSN 1991-346X (Print).
- [22] Tasmambetov Zh.N., Rajabov N., Issenova A.A. (2019) The construction of a solution of a related system of the laguerre type. New of the National Academy of Sciences of the Republic of Kazakhstan. Series of physic-mathematical sciences. Volume 1, Number 323(2019), pp. 38-45. <https://doi.org/10.32014/2019.2518-1726.5> ISSN 1991- 346X. ISSN 2518-1726 (Online), ISSN 1991-346X (Print).

**Publication Ethics and Publication Malpractice  
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

**ISSN 2518-1726 (Online), ISSN 1991-346X (Print)**

Редакторы *М. С. Ахметова, Т.А. Апендиев, Д.С. Аленов*  
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 10.08.2019.  
Формат 60x881/8. Бумага офсетная. Печать – ризограф.  
9,6 п.л. Тираж 300. Заказ 4.

---

*Национальная академия наук РК*  
*050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19*