

**ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)**

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Әль-фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Казахский национальный университет
имени Аль-фараби

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-farabi kazakh
national university

SERIES
PHYSICO-MATHEMATICAL

4 (326)

JULY-AUGUST 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Бас редакторы
ф.-м.ғ.д., проф., КР ҮФА академигі **F.M. Мұтанов**

Редакция алқасы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев Ү.Ү. проф. корр.-мүшесі (Қазақстан)
Жусіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Ә. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорус)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«КР ҮФА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Үлттық ғылым академиясы» РКБ (Алматы қ.)
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы қуәлік

Мерзімділігі: жылдана 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Үлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Р е д а к ц и о н на я кол л е г и я:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жантаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф. чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

E d i t o r i n c h i e f
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d:

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskyi I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.40>

Volume 4, Number 326 (2019), 22 – 29

UDK 517.929

A.Sh.Shaldanbayev¹, M.I.Akylbayev², A.A.Shaldanbayeva², A.Zh.Beisebayeva³

¹Silkway International University, Shymkent, Kazakhstan;

²Regional social-innovative University, Shymkent, Kazakhstan;

³South Kazakhstan State University M.Auezova, Shymkent, Kazakhstan

shaldanbaev51@mail.ru, musabek_kz@mail.ru, alitma_a@mail.ru, akbope_a@mail.ru

**ON THE SPECTRAL PROPERTIES OF A WAVE OPERATOR
PERTURBED BY A LOWER-ORDER TERM**

Abstract. The incorrectness of the minimal wave operator is well known, since zero is an infinite-to-one eigenvalue for it. As our study showed, the situation changes if the operator is perturbed by a low-order term containing the spectral parameter as a coefficient, and eventually the problem takes the form of a beam of operators. The resulting beam of operators is easily factorized by first-order functional-differential operators which spectral properties are easily studied by the classical method of separation of variables. Direct application of the method of separation of variables to the original beam of operators encounters the insurmountable difficulties.

Keywords: deviating argument, beam of operators, strong solvability, spectrum, functional - differential operator.

Introduction.

Investigations of the Dirichlet problem for the string oscillation equation in a bounded domain go back to J. Hadamard [1], who for the first time noted nonuniqueness of solution of the Dirichlet problem for a wave equation in a rectangle. Burgin and Duffin [2] considered the Dirichlet problem for the equation $U_{xx} = Utt$ in the rectangle $\{0 < x < X ; 0 < t < T\}$. It was described that the nonuniqueness of a solution in a specified space arises if and only if the ratio X/T is rational. By using Laplace transformation, they showed that if the number X/T is irrational, then there is the uniqueness of the solution of the problem in the class of continuously differentiable functions with the second derivatives integrable according to Lebesgue.

Later these results were refined and generalized by various authors (see, for example, [3], [4], [5], [6]). S.L. Sobolev [7] constructed an example of a well-posed boundary-value problem in a rectangle for a hyperbolic system of equations. Yu.M. Berezanskii [8] constructed a class of regions with angles, a change in the domain inside which leads to a continuous change in the solution of the Dirichlet problem.

For domains with a smooth boundary in smooth spaces, only the question of the uniqueness of the solution of the Dirichlet problem was studied (see, for example, the paper of Aleksandryan [9]). In paper [3] V.I. Arnol'd, applying his results on the mapping of the circle into itself, refines the results of [2], indicating that the proof of theorems on the existence of classical solutions of the Dirichlet problem can be carried over to the case of an ellipse. A number of studies by T.Sh. Kal'menov and M.A. Sadybekov are also devoted to boundary value problems of hyperbolic equations [10] - [12].

In [13], using the new general method, the properties of solutions of the Cauchy problem, as well as of the first, second and third boundary-value problems in the disk for a second-order hyperbolic equation with constant coefficients are investigated. The application of this method to higher-order equations can be found in [14]. A new and relatively simple method for constructing a system of polynomial solutions of the Dirichlet problem for second-order hyperbolic equations with constant coefficients in the disk is proposed in [15], and it is also proposed to construct a complete set of eigenfunctions for the Dirichlet

problem for the string oscillation equation. The eigenfunctions constructed in this paper coincide with the eigenfunctions constructed earlier in the work of R.A. Aleksandaryan [9]. In this paper, the spectral properties of a beam of operators with a wave operator in the principal part were studied by the methods of [16]–[21].




Figure 1

1. Setting of a Problem.

Let $\Omega \subset R^2$ be a quadrangle bounded by following segments:

$AB : 0 \leq y \leq 1, x = 0; BC : 0 \leq x \leq 1, y = 1; CD : 0 \leq y \leq 1, x = 1; DA : 0 \leq x \leq 1, y = 0.$ /See fig.1/

We denote by $C^{1,1}(\Omega)$ the set of functions $u(x, t)$ that are twice continuously differentiable with respect to the variables x and t . The boundary of the area Ω is a set of segments $\Gamma = AB \cup BC \cup CD \cup DA$.

We investigate the spectral properties of the operator beam

$$u_{xx} - u_{yy} = -2\lambda u_x + \lambda^2 u, \quad (1)$$

$$u|_{y=0} = 0, \quad (2)$$

$$u|_{x=0} = \alpha u|_{x=1}, \quad |\alpha| = 1. \quad (3)$$

2. Research Methods.

We shall need the following lemmas.

Lemma 2.1. [22]. Let A be a densely defined operator in a Hilbert space H . Then

- (a) A^* exists and is closed;
- (b) A admits a closure if and only if $D(A^*)$ is dense in H , and in this case $\overline{A} = A^{**}$.

Lemma 2.2. The set of functions that are finite in the domain Ω is dense in the space $L^2(\Omega)$ [23].

Lemma 2.3. If the symmetric operator A has a complete system of eigenvectors, then the closure of this operator \overline{A} is self-adjoint in H , in other words, the operator A is an essentially self-adjoint in H [22].

Lemma 2.4. Operator

$$Lu = iu_x(x, y) + u_y(x, 1-y) \quad (4)$$

$$D(L) = \{u \in C^{1,1}(\Omega) \cap C(\overline{\Omega}); u|_{y=0} = 0, u|_{x=0} = \alpha u|_{x=1}, |\alpha| = 1\}. \quad (5)$$

is a symmetric operator in space $L^2(\Omega)$.

{The proof is omitted, because it is done in the standard way}.

Proof. Let $u, v \in D(L)$, then

$$(Lu, v) = \int_0^1 \int_0^1 [iu_x(x, y) + u_y(x, 1-y)] \cdot \bar{v}(x, y) dx dy = \int_0^1 \int_0^1 iu_x(x, y) \bar{v}(x, y) dx dy + \\ \int_0^1 \int_0^1 u_y(x, 1-y) \bar{v}(x, y) dx dy = J_1 + J_2.$$

Using Fubini's theorem and integrating by parts, we transform the integrals J_1, J_2 .

$$J_1 = \int_0^1 \int_0^1 iu_x(x, y) \bar{v}(x, y) dx dy = \int_0^1 \left[\int_0^1 iu_x(x, y) \bar{v}(x, y) dx \right] dy = \int_0^1 \left[\int_0^1 i\bar{v}(x, y) d_x u \right] dy = \\ \int_0^1 \left[i\bar{v}(x, y) u(x, y) \Big|_0^1 - i \int_0^1 u(x, y) \bar{v}_x(x, y) dx \right] dy = \int_0^1 \left[\int_0^1 u(x, y) i\bar{v}_x(x, y) dx \right] dy = \\ \int_0^1 \int_0^1 u(x, y) i\bar{v}_x(x, y) dx dy; \\ J_2 = \int_0^1 \int_0^1 u_y(x, 1-y) \bar{v}(x, y) dx dy = \int_0^1 \left[\int_0^1 u_y(x, 1-y) \bar{v}(x, y) dy \right] dx = \int_0^1 \left[- \int_0^1 \bar{v}(x, y) d_y u(x, 1-y) \right] dx = \\ \int_0^1 \left[- \bar{v}(x, y) u(x, 1-y) \Big|_0^1 + \int_0^1 u(x, 1-y) \bar{v}_y(x, y) dy \right] dx = \int_0^1 \left[\int_0^1 u(x, 1-y) \bar{v}_y(x, y) dy \right] dx = \\ \int_0^1 \int_0^1 u(x, y) \bar{v}_y(x, 1-y) dx dy;$$

Consequently, $(Lu, v) = \int_0^1 \int_0^1 u(x, y) [iv_x(x, y) + v_y(x, 1-y)] dx dy = (u, Lv)$

Lemma 2.5. The spectral problem

$$Lw = -w''(y) = \nu^2 w(y), \\ w(0) = w'(1) = 0$$

has an infinite set of eigenvalues

$$\nu_n = n\pi - \frac{\pi}{2}, \quad n = 1, 2, \dots$$

and corresponding eigenfunctions

$$w_n(y) = \sqrt{2} \sin\left(n\pi - \frac{\pi}{2}\right)y, \quad n = 1, 2, \dots,$$

which form an orthonormal basis of the space $L^2(0,1)$ [24].

Lemma 2.6. The spectral problem

$$iv_x = \mu v(x), \quad v(0) = \alpha v(1), \quad |\alpha| = 1 \quad (6)$$

has an infinite set of real eigenvalues

$$\mu_m = arg\alpha + 2m\pi i, \quad m = 0, \pm 1, \pm 2, \dots \quad (7)$$

and their corresponding eigenfunctions

$$v_m(x) = \exp[-i(\arg \alpha + 2m\pi)x], m = 0, \pm 1, \pm 2, \dots \quad (8)$$

which form an orthonormal basis of space $L^2(0,1)$ [16].

Lemma 2.7. If the orthogonal systems $\{\Phi_n(x)\}$ and $\{\psi_n(x)\}$, $n = 1, 2, \dots$ are complete in space $L^2(0,1)$, then their product $\{\Phi_n(x) \cdot \psi_n(x)\}$, $m, n = 1, 2, \dots$ is complete in the space $L^2(0,1)$, where $\Omega = [0,1] \times [0,1]$ [25].

3. Results of the research.

Theorem 1. The boundary value problem

$$Lu = iu_x(x, y) + u_y(x, 1-y) = f(x, y), \quad (9)$$

$$u|_{y=0} = 0, \quad u|_{x=0} = \alpha u|_{x=1}, \quad |\alpha| = 1, \quad (10)$$

has an infinite set of real eigenvalues

$$\lambda_{mn} = \arg \alpha + 2m\pi + (-1)^{n+1} \left(n\pi - \frac{\pi}{2} \right), \quad n = 1, 2, \dots, m = 0, \pm 1, \pm 2, \dots \quad (11)$$

and the corresponding eigenfunctions

$$u_{mn}(x, y) = \sqrt{2} \exp[-i(\arg \alpha + 2m\pi)x] \cdot \sin\left(n\pi - \frac{\pi}{2}\right)y \quad (12)$$

which form an orthonormal basis of space $L^2(0,1)$.

Proof. Let $Su(x, y) = u(x, 1-y)$, then

$$iu_x(x, y) + u_y(x, 1-y) = \left(i \frac{\partial}{\partial x} + S \frac{\partial}{\partial y} \right) u(x, y)$$

we use this formula in the calculations.

$$\text{Let } u_{mn}(x, y) = \sqrt{2} \exp[-i(\arg \alpha + 2m\pi)x] \cdot \sin\left(n\pi - \frac{\pi}{2}\right)y, \quad n = 1, 2, \dots, m = 0, \pm 1, \pm 2, \dots$$

Then the following formulas hold:

$$\begin{aligned} i \frac{\partial}{\partial x} u_{mn}(x, y) &= (\arg \alpha + 2m\pi) u_{mn}(x, y); \\ S \frac{\partial}{\partial y} u_{mn}(x, y) &= \sqrt{2} \left(n\pi - \frac{\pi}{2} \right) \exp[-i(\arg \alpha + 2m\pi)] \cdot \cos\left(n\pi - \frac{\pi}{2}\right)(1-y) = \\ \sqrt{2} \left(n\pi - \frac{\pi}{2} \right) \exp[-i(\arg \alpha + 2m\pi)x] \cdot (-1)^{n+1} \sin\left(n\pi - \frac{\pi}{2}\right)y &= (-1)^{n+1} \left(n\pi - \frac{\pi}{2} \right) u_{mn}(x, y), \\ \left(i \frac{\partial}{\partial x} + S \frac{\partial}{\partial y} \right) u_{mn}(x, y) &= \left[\arg \alpha + 2m\pi + (-1)^{n+1} \left(n\pi - \frac{\pi}{2} \right) \right] u_{mn}(x, y). \end{aligned}$$

$$\text{Consequently, } i \frac{\partial}{\partial x} u_{mn}(x, y) + u_{mn}(x, 1-y) = \lambda_{mn} u_{mn}(x, y),$$

where

$$\lambda_{mn} = \arg \alpha + 2m\pi + (-1)^{n+1} \left(n\pi - \frac{\pi}{2} \right), \quad n = 1, 2, \dots, m = 0, \pm 1, \pm 2, \dots$$

Theorem 2. The operator

$$Lu = iu_x(x, y) + u_y(x, 1-y), \quad (13)$$

$$D(L) = \{u \in C^{1,1}(\Omega) \cap C(\bar{\Omega}) : u|_{y=0} = 0, u|_{x=0} = \alpha u|_{x=1}, |\alpha| = 1\} \quad (14)$$

is essentially self-adjoint in the space $L^2(\Omega)$.

Proof. This theorem is a simple consequence of Theorem 1, Lemma 2.3 and Lemma 2.4.

Up to the present time, we have not deliberately spoken about the spectrum of the operator L , because this concept is inherent only in a closed operator, and our operator has until now been not closed. In virtue of the Theorem 2 the equality $\bar{L} = L^*$ holds.

Further, by an operator L we mean the closure of the operator (13) – (14) and investigate its spectrum. The eigenvalues of this operator have the form:

$$\lambda_{mn} = \arg \alpha + 2m\pi + (-1)^{n+1} \left(n\pi - \frac{\pi}{2} \right), \quad n = 1, 2, \dots, m = 0, \pm 1, \pm 2, \dots$$

a) Let $n = 2k, k = 1, 2, \dots$, then

$$\lambda_{m,2k} = \arg \alpha + 2m\pi - \left(2k\pi - \frac{\pi}{2} \right) = \pi \left[\frac{\arg \alpha}{\pi} + 2(m-k) + \frac{1}{2} \right] = \pi \left[\frac{\arg \alpha}{\pi} + \frac{1}{2} + 2(m-k) \right].$$

The value $2(m-k)$ runs through all even numbers, the argument α lies within $0 \leq \arg \alpha < 2\pi$, therefore

$$0 \leq \frac{\arg \alpha}{\pi} < 2, \quad \frac{1}{2} \leq \frac{\arg \alpha}{\pi} + \frac{1}{2} < \frac{5}{2}.$$

Between the numbers $\frac{1}{2}$ and $\frac{5}{2}$ there is only one even number 2, which is reached at $\frac{\arg \alpha}{\pi} = \frac{3}{2}$.

b) Let $n = 2k-1, k = 1, 2, \dots$, then

$$\lambda_{m,2k-1} = \arg \alpha + 2m\pi(2k-1)\pi - \frac{\pi}{2} = \pi \left[\frac{\arg \alpha}{\pi} + 2m + 2k - 1 - \frac{1}{2} \right] = \pi \left[\frac{\arg \alpha}{\pi} - \frac{3}{2} + 2(m+k) \right];$$

The quantity $2(m+k)$ runs through all even integers. The following inequality holds:

$$-\frac{3}{2} \leq \frac{\arg \alpha}{\pi} - \frac{3}{2} < \frac{1}{2}.$$

Between the numbers $-\frac{3}{2}$ and $\frac{1}{2}$ there is only one even number 0, which is reached when

$$\frac{\arg \alpha}{\pi} = \frac{3}{2}.$$

Theorem 3. The spectrum of the operator (13)-(14) consists of two series of infinite eigenvalues:

a) $\lambda_m^{(1)} = \pi \left[\frac{\arg \alpha}{\pi} + \frac{1}{2} + 2m \right], m = 0, \pm 1, \pm 2, \dots;$

b) $\lambda_m^{(2)} = \pi \left[\frac{\arg \alpha}{\pi} - \frac{3}{2} + 2m \right], m = 0, \pm 1, \pm 2, \dots$

i.e. each of these values is taken an infinite number of times, the corresponding eigenfunctions form an orthonormal basis of the space $L^2(\Omega)$.

The inverse operator L^{-1} exists if and only if

$$\frac{\arg \alpha}{\pi} \neq \frac{3}{2} \quad (15)$$

Theorem 4. The boundary-value problem

$$\begin{aligned} Lu = iu_x(x, y) + u_y(x, 1-y) &= f(x, y), (x, y) \in \Omega; \\ u|_{y=0} &= 0, \quad u|_{x=0} = \alpha u|_{x=1}, \quad |\alpha| = 1, \end{aligned}$$

is strongly solvable in space $L^2(\Omega)$ if and only if

$$\frac{\arg \alpha}{\pi} \neq \frac{3}{2}.$$

When condition (15) is satisfied, the inverse operator $(\bar{L})^{-1}$ exists, is bounded, but not compact, since there is a continuous spectrum of the operator \bar{L} .

Theorem 5. The spectral problem (1)-(3) has an infinite set of eigenvalues

$$\lambda_{mn} = \arg \alpha + 2m\pi + (-1)^{n+1} \left(n\pi - \frac{\pi}{2} \right), \quad n = 1, 2, \dots, \quad m = 0, \pm 1, \pm 2, \dots,$$

and the corresponding eigenfunctions

$$u_{mn}(x, y) = \sqrt{2} \exp[-i(\arg \alpha + 2m\pi)x] \cdot \sin\left(n\pi - \frac{\pi}{2}\right)y,$$

which form an orthonormal basis of space $L_2(\Omega)$.

The spectrum of the beam of operators (1)-(3) consists of two series of infinite eigenvalues:

a) $\lambda_m^{(1)} = \pi \left[\frac{\arg \alpha}{\pi} + \frac{1}{2} + 2m \right], \quad m = 0, \pm 1, \pm 2, \dots;$

b) $\lambda_m^{(2)} = \pi \left[\frac{\arg \alpha}{\pi} - \frac{3}{2} + 2m \right], \quad m = 0, \pm 1, \pm 2, \dots$

each of these values is taken an infinite number of times, the corresponding eigenfunctions form an orthonormal basis of the space $L^2(\Omega)$.

The proof of the theorem follows easily from the above Theorems 1 - 4. For this it is sufficient to note that the equation coincides with the equation (1), where

$$Lu = iu_x(x, y) + u_y(x, 1-y) = i \left(\frac{\partial}{\partial x} + S \frac{\partial}{\partial x} \right) u(x, y),$$

and $Su(x, y) = u(x, 1-y)$.

4. Conclusions.

Note that the operator (13)-(14) is a two-dimensional generalization of the operator discussed in [16] - [17], which has found application to singularly perturbed Cauchy problem [18], to the operator of the heat conductivity in [19], and to the ill-posed problems of mathematical physics in [20] - [21], and Volterra problems in [26] - [27].

The spectral properties of the wave operator change dramatically if it is perturbed by the low order term containing the spectral parameter, in particular, it turns out to be reversible for certain values of the coefficient of the boundary condition. Zero is an infinite eigenvalue of the Dirichlet problem of the wave equation. Adding a low-order term with a spectral parameter and expanding the domain of definition change the situation, the operator becomes reversible.

Authors thank the leadership of the International Silkway University and the Regional Social Innovation University of Shymkent for the partial funding of the research.

А.Ш.Шалданбаев¹, М.И.Ақылбаев², А.А.Шалданбаева², А.Ж.Бейсебаева³

¹Халықаралық Silkway университеті, Шымкент, Қазақстан;

²Аймақтық әлеуметтік-инновациялық университеті, Шымкент, Қазақстан;

³М.О.Әуезов атындағы Оңтүстік Қазақстан мемлекеттік университеті, Шымкент, Қазақстан

КІШІ МУШЕЛІ ТОЛҚЫН ОПЕРАТОРЫНЫҢ СПЕКТРЭЛДІК ҚАСИЕТТЕРИ

Аннотация. Толқындық кішік оператордың жайсыз екені көпке мәлім, себебі нөл нүктесі шексіз еселі меншікті мән. Біздің зерттеулеріміздің көрсетуінше, жағдайды өзгертуге болады, бұл, үшін операторды спектрелді параметрге көбейтілген кіші мүшемен тітіркендіру жеткілікті, сонда есебіміз операторлар шоғырының кейіпіне енеді. Бұл операторлар шоғыры қарапайым операторлардың көбейтіндісіне жіктеледі, ал ол операторлар оңай зерттеледі.

Тірек сөздер: ауытқыған аргумент, әлді шешілу, спектр, операторлар шоғыры, функционал-дифференциал оператор.

УДК 517. 929

А.Ш. Шалданбаев¹, М.И. Ақылбаев², А.А. Шалданбаева², А.Ж. Бейсебаева³

¹Международный университет Silkway, г. Шымкент, Казахстан;

²Региональный социально-инновационный университет, г. Шымкент, Казахстан;

³Южно-Казахстанский Государственный университет им. М.Ауезова, г.Шымкент, Казахстан

О СПЕКТРАЛЬНЫХ СВОЙСТВАХ ВОЛНОВОГО ОПЕРАТОРА, ВОЗМУЩЁННОГО МЛАДШИМ ЧЛЕНОМ

Аннотация. Некорректность минимального волнового оператора общеизвестна, так как нуль для него является бесконечнократным собственным значением. Как показали наши исследования, положение изменится, если возмутить его младшим членом, содержащим в качестве коэффициента спектральный параметр, в итоге задача принимает вид операторного пучка. Полученный пучок операторов легко факторизуется с помощью функционально–дифференциальных операторов первого порядка, спектральные свойства которых легко изучаются классическим методом разделения переменных. Непосредственное применение метода разделения переменных к исходному пучку операторов наталкивается на непреодолимые трудности.

Ключевые слова: отклоняющиеся аргумент, сильная разрешимость, спектр, пучок операторов, функционально-дифференциальный оператор

Information about authors:

Shaldanbayev A. Sh. - doctor of physico-mathematical Sciences, associate Professor, head of the center for mathematical modeling Silkway International University, Shymkent, Kazakhstan; <http://orcid.org/0000-0002-7577-8402>;

Akylbaev M. I. - candidate of technical Sciences, associate Professor, Vice-rector for research of the Regional socio-innovative University, Shymkent, Kazakhstan, <https://orcid.org/0000-0003-1383-4592>;

Shaldanbayeva A.A. - master of science, "Regional Social-Innovative University", Shymkent; <https://orcid.org/0000-0003-2667-3097>;

Beisebayeva A.Zh. - senior lecturer of Mathematics, South Kazakhstan state University named after M.O.Auezova, Shymkent, Kazakhstan, <https://orcid.org/0000-0003-4839-9156>

REFERENCES

- [1] J. Hadamard. On some topics connected with linear partial differential equations, Proc.Benares Math. Soc, 1921, 3, p. 39-48.
- [2] D.G. Bourgin, R. Duffin. The Dirichlet problem for the vibrating string equation. Bull.Amer.Math.Soc., 45 (12) (1939), pp.851-858.
- [3] V.I. Arnold. Small denominators. I: Mappings of the circumference onto itself. Izv. Akad. Nauk SSSR Ser. Mat., 25 (1) (1961), pp. 21–86 [Am. Math. Soc., Transl., II. Ser. 46, 213284 (1965)].
- [4] O.I. Bobik, P.I. Bondarchuk, B.I. Ptashnik. Elements of the Qualitative Theory of Partial Differential Equations. Kiev, 1972. 175 p.(In Russian).

- [5] B. D. Sleeman. Boundary value problems for the vibrating string equation. Proc. Roy.Soc. Edinburgh, 1981, v. A-88, Us. 1-2, p. 185-194. (in Eng.).
- [6] P. P. Mosolov. The Dirichlet problem for partial differential equations, Izv. Vyssh. Uchebn. Zaved. Mat., 1960, no. 3, 213-218.(In Russian).
- [7] S.L.Sobolev. An example of a correct boundary-value problem for the equation of oscillation of a string with data on the entire boundary. Dokl. AN SSSR, 1956, v. 109, No. 4, p. 707–709. (In Russian).
- [8] Yu.M. Berezanskii. Expansions in Eigenfunctions of Self-Adjoint Operators. Kiev: Naukova Dumka (1965), 799 pp. [in Russian].
- [9] R.A. Aleksandryan. Spectral properties of operators arising from systems of differential equations of Sobolev type, Tr. Mosk. Mat. Obs., 9, 1960, 455-505. (In Russian)
- [10] T.Sh. Kal'menov. Regular boundary value problems for the wave equation, Differ. Uravn., 17:6 (1981), 1105-1121. [in Russian].
- [11] T.Sh. Kal'menov. On the Spectrum of a Self-Adjoint Problem for the Wave Equation, Vestnik Akad. Nauk KazSSR, 1982, N2, p.63–66.(In Russian).
- [12] T.Sh. Kal'menov, M.A. Sadybekov. The Dirichlet problem and nonlocal boundary value problems for the wave equation, Differential Equations, 26:1 (1990), 55-59. (In Russian).
- [13] V.P. Burskiy. Boundary value problems for a second-order hyperbolic equation in a disk, Izv. Vyssh. Uchebn. Zaved. Mat., 1987, no. 2, 22-29. (In Russian).
- [14] V.P. Burskiy. On the kernel of a differential operator with constant coefficients of lower order in the disk. -VINITI, No. 3792-82 Dep., 1982. (In Russian).
- [15] G.A. Sargsyan. Polynomial solutions of the Dirichlet problem for hyperbolic equations with constant coefficients in the disk, Reports of the National Academy of Sciences of Armenia, Volume 112, 2012, N 4. (In Russian).
- [16] T.Sh. Kal'menov, S.T. Akhmetova, A.Sh. Shaldanbaev. To the spectral theory of equations with deviating argument // Mathematical Journal, Almaty, 2004, Vol.4, No. 3 (13). Pp. 41-48. (In Russian).
- [17] M. T. Shomanbayeva. On the basis property of the eigenfunctions of the heat equation with a deviating argument, "Poisk", Almaty, 2, 2006. (In Russian).
- [18] T. Sh. Kal'menov, A.Sh. Shaldanbaev. On a recurrence method for solving a singularly perturbed Cauchy problem for a second order equation, Mat. Tr., 13:2 (2010), 128-138. (In Russian).
- [19] Orazov, A. Shaldanbayev, and M. Shomanbayeva. About the nature of the spectrum of the periodic problem for the heat equation with a deviating argument. Hindawi Publishing Corporation. Abstract and Applied Analysis. Volume 2013, Article ID 128363, 6 pages. (in Eng.).
- [20] T.Sh. Kal'menov and A.Sh. Shaldanbaev. On a criterion of solvability of the inverse problem of heat conduction. Journal of Inverse and Ill-Posed Problems 18:4, 352-369 (2010). (in Eng.).
- [21] Sh. Shaldanbayev, I. Orazov, and M.T. Shomanbayeva. On the restoration of an operator of Sturm- Liouville by one spectrum. AIP Conference Proceedings 1611, 53 (2014); doi: 10.1063/1.4893803. (in Eng.).
- [22] M. Reed and B. Simon. Methods of Modern Mathematical Physics-vol. 1-Functional Analysis, Academic Press, London, UK, 1972. (in Eng.).
- [23] S.G. Mikhlin. Course of Mathematical Physics, M., "Nauka", 1968. 576 pp. [in Russian].
- [24] V.A. Marchenko. Operators of Sturm-Liouville and their applications. Kiev: Naukova Dumka (1977), 330 pp.[in Russian].
- [25] A.N. Kolmogorov, S.V. Fomin. Elements of the theory of functions and functional analysis. M., "Nauka". 1976, 543 pp. [in Russian].
- [26] Akylbayev M.I., Beysebayeva A., Shaldanbayev A. Sh. On the Periodic Solution of the Goursat Problem for a Wave Equation of a Special Form with Variable Coefficients. News of the National Academy of Sciences of the republic of Kazakhstan. Volume 1, Number 317 (2018), 34 – 50. (in Eng.).
- [27] Shaldanbaeva A. A., Akylbayev M.I., Shaldanbaev A. Sh., Beisebaeva A.Zh. The Spectral Decomposition of Cauchy Problem's Solution for Laplace Equation. News of the National Academy of Sciences of the republic of Kazakhstan. Volume 5, Number 321 (2018), 75 – 87. <https://doi.org/10.32014/2018.2518-1726.10> (in Eng.).

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

www:nauka-nanrk.kz

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы *M. С. Ахметова, Т.А. Апендиев, Д.С. Алеков*
Верстка на компьютере *A.M. Кульгинбаевой*

Подписано в печать 10.08.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
9,6 п.л. Тираж 300. Заказ 4.

*Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19*