

**ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)**

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Әль-фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Казахский национальный университет
имени Аль-фараби

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-farabi kazakh
national university

SERIES
PHYSICO-MATHEMATICAL

4 (326)

JULY-AUGUST 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Бас редакторы
ф.-м.ғ.д., проф., КР ҮФА академигі **F.M. Мұтанов**

Редакция алқасы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев Ү.Ү. проф. корр.-мүшесі (Қазақстан)
Жусіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Ә. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорус)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«КР ҮФА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Үлттық ғылым академиясы» РКБ (Алматы қ.)
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы қуәлік

Мерзімділігі: жылдана 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Үлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Р е д а к ц и о н на я кол л е г и я:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жантаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф. чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

E d i t o r i n c h i e f
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d:

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskyi I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

N E W S

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.46>

Volume 4, Number 326 (2019), 76 – 82

UDK 517.958

B. Bekbolat¹, B. Kanguzhin², N. Tokmagambetov³

^{1,2,3}Al-Farabi Kazakh National University, Almaty, Kazakhstan;

³Ghent University, Ghent, Belgium;

^{1,2,3}Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan
bekbolat@math.kz, kanbalta@mail.ru, niyaz.tokmagambetov@gmail.com

**TO THE QUESTION OF A MULTIPoint MIXED
BOUNDARY VALUE PROBLEM FOR A WAVE EQUATION**

Abstract. It is well known that some problems in mechanics and physics lead to partial differential equations of the hyperbolic type. A classical example of the hyperbolic type is wave equation. When posed, the task sometimes lacks the classical boundary condition and the need arises to have a nonlocal boundary condition. Aim our work is get D'Alembert formula for mixed boundary value problem generated by a wave equation. In the classical case, given D'Alembert formula for boundary value problem generated by a wave equation. In our case, we must give D'Alembert formula for mixed boundary value problem. For this, we consider ordinary differential operator \mathcal{L} with non-local boundary conditions. We search the solution of the wave equation like a sum with eigenfunction of the operator \mathcal{L} . There are we use that fact, that eigenfunction of the operator \mathcal{L} is Riesz basis in $L^2(0, l)$. Through this method and calculation we get D'Alembert formula.

Key words: D'Alembert formula, wave equation, mixed boundary value problem, nonlocal boundary condition.

1 Introduction

It is well known that some problems in mechanics and physics lead to partial differential equations of the hyperbolic type. When posed, the task sometimes lacks the classical boundary condition and the need arises to have a nonlocal boundary condition (see, [2, 3, 4, 5]). A simple example of such nonlocal conditions are multipoint conditions relating the value of the solution at the boundary points with the values at some interior points. For example, we refer the reader to [6, 7, 8, 9, 10].

2 Main result

Let us consider mixed boundary value problem generated by the homogeneous wave equation

$$\frac{\partial^2 U}{\partial t^2} - \frac{\partial^2 U}{\partial x^2} = 0, (x, t) \in S \quad (1)$$

with inhomogeneous initial data

$$U(x, 0) = f(x), U_t(x, 0) = F(x), x \in [0, l], \quad (2)$$

and with non-local conditions

$$U(0, t) = 0, \sum_{j=0}^N \alpha_j U_x'(x_j, t) = 0, \quad (3)$$

where $S = \{(x, t) : 0 < x < l, t > 0\}$,

$$0 = x_0 < x_1 < \dots < x_N = l, \alpha_0 \neq 0, \alpha_N \neq 0, \sum_{j=0}^N \alpha_j = 1, l < \infty,$$

the system of point $\{x_j\}_{j=0}^N$ on the segment $[0, l]$ is chosen such that the relation $\frac{x_j}{x_{j+1}}$ is a rational number for all $j \geq 0$.

Also consider an ordinary differential operator \mathcal{L} with the expression

$$\mathcal{L}(y) \equiv -y''(x), 0 < x < b, \quad (4)$$

with non-local boundary conditions

$$y(0) = 0, \sum_{j=0}^N \alpha_j y'(x_j) = 0. \quad (5)$$

By $\{\lambda_k\}_{k=1}^\infty$ denote eigenvalues of \mathcal{L} , which are zeros of the characteristic function

$$\Phi(\lambda) = \sum_{j=0}^N \alpha_j \cos \sqrt{\lambda} x_j.$$

Theorem 1 A solution of the boundary value problem (1)-(3) has the form

$$U(x, t) = \frac{\tilde{f}(x+t) + \tilde{f}(x-t)}{2} + \frac{1}{2} \int_{x-t}^{x+t} \tilde{F}(\tau) d\tau.$$

Proof. The system of eigen- and associated functions has the form

$$\{X_{kn}(x) = \frac{1}{k!} \frac{\partial^k}{\partial \lambda^k} (\frac{\sin \sqrt{\lambda} x}{\sqrt{\lambda}})|_{\lambda=\lambda_n}, k = 0, 1, \dots, m_n - 1\}_{n=1}^\infty, \quad (6)$$

which is the Riesz basis in $L^2(0, l)$ (see [1]). Here m_n is a multiplicity of the corresponding eigenvalue λ_n for all $n \in \mathbb{N}$.

Using this fact, let us prove solvability of the problem (1)-(3). The solution of the problem (1)-(3) we will seek in the view

$$U(x, t) = \sum_{n=1}^\infty \sum_{k=0}^{m_n-1} d_{kn}(t) X_{kn}(x), \quad (7)$$

where d_{kn} is a coefficient of the Fourier decomposition of U , depends on second argument.

By differentiating twice respect to x (7), we get

$$\begin{aligned} \frac{\partial^2 U}{\partial x^2} &= \sum_{n=1}^\infty \sum_{k=0}^{m_n-1} d_{kn}(t) [-\lambda_n X_{kn}(x) - X_{k-1,n}(x)] = \\ &= \sum_{n=1}^\infty \sum_{k=0}^{m_n-1} [-\lambda_n d_{kn}(t) - d_{k+1,n}(t)] X_{kn}(x), \end{aligned} \quad (8)$$

where $d_{m_n,n} = 0$.

By differentiating twice respect to t (7), we have

$$\frac{\partial^2 U}{\partial t^2} = \sum_{n=1}^\infty \sum_{k=0}^{m_n-1} d_{kn}''(t) X_{kn}(x), \quad (9)$$

By using the condition $U(0, t) = 0$, from (8)-(9) we take

$$d_{kn}''(t) = -\lambda_n d_{kn}(t) - d_{k+1,n}(t), k < m, \text{ where } d_{m_n,n} = 0. \quad (10)$$

Use initial conditions (2):

$$U(x, 0) = \sum_{n=1}^{\infty} \sum_{k=0}^{m_n-1} d_{kn}(0) X_{kn}(x) = f(x),$$

$$U_t(x, 0) = \sum_{n=1}^{\infty} \sum_{k=0}^{m_n-1} d_{kn}'(0) X_{kn}(x) = F(x).$$

We calculate Fourier coefficients using the biorthogonal system of $\{X_{kn}(x)\}$ by formulas

$$\begin{aligned} d_{kn}(0) &= d_{kn}^f = \int_0^b f(x) \overline{h_{kn}(x)} dx, \\ d_{kn}'(0) &= d_{kn}^F = \int_0^b F(x) \overline{h_{kn}(x)} dx. \end{aligned} \quad (11)$$

Indeed, (10) – (11) is a Cauchy problem, and it's solution has the form

$$d_{kn}(t) = d_{kn}^f \cos \sqrt{\lambda_n} t + d_{kn}^F \frac{\sin \sqrt{\lambda_n} t}{\sqrt{\lambda_n}} - \int_0^t \frac{\sin \sqrt{\lambda_n} (t-\xi)}{\sqrt{\lambda_n}} d_{k+1,n} d\xi. \quad (12)$$

Taking into account $d_{m_n,n} = 0$, for $k = m_n - 1$ we have

$$d_{m_n-1,n}(t) = d_{m_n-1,n}^f \cos \sqrt{\lambda_n} t + d_{m_n-1,n}^F \frac{\sin \sqrt{\lambda_n} t}{\sqrt{\lambda_n}}.$$

And for $k = m_n - 2$ we have

$$\begin{aligned} d_{m_n-2,n}(t) &= d_{m_n-2,n}^f \cos \sqrt{\lambda_n} t + \\ &d_{m_n-2,n}^F \frac{\sin \sqrt{\lambda_n} t}{\sqrt{\lambda_n}} - \int_0^t \frac{\sin \sqrt{\lambda_n} (t-\zeta)}{\sqrt{\lambda_n}} d_{m_n-1,n}(t) d\zeta \\ &= d_{m_n-2,n}^f \cos \sqrt{\lambda_n} t - \int_0^t \frac{\sin \sqrt{\lambda_n} (t-\zeta)}{\sqrt{\lambda_n}} d_{m_n-1,n}^f \cos \sqrt{\lambda_n} \zeta d\zeta \\ &+ d_{m_n-2,n}^F \frac{\sin \sqrt{\lambda_n} t}{\sqrt{\lambda_n}} - \int_0^t \frac{\sin \sqrt{\lambda_n} (t-\zeta)}{\sqrt{\lambda_n}} d_{m_n-1,n}^F \frac{\sin \sqrt{\lambda_n} \zeta}{\sqrt{\lambda_n}} d\zeta. \end{aligned}$$

We denote

$$C_{0n}(t) = \cos \sqrt{\lambda_n} t, S_{0n}(t) = \frac{\sin \sqrt{\lambda_n} t}{\sqrt{\lambda_n}},$$

$$C_{kn}(t) = - \int_0^t \frac{\sin \sqrt{\lambda_n} (t-\zeta)}{\sqrt{\lambda_n}} \cos \sqrt{\lambda_n} \zeta d\zeta,$$

$$S_{kn}(t) = - \int_0^t \frac{\sin \sqrt{\lambda_n} (t-\zeta)}{\sqrt{\lambda_n}} \frac{\sin \sqrt{\lambda_n} \zeta}{\sqrt{\lambda_n}} d\zeta,$$

and

$$\begin{aligned} C_{k+1,n}(t) &= \frac{1}{k!} \frac{\partial^{k+1}}{\partial \lambda^{k+1}} C_{0n}(t), \\ S_{k+1,n}(t) &= \frac{1}{k!} \frac{\partial^{k+1}}{\partial \lambda^{k+1}} S_{0n}(t). \end{aligned} \quad (13)$$

Therefore

$$d_{m_n-2,n}(t) = d_{m_n-2,n}^f \cos \sqrt{\lambda_n} t + d_{m_n-2,n}^F \frac{\sin \sqrt{\lambda_n} t}{\sqrt{\lambda_n}} -$$

$$d_{m_n-1,n}^f C_{1,n}(t) - d_{m_n-1,n}^F S_{1,n}(t).$$

Analogically for another k the function $d_{kn}(t)$ can be written as

$$d_{kn}(t) = \sum_{j=k}^{m_n-1} [d_{jn}^f C_{j-k,n}(t) + d_{jn}^F S_{j-k,n}(t)]. \quad (14)$$

By substituting found $d_{kn}(t)$ in (7), we get

$$\begin{aligned} U(x, t) &= \sum_{n=1}^{\infty} \sum_{k=0}^{m_n-1} \sum_{j=k}^{m_n-1} [d_{jn}^f C_{j-k,n}(t) + d_{jn}^F S_{j-k,n}(t)] X_{kn}(x) \\ &= \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} \{d_{jn}^f \sum_{k=0}^j C_{j-k,n}(t) X_{kn}(x) + d_{jn}^F \sum_{k=0}^j S_{j-k,n}(t) X_{kn}(x)\}. \end{aligned}$$

By virtue of (6) and (13), we have a new presentation

$$\begin{aligned} U(x, t) &= \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} \{d_{jn}^f \sum_{k=0}^j \frac{1}{(j-k)!} \frac{\partial^{j-k}}{\partial \lambda_n^{j-k}} (\cos \sqrt{\lambda_n} t) \frac{1}{k!} \frac{\partial^k}{\partial \lambda_n^k} \left(\frac{\sin \sqrt{\lambda_n} t}{\sqrt{\lambda_n}} \right) \\ &\quad + d_{jn}^F \sum_{k=0}^j \frac{1}{(j-k)!} \frac{\partial^{j-k}}{\partial \lambda_n^{j-k}} \left(\frac{\sin \sqrt{\lambda_n} t}{\sqrt{\lambda_n}} \right) \frac{1}{k!} \frac{\partial^k}{\partial \lambda_n^k} \left(\frac{\sin \sqrt{\lambda_n} x}{\sqrt{\lambda_n}} \right)\}. \end{aligned}$$

Syne $\frac{\sin \sqrt{\lambda_n} t}{\sqrt{\lambda_n}} = \int_0^t \cos \sqrt{\lambda_n} \tau d\tau$, then

$$\begin{aligned} U(x, t) &= \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^f \frac{1}{j!} \sum_{k=0}^j \frac{j!}{k! (j-k)!} \frac{\partial^{j-k}}{\partial \lambda_n^{j-k}} (\cos \sqrt{\lambda_n} t) \frac{\partial^k}{\partial \lambda_n^k} \left(\frac{\sin \sqrt{\lambda_n} x}{\sqrt{\lambda_n}} \right) \\ &\quad + \int_0^t d\tau \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^F \frac{1}{j!} \sum_{k=0}^j \frac{j!}{k! (j-k)!} \frac{\partial^{j-k}}{\partial \lambda_n^{j-k}} (\cos \sqrt{\lambda_n} \tau) \frac{\partial^k}{\partial \lambda_n^k} \left(\frac{\sin \sqrt{\lambda_n} x}{\sqrt{\lambda_n}} \right). \end{aligned}$$

And using $\sum_{k=0}^j C_k^j U^{(k)}(x) V^{(j-k)}(x) = (UV)^{(j)}$, we take

$$\begin{aligned} U(x, t) &= \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^f \frac{1}{j!} \frac{\partial^j}{\partial \lambda_n^j} (\cos \sqrt{\lambda_n} t) \frac{\sin \sqrt{\lambda_n} x}{\sqrt{\lambda_n}} \\ &\quad + \int_0^t d\tau \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^F \frac{1}{j!} \frac{\partial^j}{\partial \lambda_n^j} (\cos \sqrt{\lambda_n} \tau) \frac{\sin \sqrt{\lambda_n} x}{\sqrt{\lambda_n}} \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2} \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^f \frac{1}{j!} \frac{\partial^j}{\partial \lambda_n^j} \left(\frac{\sin \sqrt{\lambda_n}(x+t)}{\sqrt{\lambda_n}} \right) \\
 &+ \frac{1}{2} \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^f \frac{1}{j!} \frac{\partial^j}{\partial \lambda_n^j} \left(\frac{\sin \sqrt{\lambda_n}(x-t)}{\sqrt{\lambda_n}} \right) \\
 &+ \frac{1}{2} \int_0^t d\tau \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^F \frac{1}{j!} \frac{\partial^j}{\partial \lambda_n^j} \left(\frac{\sin \sqrt{\lambda_n}(x+\tau)}{\sqrt{\lambda_n}} \right) \\
 &+ \frac{1}{2} \int_0^t d\tau \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^F \frac{1}{j!} \frac{\partial^j}{\partial \lambda_n^j} \left(\frac{\sin \sqrt{\lambda_n}(x-\tau)}{\sqrt{\lambda_n}} \right).
 \end{aligned}$$

So,

$$\begin{aligned}
 U(x, t) &= \frac{1}{2} \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^f X_{jn}(x+t) + \frac{1}{2} \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^f X_{jn}(x-t) \\
 &+ \frac{1}{2} \int_x^{x+t} \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^F X_{jn}(\tau) d\tau + \frac{1}{2} \int_{x-t}^x \sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^F X_{jn}(\tau) d\tau.
 \end{aligned} \tag{15}$$

When $0 \leq x - t \leq x + t \leq b$, then sums of series are coincides with the initial data

$$\sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^f X_{jn}(x+t) = f(x+t),$$

$$\sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^f X_{jn}(x-t) = f(x-t),$$

$$\sum_{n=1}^{\infty} \sum_{j=0}^{m_n-1} d_{jn}^F X_{jn}(\tau) = F(\tau).$$

For $0 \leq x - t \leq x + t \leq b$ the solution is well-known D'Alembert formula

$$U(x, t) = \frac{f(x+t) + f(x-t)}{2} + \frac{1}{2} \int_{x-t}^{x+t} F(\tau) d\tau.$$

Thus, the formula (15) can be interpreted as a generalization of the D'Alembert formula for arbitrary $0 \leq x \leq b, t \geq 0$.

As a result, we conclude

$$U(x, t) = \frac{\tilde{f}(x+t) + \tilde{f}(x-t)}{2} + \frac{1}{2} \int_{x-t}^{x+t} \tilde{F}(\tau) d\tau,$$

where $\tilde{f}(x)$ and $\tilde{F}(x)$ extended from the segment $[0, l]$ to the whole real axis by the analytical continuation of the basis system

$$\{X_{kn}(x) = \frac{1}{k!} \frac{\partial^k}{\partial \lambda^k} \frac{\sin \sqrt{\lambda_n} x}{\sqrt{\lambda_n}}\}.$$

Since $\{X_{kn}(x)\}$ is defined on $(-\infty, +\infty)$, then we can always to continue functions $f(x)$ and $F(x)$ outside of the segment $[0, l]$. From [1] follows that the system $\{X_{kn}(x)\}$ is Riesz basis in $L_2(-\infty, +\infty)$.

ACKNOWLEDGEMENTS

The authors were supported by the Ministry of Education and Science of the Republic of Kazakhstan (MESRK) Grant AP05130994. No new data was collected or generated during the course of research.

Б. Бекболат¹, Б. Кангужин², Н. Токмагамбетов³,

^{1,2,3}Өл-Фараби атындағы ҚазҰУ, Алматы, Қазақстан;

³Гент университеті, Гент, Бельгия;

^{1,2,3}Математика және математикалық модельдеу институты, Алматы, Қазақстан

ТОЛҚЫН ТЕНДЕУІ ҮШІН ҚӨП НҮКТЕЛІ АРАЛАС ШЕКАРАЛЫҚ ЕСЕП

Аннотация.Механика мен физиканың кейір есептері дербес туындылы дифференциалдық тендеулердің гиперболалық түріне алып келетіні белгілі. Гиперболалық тендеудің классикалық өкіліне толқын тендеуі жатады. Кейде есеп шығару барысында тек шекаралық шарт жеткіліксіз болады, сондықтан қосымша локалды емес шекаралық шартта қолданылады. Біздің жұмыстың мақсаты толқын тендеуі арқылы туындаған аралас шекаралық есептің Даламбер формуласын табу. Классикада толқын тендеуі арқылы туындаған шекаралық есеп үшін Даламбер формуласы берілген. Біз аралас шекаралық есеп үшін Даламбер формуласын табу керекпіз. Ол үшін біз қосымша \mathcal{L} дифференциалдық операторын қарастырамыз. Себебі біз шешімді \mathcal{L} операторының меншікті функциялары арқылы құрылған катар арқылы іздеміз. Біз бұл жерде \mathcal{L} операторының меншікті функциялары $L^2(0, l)$ кеңістігінде Рисс базисы болатынын пайдаланамыз. Біз осы әдіс және есептеулер арқылы Даламбер формуласын аламыз.

Түйін сөздер:Даламбер формуласы, толқын тендеуі, аралас шекаралық есеп, локалды емес шекаралық шарт.

Б. Бекболат¹, Б. Кангужин², Н. Токмагамбетов³

^{1,2,3}КазНУ им. Аль-Фараби, Алматы, Казахстан;

³Гентский университет, Гент, Бельгия;

^{1,2,3}Институт математики и математического моделирования, Алматы, Казахстан

О МНОГОТОЧЕЧНОЙ ЗАДАЧЕ СМЕШАННОЙ ГРАНИЦЫ ДЛЯ ВОЛНОВОГО УРАВНЕНИЯ

Аннотация.Хорошо известно, что некоторые проблемы механики и физики приводят к уравнениям в частных производных гиперболического типа. Классическим примером гиперболического типа является волновое уравнение. При постановке задачи иногда не хватает классического граничного условия, и возникает необходимость иметь нелокальное граничное условие. Цель нашей работы - получить формулу Даламбера для смешанной краевой задачи, порожденной волновым уравнением. В классическом случае дана формула Даламбера для краевой задачи, порожденная волновым уравнением. В нашем случае мы должны дать формулу Даламбера для смешанной краевой задачи. Для этого рассмотрим обыкновенный дифференциальный оператор \mathcal{L} с нелокальными граничными условиями. Мы ищем решение волнового уравнения как сумму с собственной функцией оператора \mathcal{L} . Мы используем тот факт, что собственная функция оператора \mathcal{L} является базисом Рисса в $L^2(0, l)$. С помощью этого метода и расчета мы получаем формулу Даламбера.

Ключевые слова:Формула Даламбера, волновое уравнение, смешанная краевая задача, нелокальные краевые условия.

Information about authors:

Kanguzhin Baltabek – Doctor of Physical and Mathematical Sciences, Professor, Al-Farabi Kazakh National University, Almaty, Kazakhstan;

Tokmagambetov Niyaz – PhD doctor, Department of Mathematics: Analysis, Logic and Discrete Mathematics, Ghent University, Ghent, Belgium;

Bekbolat Bayan – Doctorate, al-Farabi Kazakh National University, Almaty, Kazakhstan.

REFERENCES

- [1] Gubreev G. M., *Spectral analysis of biorthogonal expansions of functions, and exponential series*, Izv. Akad. Nauk SSSR Ser. Mat., Vol.53, №6. 1989. P.1236–1268.
- [2] Kanguzhin B. E., Anijarov A. A., *Well-Posed Problems for the Laplace Operator in a Punctured Disk*, Mathematical Notes, Vol.89, №6. 2011. P.819–829.
- [3] Kanguzhin B. E., Tokmagambetov N. E., *On Regularized Trace Formulas for a Well-Posed Perturbation of the m-Laplace Operator*. Differential Equations. **51**:1583–1588 (2015).
- [4] Kanguzhin B. E., Tokmagambetov N. E., *Resolvents of well-posed problems for finite-rank perturbations of the polyharmonic operator in a punctured domain*. Siberian Mathematical Journal. **57**:265–273 (2016)
- [5] Kanguzhin B. E., Tokmagambetov N. E., *A regularized trace formula for a well-perturbed Laplace operator*, Doklady Mathematics. Vol.91, №1. 2015.P.1–4.
- [6] Bastys A., Ivanauskas F., Sapagovas M., *An explicit solution of a parabolic equation with nonlocal boundary conditions*, Lithuanian Mathematical Journal. –Vol.45, №3. 2005. P.257–271.
- [7] Zhanbing Bai, *On positive solutions of a nonlocal fractional boundary value problem*, Nonlinear analysis-theory methods and applications. Vol.72, №2. 2005. P.916–924.
- [8] Webb J. R. L., Gennaro I., *Positive solutions of nonlocal boundary value problems: A unified approach*, Jornal of the London Mathematical society series. Vol.74, №3. 2006. P.973–693.
- [9] Webb J. R. L., Gennaro I., *Positive solutions of nonlocal boundary value problems involving integral conditions*, Node-nonlinear differential equations and applications. Vol. 15, №1-2. 2008. P.45–67.
- [10] Goodrich, Christopher S., *Existence and uniqueness of solutions to a fractional difference equation with nonlocal conditions*, Computers and mathematics with applications. Vol.61, №2. 2011. P.191–202.
- [11] Seitmuratov A., Zharmenova B., Dauitbayeva A., Bekmuratova A. K., Tulegenova E., Ussenova G., *Numerical analysis of the solution of some oscillation problems by the decomposition method*, News of the national academy of sciences of the republic of Kazakhstan, Series physic-mathematical. Vol.323, №1.2019. P.28–37.ISSN 1991-346X
<https://doi.org/10.32014/2019.2518-1726.4>

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

www:nauka-nanrk.kz

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы *M. С. Ахметова, Т.А. Апендиев, Д.С. Алеков*
Верстка на компьютере *A.M. Кульгинбаевой*

Подписано в печать 10.08.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
9,6 п.л. Тираж 300. Заказ 4.

*Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19*