

ISSN 2518-1726 (Online),  
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ  
Әль-фараби атындағы Қазақ ұлттық университетінің

# Х А Б А Р Л А Р Ы

---

---

## ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН  
Қазақстан Республикасының Ғылым  
Академиясының Әль-Фараби  
атаулы университетінің

## NEWS

OF THE NATIONAL ACADEMY OF SCIENCES  
OF THE REPUBLIC OF KAZAKHSTAN  
Al-farabi kazakh  
national university

**SERIES**  
**PHYSICO-MATHEMATICAL**

**6(328)**

**NOVEMBER – DECEMBER 2019**

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Бас редакторы  
ф.-м.ғ.д., проф., ҚР ҰҒА академигі **Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

**Жұмаділдаев А.С.** проф., академик (Қазақстан)  
**Кальменов Т.Ш.** проф., академик (Қазақстан)  
**Жантаев Ж.Ш.** проф., корр.-мүшесі (Қазақстан)  
**Өмірбаев У.У.** проф. корр.-мүшесі(Қазақстан)  
**Жүсіпов М.А.** проф. (Қазақстан)  
**Жұмабаев Д.С.** проф. (Қазақстан)  
**Асанова А.Т.** проф. (Қазақстан)  
**Бошкаев К.А.** PhD докторы (Қазақстан)  
**Сұраған Д.** корр.-мүшесі (Қазақстан)  
**Quevedo Hernando** проф. (Мексика),  
**Джунушалиев В.Д.** проф. (Қырғыстан)  
**Вишневский И.Н.** проф., академик (Украина)  
**Ковалев А.М.** проф., академик (Украина)  
**Михалевич А.А.** проф., академик (Белорус)  
**Пашаев А.** проф., академик (Әзірбайжан)  
**Такибаев Н.Ж.** проф., академик (Қазақстан), бас ред. орынбасары  
**Тигиняну И.** проф., академик(Молдова)

**«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».**

**ISSN 2518-1726 (Online), ISSN 1991-346X (Print)**

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы»РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде  
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18, <http://physics-mathematics.kz/index.php/en/archive>

---

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор  
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Редакционная коллегия:

**Джумадильдаев А.С.** проф., академик (Казахстан)  
**Кальменов Т.Ш.** проф., академик (Казахстан)  
**Жангаев Ж.Ш.** проф., чл.-корр. (Казахстан)  
**Умирбаев У.У.** проф., чл.-корр. (Казахстан)  
**Жусупов М.А.** проф. (Казахстан)  
**Джумабаев Д.С.** проф. (Казахстан)  
**Асанова А.Т.** проф. (Казахстан)  
**Бошкаев К.А.** доктор PhD (Казахстан)  
**Сураган Д.** чл.-корр. (Казахстан)  
**Quevedo Hernando** проф. (Мексика),  
**Джунушалиев В.Д.** проф. (Кыргызстан)  
**Вишневский И.Н.** проф., академик (Украина)  
**Ковалев А.М.** проф., академик (Украина)  
**Михалевич А.А.** проф., академик (Беларусь)  
**Пашаев А.** проф., академик (Азербайджан)  
**Такибаев Н.Ж.** проф., академик (Казахстан), зам. гл. ред.  
**Тигиняну И.** проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов  
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

Editor in chief  
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l   b o a r d :

**Dzhumadildayev A.S.** prof., academician (Kazakhstan)  
**Kalmenov T.Sh.** prof., academician (Kazakhstan)  
**Zhantayev Zh.Sh.** prof., corr. member. (Kazakhstan)  
**Umirbayev U.U.** prof. corr. member. (Kazakhstan)  
**Zhusupov M.A.** prof. (Kazakhstan)  
**Dzhumabayev D.S.** prof. (Kazakhstan)  
**Asanova A.T.** prof. (Kazakhstan)  
**Boshkayev K.A.** PhD (Kazakhstan)  
**Suragan D.** corr. member. (Kazakhstan)  
**Quevedo Hernando** prof. (Mexico),  
**Dzhunushaliyev V.D.** prof. (Kyrgyzstan)  
**Vishnevskiy I.N.** prof., academician(Ukraine)  
**Kovalev A.M.** prof., academician(Ukraine)  
**Mikhalevich A.A.** prof., academician(Belarus)  
**Pashayev A.** prof., academician(Azerbaijan)  
**Takibayev N.Zh.** prof., academician (Kazakhstan), deputy editor in chief.  
**Tiginyanu I.** prof., academician(Moldova)

**News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.**

**ISSN 2518-1726 (Online), ISSN 1991-346X (Print)**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.80>

Volume 6, Number 328 (2019), 130 – 137

MPHTI 27.31.21

**M.B. Zhassybayeva, K.R. Yesmakhanova**

L.N. Gumilyov Eurasian National University, Astana, Kazakhstan  
E-mail: mzhassybaeva@yahoo.com, kryesmakhanova@gmail.com

**SOLITON SOLUTIONS FOR THE (2+1)-DIMENSIONAL  
INTEGRABLE FOKAS-LENELLS EQUATION**

**Abstract.** Studying of solitons led to the discovery of a number of new directions related to it. There is interest in which is also enhanced in connection with the discovery of new examples in which soliton processes are manifested. The number and variety of nonlinear equations containing solitons as the most interesting solutions significantly increase due to generalizations to the two-dimensional and three-dimensional cases. Such popular transformations as Darboux, Backlund and Hirota's bilinear method are often used to find exact different kind of the solutions of nonlinear equations.

In the present paper, we present Lax pair of the (2+1)-dimensional integrable Fokas-Lenells equation. The bilinear form of the (2+1)-dimensional integrable Fokas-Lenells equation was obtained by the Hirota's bilinear method. By using Hirota's bilinear method, we construct exact one-soliton and two-soliton solutions of the (2+1)-dimensional Fokas-Lenells equation. The graphics of the obtained solutions are presented. The obtained new results have important physical applications.

**Keywords:** Hirota method, Lax representation, soliton solution, Fokas-Lenells equation.

**Introduction.** In many areas of science, the object of intensive theoretical and experimental research is soliton which is mean the "solitary" wave (solitary wave). Soliton can be used to transmit information, where the main idea is to use in each bit interval to represent units in the stream of binary signals. Mathematically, solitons are localized stationary solutions of nonlinear partial differential equations or their generalizations.

One of the generalization of nonlinear Schrödinger is the integrable (1+1)-dimensional Fokas-Lenells equation (FL) [1, 2] which was proposed by J. Fokas and A.S. Lenells. At present, studying multidimensional integrable systems containing derivatives in more than two variables is very interesting and relevant and namely finding exact soliton solutions. In this regard, by considering the (1+1)-dimensional FL equation, we obtain the (2+1)-dimensional FL equation and present its Lax pair [1, 2].

The paper is organized as follows. In Section 2, we present Lax representation of the (2+1)-dimensional FL equation. Hirota method is presented in Section 3. Namely, we apply Hirota method for FL equation and find one soliton and two soliton solutions by obtained bilinear equation. In Section 4, we give conclusion.

**Lax representation.** The (2+1)-dimensional FL equation is given by next form

$$iq_{xt} - iq_{xy} + 2q_x - q_x |q|^2 + iq = 0, \quad (1)$$

where  $q$  is the complex shell of the field, the indices  $x$ ,  $y$  and  $t$  denote the partial derivatives with respect to the arguments  $x$ ,  $y$  and  $t$ , and  $i$  is the complex number.

To construct solutions of differential equations, a number of conditions must be fulfilled, one of which is to satisfy the compatibility condition [1]. The studied equation (1) satisfies the "compatibility condition" and has the following has a Lax representation

$$\begin{aligned}\Psi_x &= U\Psi, & U &= -i\lambda^2\sigma_3 + \lambda Q, \\ \Psi_t &= \Psi_y + W\Psi, & W &= W_0 + \frac{1}{\lambda}W_{-1} - \frac{i}{4\lambda^2}\sigma_3,\end{aligned}$$

where  $\Psi = \Psi(x,t)$  is a  $2 \times 2$  matrix-valued eigenfunction,  $\lambda$  is an isospectral parameter, and matrices are given in the form:

$$Q = \begin{pmatrix} 0 & q_x \\ r_x & 0 \end{pmatrix}, \quad W_0 = i\sigma_3 - \frac{iqr}{2}\sigma_3, \quad W_{-1} = \frac{i}{2} \begin{pmatrix} 0 & q \\ -r & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

**Hirota's bilinear method.** Soliton solutions of the (2+1)-dimensional FL equation will be constructed by the so-called bilinear Hirota's method. The outline for constructing soliton solutions by this method is as follows:

1. The first we apply the dependent variable transformation for FL equation in order to obtain its bilinear form.
2. The second we consider the formal series of perturbation theory.
3. The third we build multi soliton solutions.

**One-soliton solution of the (2+1)-dimensional FL equation.** In order to construct soliton solutions of equation (1), we use the bilinear form of FL equation has the form [6]  $g = \frac{g}{f}$ . Then the bilinear form of the FL equation

$$iD_x D_t (g \cdot f) - iD_x D_y (g \cdot f) + 2D_x (g \cdot f) + igf = 0, \quad (2a)$$

$$iD_x D_t (f \cdot f^*) - iD_x D_y (f \cdot f^*) - \frac{1}{2} D_x (g \cdot g^*) = 0, \quad (2b)$$

$$iD_t (f \cdot f^*) - iD_y (f \cdot f^*) - \frac{1}{2} g \cdot g^* = 0, \quad (2B)$$

$$iD_t (f \cdot f_x^*) - iD_y (f \cdot f_x^*) - \frac{1}{2} g \cdot g_x^* = 0, \quad (2r)$$

where  $g$  is complex function,  $f$  is real one, the “\*” sign means complex conjugation and  $D_x, D_y, D_t$ , are bilinear differential operators which defined by

$$D_x^m D_t^n f \cdot f^* = \left( \frac{\partial}{\partial x} - \frac{\partial}{\partial x'} \right)^m \left( \frac{\partial}{\partial t} - \frac{\partial}{\partial t'} \right)^n f(x,t) f^*(x',t') \Big|_{x'=x, t'=t},$$

where  $x', t'$  as two formal variables,  $f(x,t)$  and  $f^*(x',t')$  - two functions,  $m$  and  $n$  - two non-negative integers.

Then, we expand the functions  $g$  and  $f$  in bilinear equation (2) with respect to small parameter  $\varepsilon$  as follows [2]:

$$g(x,y,t) = \varepsilon g_1(x,y,t) + \varepsilon^3 g_3(x,y,t) + \dots, \quad (3a)$$

$$f(x,y,t) = 1 + \varepsilon^2 f_2(x,y,t) + \varepsilon^4 f_4(x,y,t) + \dots, \quad (3b)$$

where  $g_j$  is complex function,  $f_n$  is real ones ( $j = 1, 3, 5, \dots; n = 2, 4, 6, \dots$ ).

In the case of a one-soliton solution of equation (1), the formal parameter in equation (3) is taken as  $\mathcal{E} = 1$  and  $j = 1$ ,  $n = 2$ , i.e.

$$\mathbf{g} = \mathcal{E} \mathbf{g}_1, \quad (4a)$$

$$f = 1 + \mathcal{E}^2 f_2 \quad (4b)$$

and we find the solution according to the following statement:

$$q = \frac{\mathbf{g}}{f} = \frac{\mathbf{g}_1}{1 + f_2}, \quad (5)$$

where  $\mathbf{g}_1 = e^{\theta_1}$ ,  $\theta_1 = a_1 x + b_1 t + c_1 y + d_1$  and  $\theta_1^* = a_1^* x + b_1^* t + c_1^* y + d_1^*$ , here  $a_1, b_1, c_1, d_1$  - complex constants.

Substituting expression (4) into equation (2) and collecting in powers of the parameter  $\mathcal{E}$ , we get next system

$$\mathcal{E}^1 : (iD_x D_t - iD_x D_y + 2D_x)(\mathbf{g}_1 \cdot 1) + i\mathbf{g}_1 = 0, \quad (6)$$

$$\mathcal{E}^3 : (iD_x D_t - iD_x D_y + 2D_x)(\mathbf{g}_1 \cdot f_2) + i\mathbf{g}_1 f_2 = 0, \quad (7)$$

$$\mathcal{E}^2 : i(D_x D_t - D_x D_y)(f_2 + f_2^*) - \frac{1}{2} D_x(\mathbf{g}_1 \cdot \mathbf{g}_1^*) = 0, \quad (8)$$

$$\mathcal{E}^4 : iD_x D_t(f_2 \cdot f_2^*) - iD_x D_y(f_2 \cdot f_2^*) = 0, \quad (9)$$

$$\mathcal{E}^2 : i(D_t - D_y)(f_2 - f_2^*) - \frac{1}{2} \mathbf{g}_1 \cdot \mathbf{g}_1^* = 0, \quad (10)$$

$$\mathcal{E}^4 : i(D_t - D_y)(f_2 \cdot f_2^*) = 0, \quad (11)$$

$$\mathcal{E}^2 : (-iD_x D_t + iD_x D_y)(f_2^* \cdot 1) - \frac{1}{2} \mathbf{g}_1 \mathbf{g}_{1x}^* = 0, \quad (12)$$

$$\mathcal{E}^4 : i(D_x - D_y)(f_2 \cdot f_{2x}^*) = 0. \quad (13)$$

Now, applying the properties of the Hirota operator to equations (6) - (13), we obtain

$$\mathcal{E}^1 : i\mathbf{g}_{1tx} - i\mathbf{g}_{1yx} + 2\mathbf{g}_{1x} + i\mathbf{g}_1 = 0, \quad (14)$$

$$\begin{aligned} \mathcal{E}^3 : & i\mathbf{g}_{1tx} f_2 - i\mathbf{g}_{1t} f_{2x} - i\mathbf{g}_{1x} f_{2t} + i\mathbf{g}_1 f_{2tx} - i\mathbf{g}_{1yx} f_2 + i\mathbf{g}_{1y} f_{2x} + \\ & + i\mathbf{g}_{1x} f_{2y} - i\mathbf{g}_1 f_{2yx} + 2\mathbf{g}_{1x} f_2 - 2\mathbf{g}_1 f_{2x} + i\mathbf{g}_1 f_2 = 0 \end{aligned} \quad (15)$$

$$\mathcal{E}^2 : if_{2tx} + if_{2tx}^* - if_{2yx} - if_{2yx}^* - \frac{1}{2} \mathbf{g}_{1x} \mathbf{g}_1^* + \frac{1}{2} \mathbf{g}_1 \mathbf{g}_{1x}^* = 0, \quad (16)$$

$$\begin{aligned} \mathcal{E}^4 : & if_{2tx} f_2^* - if_{2t} f_{2x}^* - if_{2x} f_{2t}^* + if_2 f_{2tx}^* - if_{2yx} f_2^* + \\ & + if_{2y} f_{2x}^* + if_{2x} f_{2y}^* - if_2 f_{2yx}^* = 0, \end{aligned} \quad (17)$$

$$\varepsilon^2 : if_{2t} - if_{2t}^* - if_{2y} + if_{2y}^* - \frac{1}{2}g_1g_1^* = 0, \quad (18)$$

$$\varepsilon^4 : if_{2t}f_2^* - if_2f_{2t}^* - if_{2y}f_2^* + if_2f_{2y}^* = 0, \quad (19)$$

$$\varepsilon^2 : -if_{2xt}^* + if_{2xy}^* - \frac{1}{2}g_1g_{1x}^* = 0, \quad (20)$$

$$\varepsilon^4 : if_{2t}f_{2x}^* - if_2f_{2xt}^* - if_{2y}f_{2x}^* + if_2f_{2xy}^* = 0. \quad (21)$$

By solving the system of equation (14)-(21) we can get from equation (14)

$$a_1 = \frac{1}{c_1 - b_1 + 2i}, \quad (22)$$

the equation (15) gives conjugation form of  $a_1$

$$a_1^* = \frac{1}{c_1^* - b_1^* - 2i}, \quad (23)$$

and from equations (16) - (21) we obtain

$$f_2 = \frac{ia_1^2 a_1^*}{2(a_1 + a_1^*)^2} e^{\theta_1 + \theta_1^*}. \quad (24)$$

Then by substituting equation (24) into equation (5), we obtain the one-soliton solution of the (2+1)-dimensional FL equation, which is

$$q = \frac{e^{\theta_1}}{1 + \frac{ia_1^2 a_1^*}{2(a_1 + a_1^*)^2} e^{\theta_1 + \theta_1^*}} = \frac{2(a_1 + a_1^*)^2}{2(a_1 + a_1^*)^2 e^{-\theta_1} + ia_1^2 a_1^* e^{\theta_1^*}}, \quad (25)$$

or if rewrite equation (25), with  $\theta_1 = \kappa_1 + \chi_1$ , then we get

$$q = \frac{4\alpha_1^2 e^{i\chi_1}}{4\alpha_1^2 e^{-\kappa_1} + \frac{1}{2}(\alpha_1^2 + \beta_1^2)(i\alpha_1 - \beta_1)e^{\kappa_1}}, \quad (26)$$

where

$$\chi_1 = \beta_1 x + \nu_1 t + \tau_1 y + n_1, \quad \kappa_1 = \alpha_1 x + \mu_1 t + \sigma_1 y + m_1$$

with

$$a_1 = \alpha_1 + i\beta_1, \quad b_1 = \mu_1 + i\nu_1, \quad c_1 = \sigma_1 + i\tau_1, \quad d_1 = m_1 + in_1.$$

Finally, the equation (26) has the form



$$q = \frac{p^2 e^{i\chi_1}}{p\sqrt{h} \left( \frac{\sqrt{h}}{p} e^{\kappa_1} + \frac{p}{\sqrt{h}} e^{-\kappa_1} \right)}, \quad (27)$$

where  $p = 2\alpha_1$  and  $h = \frac{i}{2}(\alpha_1^2 + \beta_1^2)\alpha_1 - \frac{1}{2}(\alpha_1^2 + \beta_1^2)\beta_1$ .

In addition, when  $\frac{\sqrt{h}}{p} = e_1^\delta$  and  $\frac{p}{\sqrt{h}} = e^{-\delta_1}$  the equation (27) can be rewritten as

$$q = \frac{pe^{i\chi_1}}{2\sqrt{h}} \operatorname{sech}(\kappa_1 + \delta_1) = \frac{pe^{i\chi_1}}{2\sqrt{h} \cosh(\kappa_1 + \delta_1)}. \quad (28)$$

So, the equation (28) is one-soliton solution to the (2+1)-dimensional FL equation. Plot of the one-soliton solution is presented in Figure 1.




Figure 1 - Dynamics of the one-soliton solution with next parameters:  
 $a_1 = 1 - i$ ,  $c_1 = 1 - i$ ,  $b_1 = 1 - i$  and  $d_1 = 1 - i$ .  $t = -5$  (a);  $t = 0$  (b);  $t = 5$  (c).

**Two-soliton solution of the (2+1)-dimensional FL equation.** To find the two-soliton solution of equation (1), the formal parameter in equation (3) is taken, as  $\varepsilon = 1$  and  $j = 1, 3$ ,  $n = 2, 4$ , i.e.

$$\varepsilon^1 : (iD_x D_t - iD_x D_y + 2D_x)(g_1 \cdot 1) + ig_1 = 0, \quad (30)$$

$$\varepsilon^3 : (iD_x D_t - iD_x D_y + 2D_x)(g_1 \cdot f_2 + g_3 \cdot 1) + ig_1 f_2 + ig_3 = 0, \quad (31)$$

$$\varepsilon^5 : (iD_x D_t - iD_x D_y + 2D_x)(g_1 \cdot f_4 + g_3 \cdot f_2) + ig_1 f_4 + ig_3 f_2 = 0, \quad (32)$$

$$\varepsilon^7 : (iD_x D_t - iD_x D_y + 2D_x)(g_3 \cdot f_4) + ig_3 f_4 = 0, \quad (33)$$

$$\varepsilon^2 : i(D_x D_t - D_x D_y)(f_2 + f_2^*) - \frac{1}{2} D_x (g_1 \cdot g_1^*) = 0, \quad (34)$$

$$\varepsilon^4 : iD_x D_t (f_2 \cdot f_2^* + f_4 + f_4^*) - iD_x D_y (f_2 \cdot f_2^* + f_4 + f_4^*) -$$

$$-\frac{1}{2}D_x(g_1g_3^* + g_3g_1^*) = 0, \quad (35)$$

$$\begin{aligned} \varepsilon^6 : \quad & iD_xD_t(f_2 \cdot f_4^* + f_4 \cdot f_2^*) - iD_xD_y(f_2 \cdot f_4^* + f_4 \cdot f_2^*) - \\ & -\frac{1}{2}D_x(g_3g_3^*) = 0, \end{aligned} \quad (36)$$

$$\varepsilon^8 : \quad iD_xD_t(f_4 \cdot f_4^*) - iD_xD_y(f_4 \cdot f_4^*) = 0, \quad (37)$$

$$\varepsilon^2 : \quad i(D_t - D_y)(f_2 - f_2^*) - \frac{1}{2}g_1 \cdot g_1^* = 0, \quad (38)$$

$$\varepsilon^4 : \quad i(D_t - D_y)(f_2 \cdot f_2^*) + i(D_t - D_y)(f_4 - f_4^*) - \frac{1}{2}(g_1 \cdot g_3^* + g_1^*g_3) = 0, \quad (39)$$

$$\varepsilon^6 : \quad i(D_t - D_y)(f_2 \cdot f_4^*) - i(D_t - D_y)(f_2^* \cdot f_4) - \frac{1}{2}(g_3 \cdot g_3^*) = 0, \quad (40)$$

$$\varepsilon^8 : \quad i(D_t - D_y)(f_4 \cdot f_4^*) = 0. \quad (41)$$

$$\varepsilon^2 : \quad (-iD_xD_t + iD_xD_y)(f_2^* \cdot 1) - \frac{1}{2}g_1g_{1x}^* = 0, \quad (42)$$

$$\varepsilon^4 : \quad i(D_x - D_y)(f_2 \cdot f_{2x}^*) - i(f_{4xt}^* - f_{4xy}^*) - \frac{1}{2}(g_1g_{3x}^* + g_3g_{1x}^*) = 0, \quad (43)$$

$$\varepsilon^6 : \quad i(D_t - D_y)(f_2 \cdot f_{4x}^* + f_4 \cdot f_{2x}^*) - \frac{1}{2}g_3g_{3x}^* = 0, \quad (44)$$

$$\varepsilon^8 : \quad i(D_t - D_y)(f_4 \cdot f_{4x}^*) = 0. \quad (45)$$

By applying the properties of the Hirota operator we solve the system of equation (30)-(45), and can get the two-soliton solution of equation (1), which has the next form

$$q = \frac{g_1 + g_3}{1 + f_2 + f_4}, \quad (46)$$

where

$$\begin{aligned} g_1 &= e^{\theta_1} + e^{\theta_2}, \\ g_3 &= k_1 e^{\theta_1 + \theta_2 + \theta_1^*} + k_2 e^{\theta_1 + \theta_2 + \theta_2^*}, \\ f_2 &= l_1 e^{\theta_1 + \theta_1^*} + l_2 e^{\theta_1 + \theta_2^*} + l_3 e^{\theta_1^* + \theta_2} + l_4 e^{\theta_2 + \theta_2^*}, \\ f_4 &= m e^{\theta_1 + \theta_1^* + \theta_2 + \theta_2^*}, \end{aligned}$$

with

$$\theta_1 = a_1x + b_1t + c_1y + d_1, \quad \theta_1^* = a_1^*x + b_1^*t + c_1^*y + d_1^*,$$

$$\theta_2 = a_2 x + b_2 t + c_2 y + d_2, \quad \theta_2^* = a_2^* x + b_2^* t + c_2^* y + d_2^*,$$

$$l_1 = \frac{ia_1^2 a_1^*}{2(a_1 + a_1^*)^2}, \quad l_2 = \frac{ia_1^2 a_2^*}{2(a_1 + a_2^*)^2}, \quad l_3 = \frac{ia_2^2 a_1^*}{2(a_1^* + a_2)^2}, \quad l_4 = \frac{ia_2^2 a_2^*}{2(a_2 + a_2^*)^2},$$

$$k_1 = \frac{i(a_1^*)^3 (a_2 - a_1)^2}{2(a_1^* + a_2)^2 (a_1 + a_1^*)^2}, \quad k_2 = \frac{i(a_2^*)^3 (a_2 - a_1)^2}{2(a_1 + a_2^*)^2 (a_2 + a_2^*)^2},$$

$$m = -\frac{(a_2 - a_1)^2 (a_2^* - a_1^*)^2 (a_1 + a_2 + a_1^* + a_2^*) a_1^* a_2^* a_1^2 a_2^2}{2(a_1 + a_1^*)^2 (a_2 + a_2^*)^2 (a_1 + a_2^*)^2 (a_2 + a_1^*)^2},$$

and  $a_n, b_n, c_n, d_n$  - complex constants,  $n = 1, 2$ .




Figure 2 - Dynamics two-soliton solution with next parameters:

$a_1 = 0.5 + 0.5i$ ,  $c_1 = 0.5 + 0.5i$ ,  $b_1 = 0.5 + 0.5i$  and  $d_1 = 0.5 + 0.5i$ .  $t = -5$  (a);  $t = 0$  (б);  $t = 5$  (c).

**Conclusion.** Thus, we studied (2+1)-dimensional Fokas-Lenells equation by the Hirota's bilinear method which is considered one of the effective methods for finding exact solutions of integrable equations. By using this method, exact one-soliton and two-soliton solutions of the (2+1)-dimensional FL equation are constructed. Additionally, we present the graphical representation of the obtained soliton solutions.

**М.В. Жасыбаева, Қ.Р. Есмаханова**

Л.Н. Гумилев атындағы Еуразия ұлттық университеті, Астана, Қазақстан

### (2+1)-ӨЛШЕМДІ ИНТЕГРАЛДАНАТЫН ФОКАС-ЛЕНЭЛЛС ТЕНДЕУІНІҢ СОЛИТОНДЫ ШЕШІМДЕРІ

**Аннотация.** Солитондарды зерттеу солитонмен байланысты бірқатар жаңа бағыттардың ашылуына алып келді. Сонымен қатар, солитонды процестер байқалатын жаңа бағыттардың ашылуымен байланысты қызығушылықтар бар. Солитон құрайтын сызықты емес теңдеулердің саны мен алуандығы екі өлшемді және үш өлшемді жалпылауларға көшу арқылы күннен-күнге артуда. Дарбу, Бэклунд түрлендіруі және Хиротаның бисызықты әдісі сияқты танымал түрлендірулер сызықты емес теңдеулердің әр түрлі шешімдерін табу кезінде жиі қолданылады.

Демек, берілген жұмыс алдыңғы табылған (2+1)-өлшемді интегралданатын Фокас-Ленэллс теңдеуі және оның бисызықты түрі атты жұмысымыздың жалғасы болып табылады. Енді, зерттеулерімізді жалғастыра отырып, Хирота әдісі арқылы интегралданатын (2+1)-өлшемді Фокас-Ленэллс теңдеуінің 1-солитонды және 2-солитонды шешімдері табылып, графиктері тұрғызылды. Біздің тапқан нәтижелеріміз маңызды физикалық

қолданысқа ие.

**Түйін сөздер:** Хирота әдісі, Лакс көрінісі, солитонды шешім, Фокас-Ленэллс теңдеуі.

**М.Б. Жасыбаева, К.Р. Есмаханова**

Евразийский национальный университет им. Л.Н. Гумилева, Астана, Казахстан

### СОЛИТОННЫЕ РЕШЕНИЯ (2+1)-МЕРНОГО ИНТЕГРИРУЕМОГО УРАВНЕНИЯ ФОКАСА-ЛЕНЭЛЛСА

**Аннотация.** Успехи в исследовании солитонов привели к открытию целого ряда новых направлений связанных с ним, тем вдохновили бурной активностью исследователей в данных направлениях. Кроме того, интерес, к которому усиливается также в связи обнаружениями новых примеров, в которых проявляются солитонные процессы. Количество и разнообразие нелинейных уравнений, содержащих солитоны в качестве наиболее интересных решений, существенно увеличиваются благодаря обобщениям на двумерные и трехмерные случаи. Для нахождения солитонных решений нелинейных уравнений часто применяются такие популярные преобразования, как Дарбу, Бэклунда и метод Хироты.

Таким образом, данная работа является продолжением нашей предыдущей работы, в которой было найдено (2+1)-мерное интегрируемое уравнение Фокаса-Ленэллса и построена ее билинейная форма методом Хироты. Теперь, продолжая наши исследования, методом Хироты найдены его точные 1-солитонное и 2-солитонное решения с помощью уже полученной нами билинейной формы (2+1)-мерного уравнения Фокаса-Ленэллса и построены их графики. Найденные нами результаты имеют важные физические приложения.

**Ключевые слова:** метод Хироты, представление Лакса, солитонное решение, уравнение Фокаса-Ленэллса.

#### REFERENCES

- [1] Ablowitz M., Sigur H. Solitons and method of the inverse problem [Solitony I metod obratnoj zadachi]. M., 1987-479 p.
- [2] Zakharov V.E., Manakov S.V., Novikov S.P., Pitaevsky L.P. Introduction to the theory of integrable systems [Vvedeniye v teoriyu integriruemyyh sistem]. M: Nauka, 1980. -320 p. (In Russian).
- [3] Jingsong H., Shuwei X., Kuppuswamy P. Rogue Waves of the Fokas-Lenells Equation. Journal of the Physical Society of Japan. **81**, 124007 (2012).
- [4] Zhassybayeva M.B., Nugmanova G.N. Spin system equivalent to the integrable Fokas-Lenells equation [Spinovaya sistema, ehkivalentnaya integriruemomu uravneniyu Fokasa-Lenehllsa] Bulletin of L.N. Gumilyov Eurasian National University [Vestnik Evrazijskogo nacional'nogo universiteta imeni L.N. Gumileva]. 1(122), 53-57, 2018.
- [5] Tasmambetov Zh.N., Rajabov N., Ubayeva Zh.K. (2019) The coordinated solution of two differential equations in private derivatives of the third order \ \ Bulletin of national academy of sciences of the Republic of Kazakhstan. Volume 4, Number 326 (2019), PP.83-91. <https://doi.org/10.32014/2019.2518-1726.47> ISSN 2518-1726 (Online), ISSN 1991-346X (Print).
- [6] Zhassybayeva M., Yesmakhanova K. The construction of the (2+1)-dimensional integrable Fokas-Lenells equation and its bilinear form by Hirota method. International Conference on Technology, Engineering and Science. October 26-29, 2018, Antalya, Turkey [in print].
- [7] Matsuno Y.A. direct method of solution for the Fokas-Lenells derivative nonlinear Schrödinger equation: I. Bright soliton solutions. Journal of Physics A: Mathematical and Theoretical. 45, 23202 (2012).

## **Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

**ISSN 2518-1726 (Online), ISSN 1991-346X (Print)**

Редакторы *М. С. Ахметова, Т. А. Апендиев, Д. С. Аленов*  
Верстка на компьютере *А. М. Кульгинбаевой*

Подписано в печать 12.12.2019.  
Формат 60x881/8. Бумага офсетная. Печать – ризограф.  
9,6 п.л. Тираж 300. Заказ 6.