

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Әль-фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
Әль-Фараби атындағы
Қазақ ұлттық университетінің

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-farabi kazakh
national university

SERIES
PHYSICO-MATHEMATICAL

6(328)

NOVEMBER – DECEMBER 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Бас редакторы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі **Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев У.У. проф. корр.-мүшесі(Қазақстан)
Жүсіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Д. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қырғыстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорус)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик(Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы»РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18, <http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жангаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф., чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

Editor in chief
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d :

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskiy I.N. prof., academician(Ukraine)
Kovalev A.M. prof., academician(Ukraine)
Mikhalevich A.A. prof., academician(Belarus)
Pashayev A. prof., academician(Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician(Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.81>

Volume 6, Number 328 (2019), 138 – 142

UDC 510.67

MPHTH 27.03.66

B.S. Baizhanov¹, S.S. Baizhanov²

¹RSE Institute of Mathematics and Mathematical Modelling, Almaty, Kazakhstan;

²Al-Farabi Kazakh National University, Almaty, Kazakhstan

E-mail: baizhanov@math.kz, sayan-5225@mail.ru

SOME QUESTIONS ON EXTERNAL DEFINABILITY

Abstract. The article discusses the various approaches to the concept of external definability developed in o-minimal theories. An example of o-minimal theories shows how external constants help determine the existence of a solution in a model of a formula with external constants. The basic concepts are formulated with the help of which external definability is proved. A brief review of the results for dependent theories is given. In conclusion, sufficient conditions are formulated so that the NSOP theory has the some property of external definability. A brief explanation of the stated theorem is given.

Keywords: externally definable, neighborhood of tuple of the set in the type, non orthogonality of two types.

External definability. Let \mathfrak{M} be elementary substructure of \mathfrak{N} . Let $\bar{a} \in N \setminus M$ and $p := tp(\bar{a}|M)$. Then for any formula $\psi(\bar{x}, \bar{y})$ define the predicate $R_{(\psi,p)}(\bar{y})$ on the set M , $\models R_{(\psi,p)}(\bar{a})$ iff $\psi(\bar{x}, \bar{a}) \in tp(\bar{a}|M)$ iff $\mathfrak{N} \models \psi(\bar{a}, \bar{a})$. Denote by $\mathfrak{M}^+ = \langle M; \Sigma^+ \rangle$, where $\Sigma^+ := \{R_{(\psi,p)}(\bar{y}) | p \in S(M), \psi \in \Sigma\}$. It follows from definition that if a pair of models (M, N) is conservative pair (type of any tuple elements from N over M is definable), then the structure \mathfrak{M}^+ is the structure obtained from \mathfrak{M} scolemisation of \mathfrak{M} . We will consider the simple cases when \mathfrak{M}^+ constructed from one 1-type for o-minimal theory from two approaches.

Let \mathfrak{M} be a model of an arbitrary complete theory T of the signature Σ . We say that \mathfrak{M}_p^+ is expansion of \mathfrak{M} by type $p \in S_1(M)$, if $\mathfrak{M}_p^+ = \langle M; \Sigma_p^+ \rangle$, where $\Sigma_p^+ := \{R_{(\psi,p)}(\bar{y}) | \psi \in \Sigma\}$.

We say that \mathfrak{M}_p^+ admits uniformly representation of Σ_p^+ -formulas by Σ -formulas, if for any formula $\phi(\bar{y})$ of Σ_p^+ there exists Σ -formula $K_\phi(\bar{y}, \bar{z})$, there exists $\bar{a} \in N \setminus M$ such that for any $\bar{a} \in M$ the following holds:

$$\mathfrak{M}_p^+ \models \phi(\bar{a}) \Leftrightarrow \models K_\phi(\bar{a}, \bar{a}).$$

Approach of Macpherson-Marker-Steinhorn. In the paper [1] (preprint 1994 Macpherson-Marker-Steinhorn proved weak o-minimality of the expansion of an o-minimal structure by unary convex predicate, such that the predicate is traversed by a uniquely realizable 1-type. Following D. Marker [5], an uniquely realizable 1-type $p \in S_1(M)$ over model is that prime model over model and one realization of this 1-type p contains just this element from the set of realization of the type. An uniquely realizable 1-type has the next property: there is no definable function acting on the set of realizations of this 1-type p . Macpherson-Marker-Steinhorn considered at the same time two structures $\mathfrak{M}^+ = \langle M; \Sigma \cup \{U^1\} \rangle$ and $\mathfrak{N} = \langle N; \Sigma \rangle$, where \mathfrak{N} is a model of an o-minimal theory of the signature Σ and a saturated elementary extension of \mathfrak{M} . They defined a new unary convex predicate U by using an element $\alpha \in N \setminus M$ from the set of realizations of an irrational 1 –type $p \in S_1(M)$ such that for every $a \in M$ the following holds:

$$\mathfrak{M}^+ \models U(a) \Leftrightarrow \mathfrak{N} \models a < \alpha.$$

By induction of construction of formulas $\phi(\bar{y})$ of the signature $\Sigma^+ = \Sigma \cup \{U^1\}$ there is a formula $K_\phi(\bar{y}, \alpha)$ of the signature Σ such that for any $\bar{a} \in M$ the following holds:

for any $n, k, m < \omega$,

$$QV_{p_n}(\alpha_n, \alpha_{n+1}, \dots, \alpha_{n+k}) < QV_{p_n}(\alpha_{n+k+1}, \alpha_{n+k+2}, \dots, \alpha_{n+k+m}),$$

and consequently, all these sets have empty intersection. D. Marker proved that for any set A in o-minimal theory, for any $q, r \in S_1(A)$ if q and r are not weakly orthogonal than there is A -definable monotonic bijection from $q(\mathfrak{N})$ to $r(\mathfrak{N})$. Then for $r \in S_1(M)$, p_i non weakly orthogonal to r , $QV_r(\bar{\alpha}) \cap QV_{p_i}(\bar{\alpha}_i, \alpha_{n+i+1}, \dots, \alpha_{2n+1}) = \emptyset$.

$$\mathfrak{M}^+ \models \phi(\bar{a}) \Leftrightarrow \mathfrak{N} \models K_\phi(\bar{a}, \alpha). \tag{1}$$

The crucial point in this construction was the case $\phi(\bar{y}) = \exists x\psi(x, \bar{y})$. They proposed

$$K_{\exists x\psi(x, \bar{y})}(\bar{y}, \alpha) := \exists z_1 \exists z_2 \exists x (z_1 < \alpha < z_2 \wedge \forall z (z_1 < z < z_2 \rightarrow K_{\psi(x, \bar{y})}(x, \bar{y}, z)).$$

Since the 1-type $p \in S_1(M)$ is uniquely realizable, two convex to right and to left from α $M\alpha - 1$ -formulas have solutions out of $p(\mathfrak{N})$. Thus for any $\bar{a} \in M$, if $\mathfrak{N} \models K_{\exists x\psi(x, \bar{y})}(\bar{a}, \alpha)$, then for some $b_1, b_2 \in M$,

$$\mathfrak{N} \models \exists x (b_1 < \alpha < b_2 \wedge \forall z (b_1 < z < b_2 \rightarrow K_{\psi(x, \bar{y})}(x, \bar{a}, z)).$$

This means that in an elementary submodel of \mathfrak{N} the part of the last formula holds on $\mathfrak{M} \models \exists x \forall z (b_1 < z < b_2 \rightarrow K_{\psi(x, \bar{y})}(x, \bar{a}, z))$. Then there is an element $c \in M$ such that $\mathfrak{M} \models \forall z (b_1 < z < b_2 \rightarrow K_{\psi(x, \bar{y})}(c, \bar{a}, z))$. So, $K_{\psi(x, \bar{y})}(c, \bar{a}, z) \in p$.

Thus, any $\Sigma^+ - M - 1$ -formula $\phi(x, \bar{a})$ has the set of its realizations, $\phi(\mathfrak{M}^+, \bar{a}) = K_\phi(\mathfrak{N}, \bar{a}) \cap M$, being a finite union of convex sets because $K_\phi(\mathfrak{N}, \bar{a})$ is a finite union of intervals and points. The elementary theory of \mathfrak{M}^+ is weakly o-minimal since the number of convex sets is bounded and consequently does not depend on parameters.

Approach of B.S. Baizhanov. For the case when $p \in S_1(M)$ is a non uniquely realizable type, B.S. Baizhanov proposed [2] (1995), on the base of theory of (non)orthogonality of 1-types and its classification made in [4], [5], [6], [8] (Pillay-Steinhorn, Marker, Mayer, Marker-Steinhorn, 1986–1994), to take the constants for $K_{\exists x\psi(x, \bar{y})}$ from an infinite indiscernible sequence $I = \langle \alpha_n \rangle_{n < \omega}$ over M and α_n from $p(\mathfrak{N})$. Taking into consideration that if $K_{\psi(x, \bar{y})}(\mathfrak{N}, \bar{a}, \bar{\alpha}_n) \cap M = \emptyset$, then there is a finite number irrational cuts (1-types over M) such that for any such 1-type $r \in S_1(M)$, $K_{\psi(x, \bar{y})}(\mathfrak{N}, \bar{a}, \bar{\alpha}_n)$ is a subset of

$$QV_r(\bar{\alpha}_n) := \{ \beta \in r(\mathfrak{N}) \mid \text{there exists an } M\bar{\alpha}_n\text{-1-formula } \Theta(x, \bar{\alpha}_n), \text{ such that } \beta \in \Theta(\mathfrak{N}, \bar{\alpha}_n) \subset r(\mathfrak{N}) \}.$$

The idea to use an indiscernible sequence consists from two parts.

B1. On the one hand, if for some $c \in M$, $\mathfrak{N} \models K_{\psi(x, \bar{y})}(c, \bar{a}, \bar{\alpha}_n)$, then for any $\bar{\gamma} = \langle \alpha_{i_0}, \dots, \alpha_{i_n} \rangle$ ($n < i_0 < \dots < i_n$), $\mathfrak{N} \models K_{\psi(x, \bar{y})}(c, \bar{a}, \bar{\gamma})$, because $\bar{\alpha}_n$ and $\bar{\gamma}$ have the same type over M .

B2. On the other hand, to find a sequence I such that for any $r \in S_1(M)$, for any $\bar{\gamma} = \langle \alpha_{i_0}, \dots, \alpha_{i_n} \rangle$ ($n < i_0 < \dots < i_n$), $QV_r(\bar{\alpha}_n) \cap QV_r(\bar{\gamma}) = \emptyset$.

For find the indiscernible sequence I define the properties A1-A3 that follow from the classification of 1-types and theory non orthogonality of 1-types over sets in o-minimal theories.

A1. [5] (Marker 1986). *Let $q, r \in S_1(A)$, and let type $q(x) \cup r(y)$ be non complete (q is non weakly orthogonal to r , Shelah, 1978). Then there is an A -definable monotonic bijection $g: q(\mathfrak{N}) \rightarrow r(\mathfrak{N})$ and consequently, q is irrational if and only if r is irrational;*

q is uniquely realizable if and only if r is uniquely realizable.

Recall that if $q \in S_1(A)$ is irrational then $q(\mathfrak{N})$ is a convex non-definable set without maximal and minimal elements.

A2. *If $q \in S_1(A)$ is irrational, then for any $\bar{\gamma}$, $QV_q(\bar{\gamma}) = V_q(\bar{\gamma})$, here*

$$V_q(\bar{\gamma}) := \{ \beta \in q(\mathfrak{N}) \mid \exists \delta_1, \delta_2 \in q(\mathfrak{N}), \text{ there exists an } A\bar{\gamma}\text{-1-formula } S(x, \bar{\gamma}), \delta_1 < S(\mathfrak{N}, \bar{\gamma}) < \delta_2, \beta \in S(\mathfrak{N}, \bar{\gamma}) \}.$$

Indeed, the *quasi-neighborhood* of \bar{y} in q ($QV_q(\bar{y})$) is the union of $A\bar{y}$ -definable sets, and any such definable set is a subset of $q(\mathfrak{M})$, a convex non-definable set without minimal and maximal elements. The last means that such a definable set is a subset of $V_q(\bar{y})$ (*neighborhood of \bar{y} in q*). This explains equality of two convex sets.

A3. If $q \in S_1(A)$ is irrational and non uniquely realizable, then for any $\bar{y} \in N$, if $QV_q(\bar{y}) \neq \emptyset$ then the 1-types $q(x) \cup \{x < QV_q(\bar{y})\}$ and $q(x) \cup \{QV_q(\bar{y}) < x\}$ are irrational and non uniquely realizable.

By A2 and theorem of compactness there exist $\delta_1, \delta_2 \in q(\mathfrak{M})$ such that $\delta_1 < V_q(\bar{y}) < \delta_2$ and because q is non uniquely realizable i.e. there is A -definable monotonic bijection $f: q(\mathfrak{M}) \rightarrow q(\mathfrak{M})$, $V_q(\mathfrak{M})$ can not have minimal and maximal element. Taking in consideration that for irrational q , $q(\mathfrak{M})$ is a convex non-definable set without maximal and minimal elements, $r_1 := tp(\delta_1 | A\bar{y})$ and $r_2 = tp(\delta_2 | A\bar{y})$ are irrational and since $f(V_q(\bar{y})) = V_q(\bar{y})$, f acts on $r_1(\mathfrak{M})$ and $r_2(\mathfrak{M})$. The last means r_1 and r_2 are non uniquely realizable.

Let $p_n(x) := p(x) \cup (QV_p(\bar{\alpha}_{n-1}) < x)$. Then by A2, A3 p_n is irrational, non uniquely realizable and finitely satisfiable in M since right sides of p and p_n coincide

For any $n, k, m < \omega$,

$$QV_{p_n}(\alpha_n, \alpha_{n+1}, \dots, \alpha_{n+k}) < QV_{p_n}(\alpha_{n+k+1}, \alpha_{n+k+2}, \dots, \alpha_{n+k+m}), \quad (2)$$

and consequently, all these sets have empty intersection.

The proof of (2) is done by induction on m . Assume (2) for m . Denote by $r_1(x) = p_n \cup (x < QV_{p_n}(x)(\alpha_{n+k+1}, \dots, \alpha_{n+k+m}))$ and $r_2(y) = p_n(y) \cup (QV_{p_n}(\alpha_{n+k+1}, \dots, \alpha_{n+k+m}) < y)$, Suppose that

$QV_{p_n}(\alpha_n, \alpha_{n+1}, \dots, \alpha_{n+k}) \cap QV_{p_n}(\alpha_{n+k+1}, \dots, \alpha_{n+k+m}, \alpha_{n+k+m+1}) \neq \emptyset$. Since the first set does not change, there exists $M\bar{\alpha}_n \alpha_{n+k+1} \dots \alpha_{n+k+m+1}$ -formula $L(x, \alpha_{n+k+m+1})$ such $L(\mathfrak{M}, \alpha_{n+k+m+1}) \subset QV_{p_n}(\alpha_n, \alpha_{n+1}, \dots, \alpha_{n+k}) \subset r(\mathfrak{M})$. Let β be end point of one of interval of formula L , then because p_n is non uniquely realizable then $\beta \in QV_{p_n}(\alpha_n, \alpha_{n+1}, \dots, \alpha_{n+k}) \subset r(\mathfrak{M})$. Since $\beta \models r_1$ and $\alpha_{n+k+m+1} \models r_2$ by A1 there exists $M\bar{\alpha}_n \alpha_{n+k+1} \dots, \alpha_{n+k+m}$ -definable monotonic function $f: r_2(\mathfrak{M}) \rightarrow r_1(\mathfrak{M})$ such that $f(\alpha_{n+k+m+1}) = \beta$. On other hand $\beta \in QV_{p_n}(\alpha_n, \alpha_{n+1}, \dots, \alpha_{n+k})$ and consequently, there is $M\bar{\alpha}_{n+k} - 1$ -formula $H(x)$ such that $\beta \in H(\mathfrak{M}) \subset QV_{p_n}(\alpha_n, \alpha_{n+1}, \dots, \alpha_{n+k}) \subset r_1(\mathfrak{M})$. Then $\alpha_{n+k+m+1}$ belongs to $M\bar{\alpha}_{n+k+m}$ -definable set $f^{-1}(H(\mathfrak{M})) \subset r_2(\mathfrak{M})$. This means $\alpha_{n+k+m+1} \in QV_p(\bar{\alpha}_{n+k+m})$. Contradiction.

It follow from (2) and A1 that for any $r \in S_1(M)$, if for any $i < n$, $p_i \perp^w r$ and $p_n \not\perp^w r$ then $QV_r(\bar{\alpha}) \cap QV_r(\bar{\alpha}_i, \alpha_{n+i+1}, \dots, \alpha_{2n+1}) = \emptyset$.

$$QV_r(\bar{\alpha}) \cap QV_r(\bar{\alpha}_i, \alpha_{n+i+1}, \dots, \alpha_{2n+1}) = \emptyset \quad (3)$$

Suppose for the formula $\psi(x, \bar{y})$ of signature Σ^+ corresponding formula of signature Σ is $K_{\psi(x, \bar{y})}(x, \bar{y}, \bar{\alpha}_n)$. Thus for any formula $K_\psi(x, \bar{\alpha}, \bar{\alpha}_n)$ to have the solution in M it is sufficient to write the formula

$$K_{\exists x \psi(x, \bar{y})}(\bar{y}, \bar{\alpha}_{2n+1}) := \exists x (K_\psi(x, \bar{y}, \bar{\alpha}_n) \wedge \bigwedge_{i \leq n} K_\psi(x, \bar{y}, \bar{\alpha}_{n-i}, \alpha_{(n-i)+n+1}, \alpha_{(n-i)+n+2}, \dots, \alpha_{2n+1})).$$

B.S. Baizhanov in 1996 obtained a classification of 1-types over a subset of a model of weakly o-minimal theory and solved the problem of expanding a model of weakly-o-minimal theory by a unary convex predicate in the preprint "Classifications of 1-types in weakly o-minimal theories and its applications" and submitted in the JSL, that revised version published in [9](2001). Ye.Baisalov and B. Poizat [10] (preprint 1996) in the paper on "beautiful" pairs of models of o-minimal theories proved the elimination of quantify $\exists x \in M$. It is difficult to say that the approach in [10] is alternative to approach elaborated in [2], because they used the same principles B1-B2 from [2].

We say that \mathfrak{M}^+ the expansion by all externally definable subsets admits quantifier elimination, if for any formula $\phi(\bar{y})$ of Σ^+ there exists Σ -formula $K_\phi(\bar{y}, \bar{z})$, there exists $\bar{\alpha} \in N \setminus M$ such that for any $\bar{a} \in M$ the following holds:

$$\mathfrak{M}^+ \models \phi(\bar{a}) \Leftrightarrow K_\phi(\bar{a}, \bar{\alpha}).$$

Approach of Shelah. In his paper, S. Shelah [11] (2004) considered a model of NIP theory and proved that the expansion by all externally definable subsets admits quantifier elimination and thereby is NIP. The key problem here is eliminating quantifier "there exists in the submodel". In his proof in the way of contradiction Shelah used an indiscernible sequence $\langle \bar{b}_n : n < \omega \rangle$ in order to show that if eliminating quantifier "there exists x in the submodel" $\varphi(x, \bar{a})$ fails, then $\varphi(\alpha, \bar{b}_n)$ holds iff n is even, for some α , which implies the independence property, for a contradiction.

V.V. Verbovskiy [12] (preprint 2005) found a somewhat simplified account of Shelah's proof, namely by using noting of a finitely realizable type. A. Pillay [13] (preprint 2006) gave two re-proofs of Shelah's theorem, the first going through quantifier-free heirs of quantifier-free types and the second through quantifier-free coheirs of quantifier-free types.

The analysis of approaches shows that the using the theory of orthogonality we can control the set of realizations of one-types. If we consider the complete theory satisfies A2, it gives the possibility to construct the indiscernible sequence satisfied the condition B2. Notice that the of indiscernible sequence constructed by mathematical induction satisfies the condition of finite realizability of an one-type of new element over model and beginning of sequence. The generalization of the approach for o-minimal model in case of non uniquely realizable one-type by introduction of generalization of the notions of (quasi)-neighborhood and almost (non)-orthogonality of two types gives the possibility to formulate the next

Theorem 1 Let T be a complete NSOP theory such that for any set A the following holds:

- 1) For any $p \in S_1(A)$, for any \bar{y} , $QV_p(\bar{y}) = V_p(\bar{y})$
- 2) For any $p, q \in S_1(A)$ the following holds. If $p \not\perp^\alpha q$, then $q \not\perp^\alpha p$.

Then for model of the theory T the expansion of this model by one-type admits uniformly representation of Σ_p^+ -formulas by Σ -formulas.

The work is financially supported by the Ministry of Education and Science of the Republic of Kazakhstan (grant AP05134992)

Б.С. Байжанов¹, С.С. Байжанов²

¹РГП «Институт математики и математического моделирования» КН МОН РК, Алматы, Казахстан,

²КазНУ им. Аль-Фараби, Алматы, Казахстан

НЕКОТОРЫЕ ВОПРОСЫ О ВНЕШНЕЙ ОПРЕДЕЛИМОСТИ

Аннотация. В статье рассматриваются различные подходы к концепции внешней определимости, разработанные в о-минимальных теориях. Пример о-минимальных теорий показывает, как внешние константы помогают определить существование решения в модели формулы с внешними константами. Сформулированы основные понятия, с помощью которых доказывается внешняя определимость. Дается краткий обзор результатов для зависимых теорий. В заключение сформулированы достаточные условия, так что теория обладает свойством внешней определимости. Дано краткое объяснение изложенной теоремы.

Ключевые слова: внешне определяемая, окрестность кортежа множества в типе, неортогональность двух типов

Б.С. Байжанов¹, С.С. Байжанов²

¹РМ «Математика және математикалық үлгілеу институты» РМ, ҚР ҚМ, Алматы, Қазақстан,

²Әл-Фараби атындағы ҚазҰУ, Алматы, Қазақстан

СЫРТҚЫ ЕРЕЖЕЛЕРГЕ ҚАТЫСТЫ САУАЛДАР

Аннотация. Мақалада о-минималды теорияларда жасалған сыртқы анықтама тұжырымдамасына әртүрлі көзқарастар қарастырылған. О-минималды теориялардың мысалы сыртқы тұрақтылар формула моделінде ерітіндінің болуын анықтауға көмектесетінін көрсетеді. Негізгі ұғымдар сыртқы анықталуы дәлелденген, тұжырымдалған. Тәуелді теориялардың нәтижелеріне қысқаша шолу келтірілген.

Қорытындылай келе, теория сыртқы анықталу қасиетіне ие болатындай жеткілікті жағдайлар жасалған. Көрсетілген теоремаға қысқаша түсініктеме беріледі.

Түйін сөздер: сыртқы анықталған, типтегі жиынның көршілес, екі түрге жатпайтындығы.

Informaiton about authors:

Baizhanov Bektur Sembiuly - Institute of mathematics and mathematical modelling, deputy director, corresponding member of NAS RK, <https://orcid.org/0000-0002-3743-7404>, baizhanov@hotmail.com

Baizhanov Sayan Samatovich – PhD student, <https://orcid.org/0000-0002-7482-7145>, sayan-5225@mail.ru

REFERENCES

- [1] D. Macpherson, D. Marker, Ch. Steinhorn, Weakly o-minimal structures and real closed fields // *Translations of the American Mathematical Society*, 352 (2000), 5435–5483.
- [2] B.S. Baizhanov, Extensions of o-minimal structures by convex unary predicates // In the book: "Research in the theory of algebraic systems" (editor T.A. Nurmagambetov), Karaganda State University, 1995, 6–24. (in Russian)
- [3] A. Pillay, C. Steinhorn, Definable sets in ordered structures // *Bulletin AMS*, 11 (1984), 159–162.
- [4] A. Pillay, C. Steinhorn, Definable sets in ordered structures I // *Transactions of American Mathematical Society*, 295 (1986), 565–592.
- [5] D. Marker, Omitting types in o-minimal theories // *The Journal of Symbolic Logic*, 51 (1986), 63–74.
- [6] L. Mayer, Vaught's conjecture for o-minimal theories // *The Journal of Symbolic Logic*, 53 (1988), 146–159.
- [7] A. Pillay, Ch. Steinhorn, Definable sets in ordered structures III // *Transactions of the American Mathematical Society* 300 (1988), 469–476.
- [8] D.Marker, Ch. Steinhorn, Definable types in o-minimal theories // *The Journal of Symbolic Logic*, 59 (1994), 185–198.
- [9] B.S. Baizhanov, Expansion of a model of a weakly o-minimal theory by a family of unary predicates // *The Journal of Symbolic Logic*, 66 (2001), 1382–1414.
- [10] Y. Baisalov, B. Poizat, Pairs of O-minimal structures // *Journal of Symbolic Logic*, 63:2 (1998), 570–578.
- [11] S. Shelah, Dependent first order theories // *Continued, Israel Journal of Mathematics* 173: 1. (2009) <https://doi.org/10.1007/s11856-009-0082-1>
- [12] V. Verbovskiy, Dependent theories. The Shelah's theorem // *Proceeding of International conference "Contemporary problems of mathematics, informatics and control"*, dedicated to 60th anniversary of M.B. Aidarkhanov, October 2-3, 2008, Almaty, 439–441
- [13] A. Pillay, On externally definable sets and a theorem of Shelah // *Algebra, logic, set theory, Stud. Log.(Lond.)*, (2007) pdfs.semanticscholar.org

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

www.nauka-nanrk.kz

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы *М. С. Ахметова, Т. А. Апендиев, Д. С. Аленов*
Верстка на компьютере *А. М. Кульгинбаевой*

Подписано в печать 12.12.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
9,6 п.л. Тираж 300. Заказ 6.