

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Әль-фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
Әль-Фараби атындағы
Қазақ ұлттық университетінің

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-farabi kazakh
national university

SERIES
PHYSICO-MATHEMATICAL

6(328)

NOVEMBER – DECEMBER 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Бас редакторы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі **Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев У.У. проф. корр.-мүшесі(Қазақстан)
Жүсіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Д. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қырғыстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорус)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик(Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы»РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18, <http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жангаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф., чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

Editor in chief
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d :

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskiy I.N. prof., academician(Ukraine)
Kovalev A.M. prof., academician(Ukraine)
Mikhalevich A.A. prof., academician(Belarus)
Pashayev A. prof., academician(Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician(Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.69>

Volume 6, Number 328 (2019), 14 – 24

UDC 517.958:536.2; 517.968.7; 519.62/.64

MRNTI 27.35.45; 27.33.19; 27.41.19

A.T.Assanova^{1,2}, E.A.Bakirova^{1,2,3}, Zh.M.Kadirbayeva^{1,2,4}

¹Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan;

²Institute of Information and Computational Technologies, Almaty, Kazakhstan;

³Kazakh National Women's Teacher Training University, Almaty, Kazakhstan;

⁴International Information Technology University, Almaty, Kazakhstan

assanova@math.kz, bakirova1974@mail.ru, zh.kadirbayeva@gmail.com

**NUMERICALLY APPROXIMATE METHOD FOR SOLVING
OF A CONTROL PROBLEM FOR INTEGRO-DIFFERENTIAL
EQUATIONS OF PARABOLIC TYPE**

Abstract. A linear boundary value problem with a parameter for integro-differential equations of parabolic type is investigated. Using the spatial variable discretization, the considering problem is approximated by a linear boundary value problem with a parameter for a system of ordinary integro-differential equations. The parameterization method is used for solving the obtained problem. The approximating problem is reduced to an equivalent problem consisting of a special Cauchy problem for the system of Fredholm integro-differential equations, boundary conditions, and continuity conditions of the solution at the partition points. The solution of the Cauchy problem for the system of ordinary differential equations with parameters is constructed using the fundamental matrix of the differential equation. The system of a linear algebraic equations with respect to the parameters are composed by substituting the values of the corresponding points in the boundary condition and the continuity conditions. Numerical method for solving of the problem is suggested, which based on the solving of the constructed system and method of Runge-Kutta 4-th order for solving of the Cauchy problem on the subintervals.

Key words: partial integro-differential equations of parabolic type, problem with parameter, approximation, numerically approximate method, algorithm.

Funding: The work was supported by grants (№№ AP05132486, AP05131220, AP05132455) of Ministry of Education and Science of the Republic of Kazakhstan (2018-2020).

Control problems, which are also called boundary value problems with parameters and the problem of parameter identification for a system of ordinary differential and integro-differential equations with parameters, have been actively investigated in recent decades. Models describing reaction-diffusion processes lead to control problems for integro-differential equations of parabolic type [1-17]. Questions of existence, uniqueness and stability of solving problems with parameters are very important for development of numerical methods of identification of parameters of the mathematical models described by integro-differential equations of parabolic type [1-17].

In the present paper, linear problem with a parameter for an integro-differential equation of parabolic type is investigated. By discretizing a spatial variable, the considering problem is approximated by a two-point boundary value problem with parameters for a system of Fredholm integro-differential equations with a degenerate kernel. By introducing additional parameters [18-23] as the values of the desired solution at some points of the interval $[0, T]$, where the problem is considered, the obtained problem is reduced to the equivalent problem consisting of a special Cauchy problem for the system of Fredholm integro-differential equations, boundary conditions, and continuity conditions for the solution at the points of partition. Using the integral equation, that equivalent to the special Cauchy problem for the system of Fredholm integro-differential equation and the property of the degeneracy of kernel of the integral term,

we obtained a representation of the solution of the special Cauchy problem using the entered parameters at the assumption of invertibility of a some matrix. Based on this representation, a system of algebraic equations with respect to the parameters is constructed from the boundary conditions and the continuity conditions of the solution. We offer algorithm for solving the linear boundary value problem for the equation with degenerate kernel, and its numerical implementation.

We consider a linear boundary value problem with a parameter for an integro-differential equation of parabolic type

$$\frac{\partial u}{\partial t} = a(x, t) \frac{\partial^2 u}{\partial x^2} + c(x, t)u + b(x, t)\mu(x) + \varphi(x, t) \int_0^T \psi(x, s)u(x, s)ds + f(x, t), \quad (x, t) \in \Omega = (0, \omega) \times (0, T), \tag{1}$$

$$u(x, 0) = 0, \quad x \in [0, \omega], \tag{2}$$

$$u(x, T) = 0, \quad x \in [0, \omega], \tag{3}$$

$$u(0, t) = \tilde{\psi}_1(t), \quad u(\omega, t) = \tilde{\psi}_2(t), \quad t \in [0, T], \tag{4}$$

where $u(x, t)$ is sought function, $\mu(x)$ is unknown functional parameter, functions $a(x, t) \geq a_0 > 0$, $c(x, t) \leq 0$, $b(x, t)$, $\varphi(x, t)$, $\psi(x, t)$, $f(x, t)$ are continuous in t and Holder continuous in x on Ω ; functions $\tilde{\psi}_1(t)$, $\tilde{\psi}_2(t)$ are continuous on $[0, T]$. It is assumed that the boundary functions are sufficiently smooth and satisfy the matching conditions.

The solution of the boundary problem (1)-(4) is a pair of functions $(u^*(x, t), \mu^*(x))$, where function $u^*(x, t)$ is continuous on Ω , that has continuous partial derivatives with respect to x of first order, with respect to t of second order, satisfies the integro-differential equation (1) at $\mu(x) = \mu^*(x)$, $x \in [0, \omega]$, and boundary conditions (2)-(4).

In view of condition (2)-(4), from (1) we obtain two groups of equations for determining $\mu(0)$ and $\mu(\omega)$:

$$\begin{aligned} b(0,0)\mu(0) &= \dot{\tilde{\psi}}_1(0) - \varphi(0,0) \int_0^T \psi(0,s)\tilde{\psi}_1(s)ds - f(0,0), \\ b(\omega,0)\mu(\omega) &= \dot{\tilde{\psi}}_2(0) - \varphi(\omega,0) \int_0^T \psi(\omega,s)\tilde{\psi}_2(s)ds - f(\omega,0), \\ b(0,T)\mu(0) &= \dot{\tilde{\psi}}_1(T) - \varphi(0,T) \int_0^T \psi(0,s)\tilde{\psi}_1(s)ds - f(0,T), \\ b(\omega,T)\mu(\omega) &= \dot{\tilde{\psi}}_2(T) - \varphi(\omega,T) \int_0^T \psi(\omega,s)\tilde{\psi}_2(s)ds - f(\omega,T). \end{aligned}$$

These relations also are the matching conditions with respect to initial data.

We take $\forall h > 0$ and produce a discretization by x : $x_i = ih, i = \overline{0, P}, Ph = \omega$.

We introduce the notations $u_i(t) = u(ih, t)$, $\mu_i = \mu(ih)$, $a_i(t) = a(ih, t)$, $c_i(t) = c(ih, t)$, $b_i(t) = b(ih, t)$, $\varphi_i(t) = \varphi(ih, t)$, $\psi_i(t) = \psi(ih, t)$, $f_i(t) = f(ih, t)$, $i = \overline{0, P}$.

Problem (1) - (4) is replaced by the following linear boundary value problem with a parameter for an integro-differential equation

$$\frac{du_i}{dt} = a_i(t) \frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} + c_i(t)u_i + b_i(t)\mu_i + \varphi_i(t) \int_0^T \psi_i(s)u_i(s)ds + f_i(t), \quad i = \overline{1, P-1}, \tag{5}$$

$$u_i(0) = 0, \quad i = \overline{0, P}, \tag{6}$$

$$u_i(T) = 0, \quad i = \overline{0, P}, \tag{7}$$

$$u_0(t) = \tilde{\psi}_1(t), \quad u_P(t) = \tilde{\psi}_2(t), \quad t \in [0, T]. \tag{8}$$

The functions $u_0(t)$, $u_p(t)$, and parameters μ_0, μ_p are known.
 Problem (5)-(8) will be rewritten in vector-matrix form

$$\frac{du}{dt} = A(t)u + B(t)\mu + \Phi(t) \int_0^T \Psi(s)u(s)ds + F(t), \quad u, \mu \in R^{P-1}, \quad t \in (0, T), \quad (9)$$

$$u(0) = 0, \quad (10)$$

$$u(T) = 0, \quad (11)$$

where $u(t) = (u_1(t), u_2(t), \dots, u_{p-1}(t))$, $\mu = (\mu_1, \mu_2, \dots, \mu_{p-1})$ -unknown function and parameter,

$$A(t) = \begin{pmatrix} -\frac{2a_1(t)}{h^2} + c_1(t) & \frac{a_1(t)}{h^2} & 0 & \dots & 0 \\ \frac{a_2(t)}{h^2} & -\frac{2a_2(t)}{h^2} + c_2(t) & \frac{a_2(t)}{h^2} & \dots & 0 \\ 0 & \frac{a_3(t)}{h^2} & -\frac{2a_3(t)}{h^2} + c_3(t) & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & -\frac{2a_{p-1}(t)}{h^2} + c_{p-1}(t) \end{pmatrix},$$

$$B(t) = \text{diag}\{b_1(t), b_2(t), \dots, b_{p-1}(t)\},$$

$$\Phi(t) = \text{diag}\{\varphi_1(t), \varphi_2(t), \dots, \varphi_{p-1}(t)\},$$

$$\Psi(s) = \text{diag}\{\psi_1(s), \psi_2(s), \dots, \psi_{p-1}(s)\},$$

$$F(t) = \left(\frac{a_1(t)}{h^2} \tilde{\psi}_1(t) + f_1(t), f_2(t), \dots, \frac{a_{p-1}(t)}{h^2} \tilde{\psi}_2(t) + f_{p-1}(t) \right)'$$

Here $(P - 1) \times (P - 1)$ -matrices $A(t)$, $B(t)$, $\Phi(t)$, $\Psi(s)$ and $(P - 1)$ -vector $F(t)$ are continuous on $[0, T]$.

The solution to problem (9) - (11) is a pair $(u^*(t), \mu^*)$, where $u^*(t)$ is continuous on $[0, T]$ and continuously differentiable on $(0, T)$ a function $u^*(t)$ satisfies the integro-differential equation (9) at $\mu = \mu^*$ and conditions (10), (11).

To solve the problem with parameter (9)-(11), the approach developed in [24-26] is used, based on the algorithms of the parameterization method and numerical methods for solving Cauchy problems.

Scheme of the method. Points $0 = t_0 < t_1 < \dots < t_{N-1} < t_N = T$ are taken and the interval $[0, T]$ is divided into N subintervals: $[0, T] = \bigcup_{r=1}^N [t_{r-1}, t_r)$, which is denoted by Δ_N [20]. The restriction of the function $u(t)$ to the r -th interval $[t_{r-1}, t_r)$ is denoted by $x_r(t)$, i.e. $u_r(t) = u(t)$ for $t \in [t_{r-1}, t_r)$, $r = \overline{1, N}$.

Let $C([0, T], R^{P-1})$ be the space of continuous on $[0, T]$ functions $u: [0, T] \rightarrow R^{P-1}$ with norm $\|u\|_1 = \max_{t \in [0, T]} \|u(t)\|$; $C([0, T], \Delta_N, R^{(P-1)N})$ - the space of systems of functions $u[t] = (u_1(t), u_2(t), \dots, u_N(t))$, where $u_r: [t_{r-1}, t_r) \rightarrow R^{P-1}$ are continuous on $[t_{r-1}, t_r)$ and have finite left-sided limits $\lim_{t \rightarrow t_{r-1}^+} u_r(t)$ for all $r = \overline{1, N}$, with norm $\|u[\cdot]\|_2 = \max_{r=\overline{1, N}} \sup_{t \in [t_{r-1}, t_r)} \|u_r(t)\|$.

We introduce additional parameters $\lambda_r = u_r(t_{r-1})$, $r = \overline{2, N}$, $\lambda_1 = \mu$. Making the substitution $u_1(t) = z_1(t)$, $u_r(t) = z_r(t) + \lambda_r$ on every r -th interval $[t_{r-1}, t_r)$, $r = \overline{2, N}$, we obtain multipoint boundary value problem with parameters

$$\frac{dz_1}{dt} = A(t)z_1 + B(t)\lambda_1 + \Phi(t) \sum_{j=1}^N \int_{t_{j-1}}^{t_j} \Psi(s)z_j(s)ds + \Phi(t) \sum_{j=2}^N \int_{t_{j-1}}^{t_j} \Psi(s)\lambda_j ds + F(t), \quad t \in [t_0, t_1), \quad (12)$$

$$z_1(t_0) = 0, \quad (13)$$

$$\frac{dz_r}{dt} = A(t)(z_r + \lambda_r) + B(t)\lambda_1 + \Phi(t) \sum_{j=1}^N \int_{t_{j-1}}^{t_j} \Psi(s)z_j(s)ds + \Phi(t) \sum_{j=2}^N \int_{t_{j-1}}^{t_j} \Psi(s)\lambda_j ds + F(t), \quad t \in [t_{r-1}, t_r), \tag{14}$$

$$z_r(t_{r-1}) = 0, \quad r = \overline{2, N}, \tag{15}$$

$$\lambda_N + \lim_{t \rightarrow T-0} z_N(t) = 0, \tag{16}$$

$$\lim_{t \rightarrow t_1-0} z_1(t) = \lambda_2, \tag{17}$$

$$\lambda_s + \lim_{t \rightarrow t_s-0} z_s(t) = \lambda_{s+1}, \quad s = \overline{2, N-1}. \tag{18}$$

The solution of the problem with parameters (12)-(18) is a pair $(z^*[t], \lambda^*)$ where the function $z^*[t] = (z_1^*(t), z_2^*(t), \dots, z_N^*(t)) \in C([0, T], \Delta_N, R^{(P-1)N})$ with continuously differentiable components $z_r^*(t)$ on $[t_{r-1}, t_r)$ and $\lambda^* = (\lambda_1^*, \lambda_2^*, \dots, \lambda_N^*) \in R^{(P-1)N}$, satisfies the integro-differential equation with parameters (12), (14), initial conditions (13), (15), relations (16)-(18) at $\lambda_j = \lambda_j^*, j = \overline{2, N}$.

If the pair $(u^*(t), \mu^*)$ is a solution of problem (9)-(11), then the pair $(z^*[t], \lambda^*)$ with elements $z^*[t] = (z_1^*(t), z_2^*(t), \dots, z_N^*(t)) \in C([0, T], \Delta_N, R^{(P-1)N})$, $\lambda^* = (\lambda_1^*, \lambda_2^*, \dots, \lambda_N^*) \in R^{(P-1)N}$, where $\lambda_1^* = \mu^* \in R^{P-1}$, $z_1^*(t) = u_1^*(t)$, $t \in [t_0, t_1)$, $\lambda_r^* = u_r^*(t_{r-1})$, $z_r^*(t) = u_r^*(t) + u_r^*(t_{r-1})$, $t \in [t_{r-1}, t_r)$, $r = \overline{2, N}$, is the solution of problem (12)-(18). Conversely, if a pair $(\tilde{z}[t], \tilde{\lambda})$ with elements $\tilde{z}[t] = (\tilde{z}_1(t), \tilde{z}_2(t), \dots, \tilde{z}_N(t)) \in C([0, T], \Delta_N, R^{(P-1)N})$, $\tilde{\lambda} = (\tilde{\lambda}_1, \tilde{\lambda}_2, \dots, \tilde{\lambda}_N) \in R^{(P-1)N}$, is a solution of (12)-(18), then the pair $(\tilde{u}(t), \tilde{\mu})$ defined by the equalities $\tilde{u}(t) = \tilde{z}_1(t)$, $t \in [t_0, t_1)$, $\tilde{u}(t) = \tilde{z}_r(t) + \tilde{\lambda}_r$, $t \in [t_{r-1}, t_r)$, $r = \overline{2, N}$, $\tilde{u}(T) = \lim_{t \rightarrow T-0} \tilde{z}_N(t) + \tilde{\lambda}_N$ and $\tilde{\mu} = \tilde{\lambda}_1$, will be the solution of the original boundary value problem with parameter (9) - (11).

Using the fundamental matrix $X_r(t)$ of the differential equation $\frac{dx}{dt} = A(t)x, t \in [t_{r-1}, t_r)$, $r = \overline{1, N}$, we reduce the solution of a special Cauchy problem for an integro-differential equation with parameters (12)-(15) to an equivalent system of integral equation

$$z_1(t) = X_1(t) \int_{t_0}^t X_1^{-1}(\tau)\Phi(\tau) \left\{ \sum_{j=1}^N \int_{t_{j-1}}^{t_j} \Psi(s)z_j(s)ds + \sum_{j=2}^N \int_{t_{j-1}}^{t_j} \Psi(s)\lambda_j ds \right\} d\tau + X_1(t) \int_{t_0}^t X_1^{-1}(\tau)B(\tau)d\tau \lambda_1 + X_1(t) \int_{t_0}^t X_1^{-1}(\tau)F(\tau)d\tau, \quad t \in [t_0, t_1), \tag{19}$$

$$z_r(t) = X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau)\Phi(\tau) \left\{ \sum_{j=1}^N \int_{t_{j-1}}^{t_j} \Psi(s)z_j(s)ds + \sum_{j=2}^N \int_{t_{j-1}}^{t_j} \Psi(s)\lambda_j ds \right\} d\tau + X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau)A(\tau)d\tau \lambda_r + X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau)B(\tau)d\tau \lambda_1 + X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau)F(\tau)d\tau, \quad t \in [t_{r-1}, t_r), \quad r = \overline{2, N}. \tag{20}$$

Let $\xi = \sum_{j=1}^N \int_{t_{j-1}}^{t_j} \Psi(s)z_j(s)ds$ and rewrite the system of integral equations (19), (20) in the form

$$z_1(t) = X_1(t) \int_{t_0}^t X_1^{-1}(\tau)\Phi(\tau) \left\{ \xi + \sum_{j=2}^N \int_{t_{j-1}}^{t_j} \Psi(s)\lambda_j ds \right\} d\tau + X_1(t) \int_{t_0}^t X_1^{-1}(\tau)B(\tau)d\tau \lambda_1 + X_1(t) \int_{t_0}^t X_1^{-1}(\tau)F(\tau)d\tau, \quad t \in [t_0, t_1), \tag{21}$$

$$\begin{aligned}
 z_r(t) = & X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau) \Phi(\tau) \left\{ \xi + \sum_{j=2}^N \int_{t_{j-1}}^{t_j} \Psi(s) \lambda_j ds \right\} d\tau + \\
 & + X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau) A(\tau) d\tau \lambda_r + X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau) B(\tau) d\tau \lambda_1 + \\
 & + X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau) F(\tau) d\tau, t \in [t_{r-1}, t_r), r = \overline{2, N}.
 \end{aligned} \tag{22}$$

Multiplying both parts of (21), (22) by $\Psi(t)$, integrating on $[t_{r-1}, t_r]$, and summing by r , we obtain a system of linear algebraic equations with respect to $\xi \in R^{P-1}$

$$\xi = G(\Delta_N) \xi + \sum_{r=1}^N V_r(\Delta_N) \lambda_r + g(F, \Delta_N), \tag{23}$$

with $(P-1) \times (P-1)$ -matrices

$$\begin{aligned}
 G(\Delta_N) &= \sum_{r=1}^N \int_{t_{r-1}}^{t_r} \Psi(\tau) X_r(\tau) \int_{t_{r-1}}^{\tau} X_r^{-1}(s) \Phi(s) ds d\tau, \\
 V_1(\Delta_N) &= \sum_{r=1}^N \int_{t_{r-1}}^{t_r} \Psi(\tau) X_r(\tau) \int_{t_{r-1}}^{\tau} X_r^{-1}(s) B(s) ds d\tau, \\
 V_r(\Delta_N) &= \int_{t_{r-1}}^{t_r} \Psi(\tau) X_r(\tau) \int_{t_{r-1}}^{\tau} X_r^{-1}(s) A(s) ds d\tau + \\
 &+ \sum_{r=1}^N \sum_{j=1}^N \int_{t_{r-1}}^{t_r} \Psi(\tau) X_r(\tau) \int_{t_{r-1}}^{\tau} X_r^{-1}(\tau_1) \Phi(\tau_1) d\tau_1 d\tau \int_{t_{j-1}}^{t_j} \Psi(s) ds, r = \overline{2, N}.
 \end{aligned}$$

and $(P-1)$ -vectors

$$g(F, \Delta_N) = \sum_{r=1}^N \int_{t_{r-1}}^{t_r} \Psi(\tau) X_r(\tau) \int_{t_{r-1}}^{\tau} X_r^{-1}(s) F(s) ds d\tau.$$

We write the system (23) in the form

$$[I - G(\Delta_N)] \xi = \sum_{r=1}^N V_r(\Delta_N) \lambda_r + g(F, \Delta_N), \tag{24}$$

where I is the identity matrix of $(P-1)$ dimension.

The special Cauchy problem (12) - (15) is equivalent to the system of integral equations (19) - (20). This system, due to the degeneracy of the kernel, will be equivalent to the system of algebraic equations (23) with respect to $\xi \in R^{P-1}$. The unique solvability of the special Cauchy problem was investigated in [19, 20]. It has been established that with a sufficiently small step $h > 0: Nh = T$ partitioning a segment $[0, T]$ the special Cauchy problem will be unique solvable.

Let the matrix $I - G(\Delta_N)$ be invertible, i.e. exists $[I - G(\Delta_N)]^{-1}$. Then, according to (24), the vector $\xi \in R^{P-1}$ is determined by the equality

$$\xi = [I - G(\Delta_N)]^{-1} \sum_{r=1}^N V_r(\Delta_N) \lambda_r + [I - G(\Delta_N)]^{-1} g(F, \Delta_N). \tag{25}$$

In (21), (22), instead of ξ substituting the right-hand side of (25), we obtain the representation of the function $z_r(t)$ in terms of λ_r , $r = \overline{1, N+1}$:

$$\begin{aligned}
 z_1(t) = & \sum_{j=2}^N \left\{ X_1(t) \int_{t_0}^t X_1^{-1}(\tau) \Phi(\tau) d\tau \left[[I - G(\Delta_N)]^{-1} V_j(\Delta_N) + \int_{t_{j-1}}^{t_j} \Psi(s) ds \right] \right\} \lambda_j + \\
 & + X_1(t) \int_{t_0}^t X_1^{-1}(\tau) [\Phi(\tau) [I - G(\Delta_N)]^{-1} V_1(\Delta_N) + B(\tau)] d\tau \lambda_1 + \\
 & + X_1(t) \int_{t_0}^t X_1^{-1}(\tau) [\Phi(\tau) [I - G(\Delta_N)]^{-1} g(F, \Delta_N) + F(\tau)] d\tau, t \in [t_0, t_1), \tag{26}
 \end{aligned}$$

$$\begin{aligned}
 z_r(t) = & \sum_{j=2}^N \left\{ X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau) \Phi(\tau) d\tau \left[[I - G(\Delta_N)]^{-1} V_j(\Delta_N) + \int_{t_{j-1}}^{t_j} \Psi(s) ds \right] \right\} \lambda_j + \\
 & + X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau) A(\tau) d\tau \lambda_r + X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau) [\Phi(\tau) [I - G(\Delta_N)]^{-1} V_1(\Delta_N) + B(\tau)] d\tau \lambda_1 + \\
 & + X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau) [\Phi(\tau) [I - G(\Delta_N)]^{-1} g(F, \Delta_N) + F(\tau)] d\tau, t \in [t_{r-1}, t_r), r = \overline{2, N}. \tag{27}
 \end{aligned}$$

We introduce the notations

$$\begin{aligned}
 D_{r,j}(\Delta_N) = & X_r(t_r) \int_{t_{r-1}}^{t_r} X_r^{-1}(\tau) \Phi(\tau) d\tau \left[[I - G(\Delta_N)]^{-1} V_j(\Delta_N) + \int_{t_{j-1}}^{t_j} \Psi(s) ds \right], r \neq j, \quad r, j = \overline{2, N}, \\
 D_{r,r}(\Delta_N) = & X_r(t_r) \int_{t_{r-1}}^{t_r} X_r^{-1}(\tau) \Phi(\tau) d\tau \left[[I - G(\Delta_N)]^{-1} V_r(\Delta_N) + \int_{t_{j-1}}^{t_j} \Psi(s) ds \right] \\
 & + X_r(t_r) \int_{t_{r-1}}^{t_r} X_r^{-1}(\tau) A(\tau) d\tau, r = \overline{2, N}, \\
 D_{r,1}(\Delta_N) = & X_r(t_r) \int_{t_{r-1}}^{t_r} X_r^{-1}(\tau) \Phi(\tau) d\tau [I - G(\Delta_N)]^{-1} V_1(\Delta_N) + X_r(t_r) \int_{t_{r-1}}^{t_r} X_r^{-1}(\tau) B(\tau) d\tau, r = \overline{1, N}, \\
 F_r(\Delta_N) = & X_r(t_r) \int_{t_{r-1}}^{t_r} X_r^{-1}(\tau) \Phi(\tau) [I - G(\Delta_N)]^{-1} g(F, \Delta_N) d\tau + X_r(t_r) \int_{t_{r-1}}^{t_r} X_r^{-1}(\tau) F(\tau) d\tau, r = \overline{1, N}.
 \end{aligned}$$

Then from (26), (27) we obtain

$$\lim_{t \rightarrow t_r-0} z_r(t) = \sum_{j=1}^N D_{r,j}(\Delta_N) \lambda_j + F_r(\Delta_N), \quad r = \overline{1, N}. \tag{28}$$

Substituting the corresponding right-hand sides of (28) into the conditions (16) – (18), we obtain a system of linear algebraic equations with respect to the parameters $\lambda_r, r = \overline{1, N+1}$:

$$[I + D_{N,N}(\Delta_N)] \lambda_N + \sum_{j=1}^{N-1} D_{N,j}(\Delta_N) \lambda_j = -F_N(\Delta_N), \tag{29}$$

$$\sum_{\substack{j=1 \\ j \neq 2}}^N D_{1,j}(\Delta_N) \lambda_j - [I - D_{1,2}(\Delta_N)] \lambda_2 = -F_1(\Delta_N), \tag{30}$$

$$\begin{aligned}
 & [I + D_{s,s}(\Delta_N)]\lambda_s - [I - D_{s,s+1}(\Delta_N)]\lambda_{s+1} + \\
 & + \sum_{\substack{j=1 \\ j \neq s, j \neq s+1}}^N D_{s,j}(\Delta_N) \lambda_j = -F_s(\Delta_N), \quad s = \overline{2, N-1}.
 \end{aligned} \tag{31}$$

We denote the matrix corresponding to the left side of the system of equations (29) - (31) by $Q_*(\Delta_N)$ and write the system in the form

$$Q_*(\Delta_N)\lambda = -F_*(\Delta_N), \quad \lambda \in R^{(P-1)N}, \tag{32}$$

where $F_*(\Delta_N) = (F_N(\Delta_N), F_1(\Delta_N), \dots, F_{N-1}(\Delta_N)) \in R^{(P-1)N}$.

Cauchy problems for ordinary differential equations on subintervals

$$\frac{dx}{dt} = A(t)x + P(t), \quad x(t_{r-1}) = 0, \quad t \in [t_{r-1}, t_r], \quad r = \overline{1, N} \tag{33}$$

are a significant part of proposed algorithm. Here $P(t)$ is either $(n \times n)$ matrix, or n vector, both continuous on $[t_{r-1}, t_r]$, $r = \overline{1, N}$. Consequently, solution to problem (33) is a square matrix or a vector of dimension n .

Denote by $a(P, t)$ the solution to the Cauchy problem (33). Obviously,

$$a(P, t) = X_r(t) \int_{t_{r-1}}^t X_r^{-1}(\tau) P(\tau) d\tau, \quad t \in [t_{r-1}, t_r],$$

where $X_r(t)$ is a fundamental matrix of differential equation (33) on the r -th interval.

We offer the following numerical implementation of algorithm based on the Runge – Kutta method of 4th order and Simpson’s method.

1. Suppose we have a partition $\Delta_N: 0 = t_0 < t_1 < \dots < t_{N-1} < t_N = T$. Divide each r -th interval $[t_{r-1}, t_r]$, $r = \overline{1, N}$, into N_r parts with step $h_r = (t_r - t_{r-1})/N_r$. Assume on each interval $[t_{r-1}, t_r]$ the variable \hat{t} takes its discrete values: $\hat{t} = t_{r-1}, \hat{t} = t_{r-1} + h_r, \dots, \hat{t} = t_{r-1} + (N_r - 1)h_r, \hat{t} = t_r$, and denote by $\{t_{r-1}, t_r\}$ the set of such points.

2. Using the Runge Kutta method of 4th order, we find the numerical solutions to Cauchy problems

$$\frac{dx}{dt} = A(t)x + \Phi(t), \quad x(t_{r-1}) = 0, \quad t \in [t_{r-1}, t_r],$$

and define the values of $(n \times n)$ matrices $a^{h_r}(\Phi, \hat{t})$ on the set $\{t_{r-1}, t_r\}$, $r = \overline{1, N}$.

3. Using the values of $(n \times n)$ matrices $\Psi(s)$ and $a^{h_r}(\Phi, \hat{t})$ on $\{t_{r-1}, t_r\}$, and Simpson’s method, we calculate the $(n \times n)$ matrices

$$\hat{\Phi}_r^{h_r}(\Phi) = \int_{t_{r-1}}^{t_r} \Psi(\tau) a^{h_r}(\Phi, \tau) d\tau, \quad r = \overline{1, N}.$$

Summing up the matrices $\hat{\Phi}_r^{h_r}(\Phi)$ over r , we find the $(n \times n)$ matrices $G^{\tilde{h}}(\Delta_N) = \sum_{j=1}^N \hat{\Phi}_j^{h_j}(\Phi)$, where $\tilde{h} = (h_1, h_2, \dots, h_N) \in R^P$.

4. Solving the Cauchy problems

$$\frac{dx}{dt} = A(t)x + A(t), \quad x(t_{r-1}) = 0, \quad t \in [t_{r-1}, t_r],$$

$$\frac{dx}{dt} = A(t)x + B(t), \quad x(t_{r-1}) = 0, \quad t \in [t_{r-1}, t_r],$$

$$\frac{dx}{dt} = A(t)x + F(t), \quad x(t_{r-1}) = 0, \quad t \in [t_{r-1}, t_r], \quad r = \overline{1, N},$$

by using again the Runge–Kutta method of 4th order, we find the values of $(n \times n)$ matrices $a(A, \hat{t})$, $a(B, \hat{t})$ and n vector $a(F, \hat{t})$ on $\{t_{r-1}, t_r\}$, $r = \overline{1, N}$.

5. Applying Simpson’s method on the set $\{t_{r-1}, t_r\}$, we evaluate the definite integrals

$$\hat{\Phi}_r^{h_r} = \int_{t_{r-1}}^{t_r} \Psi(\tau) d\tau, \quad \hat{\Phi}_r^{h_r}(A) = \int_{t_{r-1}}^{t_r} \Psi(\tau) a^{h_r}(A, \tau) d\tau,$$

$$\hat{\Phi}_r^{h_r}(B) = \int_{t_{r-1}}^{t_r} \Psi(\tau) a^{h_r}(B, \tau) d\tau, \quad \hat{\Phi}_r^{h_r}(F) = \int_{t_{r-1}}^{t_r} \Psi(\tau) a^{h_r}(F, \tau) d\tau, \quad r = \overline{1, N}.$$

By the equalities

$$\begin{aligned} V_1^{\tilde{h}}(\Delta_N) &= \sum_{j=1}^N \hat{\Phi}_j^{h_j}(B), \\ V_r^{\tilde{h}}(\Delta_N) &= \hat{\Phi}_r^{h_r}(A) + \sum_{j=1}^N \hat{\Phi}_j^{h_j}(\Phi) \cdot \hat{\Phi}_r^{h_r}, \quad r = \overline{2, N}, \\ g^{\tilde{h}}(F, \Delta_N) &= \sum_{j=1}^N \hat{\Phi}_j^{h_j}(F). \end{aligned}$$

we define the $(n \times n)$ matrices $V_r^{\tilde{h}}(\Delta_N)$ and n vectors $g^{\tilde{h}}(F, \Delta_N)$, $r = \overline{1, N}$.

6. Construct the system of linear algebraic equations with respect to parameters

$$Q_*^{\tilde{h}}(\Delta_N)\lambda = -F_*^{\tilde{h}}(\Delta_N), \quad \lambda \in R^{(P-1)N}, \quad (34)$$

Solving the system (34), we find $\lambda^{\tilde{h}}$. As noted above, the elements of $\lambda^{\tilde{h}} = (\lambda_1^{\tilde{h}}, \lambda_2^{\tilde{h}}, \dots, \lambda_N^{\tilde{h}})$ are the values of approximate solution to problem (12)-(18) at the left-end points of subintervals.

7. To define the values of approximate solution at the remaining points of set $\{t_{r-1}, t_r\}$, we first find

$$\xi^{\tilde{h}} = [I - G^{\tilde{h}}(\Delta_N)]^{-1} \sum_{r=1}^N V_r^{\tilde{h}}(\Delta_N) \lambda_r^{\tilde{h}} + [I - G^{\tilde{h}}(\Delta_N)]^{-1} g^{\tilde{h}}(F, \Delta_N).$$

and then solve the Cauchy problems

$$\frac{du}{dt} = A(t)u + \Phi(t) \left(\xi^{\tilde{h}} + \sum_{j=2}^N \hat{\Phi}_j^{h_j} \cdot \lambda_j^{\tilde{h}} \right) + B(t)\lambda_1^{\tilde{h}} + F(t),$$

$$u(t_0) = 0, \quad t \in [t_0, t_1],$$

$$\frac{du}{dt} = A(t)u + \Phi(t) \left(\xi^{\tilde{h}} + \sum_{j=2}^N \hat{\Phi}_j^{h_j} \cdot \lambda_j^{\tilde{h}} \right) + B(t)\lambda_1^{\tilde{h}} + F(t),$$

$$u(t_{r-1}) = \lambda_r^{\tilde{h}}, \quad t \in [t_{r-1}, t_r], \quad r = \overline{2, N}.$$

And the solutions to Cauchy problems are found by the Runge–Kutta method of 4th order. Thus, the algorithm allows us to find the numerical solution to the problem (9)-(11).

So, we propose the numerically approximate method for solving of the original problem (1)-(4).

Example. We consider a linear boundary value problem with a parameter for an integro-differential equation of parabolic type

$$\begin{aligned} \frac{\partial u}{\partial t} &= a(x, t) \frac{\partial^2 u}{\partial x^2} + c(x, t)u + b(x, t)\mu(x) + \\ &+ \varphi(x, t) \int_0^T \psi(x, s)u(x, s)ds + f(x, t), \quad (x, t) \in \Omega = (0, \omega) \times (0, T), \end{aligned} \quad (35)$$

$$u(x, 0) = 0, \quad x \in [0, \omega], \quad (36)$$

$$u(x, T) = 0, \quad x \in [0, \omega], \quad (37)$$

$$u(0, t) = \tilde{\psi}_1(t), \quad u(\omega, t) = \tilde{\psi}_2(t), \quad t \in [0, T], \quad (38)$$

where $\omega = 0.5, T = 0.1, a(x, t) = 1, c(x, t) = 0, b(x, t) = t^2 + 1, \varphi(x, t) = x^2, \psi(x, s) = s, f(x, t) = xe^{xt} \sin(10\pi t) + 10\pi e^{xt} \cos(10\pi t) - t^2 e^{xt} \sin(10\pi t) - (t^2 + 1)(x^3 + 1) - \frac{[e^{0.1(\pi x^2 - 20\pi x + 100\pi^3)} - 20\pi x]x^2}{(x^2 + 100\pi^2)^2}, \tilde{\psi}_1(t) = \sin(10\pi t), \tilde{\psi}_2(t) = e^{0.5t} \sin(10\pi t).$

We take $h = 0.1$ and produce a discretization by $x: x_i = ih, i = \overline{0, 5}$.

We introduce the notations $u_i(t) = u(ih, t), \mu_i = \mu(ih), f_i(t) = f(ih, t), i = \overline{0, 5}$.

Problem (35) - (38) is replaced by the following linear boundary value problem with a parameter for an integro-differential equation

$$\frac{du_i}{dt} = \frac{u_{i+1} - 2u_i + u_{i-1}}{0.01} + (t^2 + 1)\mu_i + (0.1 \cdot i)^2 \int_0^{0.1} s \cdot u_i(s) ds + f_i(t), \quad \overline{1,4}, \quad (39)$$

$$u_i(0) = 0, \quad i = \overline{0,5}, \quad (40)$$

$$u_i(0.1) = 0, \quad i = \overline{0,5}, \quad (41)$$

$$u_0(t) = \sin(10\pi t), \quad u_5(t) = e^{0.5t} \sin(10\pi t), \quad t \in [0,0.1]. \quad (42)$$

In view of condition (40)-(42), from (39) we obtain two groups of equations for determining μ_0 and μ_5 :

$$b_0(0)\mu_0 = \dot{\tilde{\psi}}_1(0) - \varphi_0(0) \int_0^T \psi_0(s) \tilde{\psi}_1(s) ds - f_0(0), \text{ then } \mu_0 = 1,$$

$$b_5(0)\mu_5 = \dot{\tilde{\psi}}_2(0) - \varphi_5(0) \int_0^T \psi_5(s) \tilde{\psi}_2(s) ds - f_5(0), \text{ then } \mu_5 = 1.125.$$

The functions $u_0(t)$, $u_5(t)$, and parameters μ_0, μ_5 are known.

Problem (39)-(42) will be rewritten in vector-matrix form

$$\frac{du}{dt} = A(t)u + B(t)\mu + \Phi(t) \int_0^{0.1} \Psi(s)u(s) ds + F(t), \quad u, \mu \in R^4, \quad t \in (0,0.1), \quad (43)$$

$$u(0) = 0, \quad (44)$$

$$u(0.1) = 0, \quad (45)$$

where $u(t) = (u_1(t), u_2(t), u_3(t), u_4(t))$, $\mu = (\mu_1, \mu_2, \mu_3, \mu_4)$ - unknown function and parameter,

$$A(t) = \begin{pmatrix} -200 & 100 & 0 & 0 \\ 100 & -200 & 100 & 0 \\ 0 & 100 & -200 & 100 \\ 0 & 0 & 100 & -200 \end{pmatrix}, \quad B(t) = \begin{pmatrix} t^2 + 1 & 0 & 0 & 0 \\ 0 & t^2 + 1 & 0 & 0 \\ 0 & 0 & t^2 + 1 & 0 \\ 0 & 0 & 0 & t^2 + 10 \end{pmatrix},$$

$$\Phi(t) = \begin{pmatrix} 0.01 & 0 & 0 & 0 \\ 0 & 0.04 & 0 & 0 \\ 0 & 0 & 0.09 & 0 \\ 0 & 0 & 0 & 0.16 \end{pmatrix}, \quad \Psi(s) = \begin{pmatrix} s & 0 & 0 & 0 \\ 0 & s & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & s \end{pmatrix},$$

$$F(t) = \begin{pmatrix} f(0.1, t) + 100 \sin(10\pi t) \\ f(0.2, t) \\ f(0.3, t) \\ f(0.4, t) + 100e^{0.5t} \sin(10\pi t) \end{pmatrix}.$$

Here we use the numerical implementation of algorithm. Accuracy of solution depends on the accuracy of solving the Cauchy problem on subintervals and evaluating definite integrals. We provide the results of the numerical implementation of algorithm by partitioning the interval $[0, 0.1]$ with step $h = 0.002$.

Solution to problem (35)-(38) is pair $(u^*(x, t), \mu^*(x))$, where $u^*(x, t) = e^{xt} \sin(10\pi t)$, $\mu^*(x) =$

$$x^3 + 1. \text{ Then solution to problem (43)-(45) is pair } (u^*(t), \mu^*), \text{ where } u^*(t) = \begin{pmatrix} e^{0.1t} \sin(10\pi t) \\ e^{0.2t} \sin(10\pi t) \\ e^{0.3t} \sin(10\pi t) \\ e^{0.4t} \sin(10\pi t) \end{pmatrix},$$

$$\mu^* = \begin{pmatrix} 1.001 \\ 1.008 \\ 1.027 \\ 1.064 \end{pmatrix} \text{ and the following estimates } \max \|\mu^* - \tilde{\mu}\| < 0.00009, \text{ and } \max_{j=\overline{0,50}} \|u^*(t_j) - \tilde{u}(t_j)\| < 0.000004 \text{ is true.}$$

УДК 517.958:536.2; 517.968.7; 519.62/.64
МРНТИ 27.35.45; 27.33.19; 27.41.19

А.Т. Асанова^{1,2}, Э.А. Бакирова^{1,2,3}, Ж.М. Кадирбаева^{1,2,4}

¹Математика және математикалық модельдеу институты, Алматы, Қазақстан;

²Ақпараттық және есептеуіш технологиялар институты, Алматы, Қазақстан;

³Қазақ Ұлттық қыздар педагогикалық университеті, Алматы, Қазақстан;

⁴Халықаралық ақпараттық технологиялар университеті, Алматы, Қазақстан

ПАРАБОЛАЛЫҚ ТЕКТЕС ИНТЕГРАЛДЫҚ-ДИФФЕРЕНЦИАЛДЫҚ ТЕНДЕУЛЕР ҮШІН БАСҚАРУ ЕСЕБІН ШЕШУДІҢ САНДЫҚ ЖУЫҚТАЛҒАН ӘДІСІ

Аннотация. Параболалық тектес интегралдық-дифференциалдық тендеулер үшін параметрі бар сызықтық шеттік есеп зерттеледі. Кеңістіктік айнымалыны дискреттеу көмегімен қарастырылатын есеп жәй интегралдық-дифференциалдық тендеулер жүйесі үшін параметрі бар сызықтық екі нүктелі шеттік есеппен аппроксимацияланады. Алынған есепті шешу үшін параметрлеу әдісі қолданылады. Аппроксимацияланған есеп Фредгольм интегралдық дифференциалдық тендеулер жүйесі үшін арнайы Коши есептерінен, шеттік шарттардан және бөлу нүктелерінде шешімнің үзіліссіз шарттарынан тұратын пара-паресепке келтіріледі. Параметрлері бар жәй дифференциалдық тендеулер жүйесі үшін Коши есебін шешу дифференциалдық тендеудің фундаменталдық матрицасы көмегімен құрылады. Параметрлерге қатысты сызықты алгебралық тендеулер жүйесі тиісті нүктелердің мәндерін шеттік шарт пен үзіліссіз шарттарына қою арқылы құрылады. Есепті шешудің құрылған жүйе мен ішкі аралықтарда Коши есебін шешудің 4-ші ретті Рунге-Кутта әдісіне негізделген сандық әдісі ұсынылады.

Кілттік сөздер: параболикалық тектес дербес туындылы интегралдық-дифференциалдық тендеулер, параметрі бар есеп, аппроксимация, сандық жуықталған әдіс, алгоритм.

УДК 517.958:536.2; 517.968.7; 519.62/.64
МРНТИ 27.35.45; 27.33.19; 27.41.19

А.Т. Асанова^{1,2}, Э.А. Бакирова^{1,2,3}, Ж.М. Кадирбаева^{1,2,4}

¹Институт математики и математического моделирования, Алматы, Казахстан;

²Институт информационных и вычислительных технологий, Алматы, Казахстан;

³Казахский Национальный Женский Педагогический Университет, Алматы, Казахстан;

⁴Международный Университет Информационных Технологий, Алматы, Казахстан

ЧИСЛЕННО ПРИБЛИЖЕННЫЙ МЕТОД РЕШЕНИЯ ЗАДАЧИ УПРАВЛЕНИЯ ДЛЯ ИНТЕГРО-ДИФФЕРЕНЦИАЛЬНЫХ УРАВНЕНИЙ ПАРАБОЛИЧЕСКОГО ТИПА

Аннотация. Исследуется линейная краевая задача с параметром для интегро-дифференциальных уравнений параболического типа. С помощью дискретизации пространственной переменной рассматриваемая задача аппроксимируется линейной двухточечной краевой задачей с параметром для системы интегро-дифференциальных уравнений. Для решения полученной задачи применяется метод параметризации. Аппроксимирующая задача сведена к эквивалентной задаче, состоящей из специальной задачи Коши для системы интегро-дифференциальных уравнений Фредгольма, краевых условий и условий непрерывности решения в точках разбиения. Решение задачи Коши для системы обыкновенных дифференциальных уравнений с параметрами строится с использованием фундаментальной матрицы дифференциального уравнения. Система линейных алгебраических уравнений относительно параметров составляется путем подстановки значений соответствующих точек в краевое условие и условия непрерывности. Предлагается численный метод решения задачи, основанный на решении построенной системы и метода Рунге-Кутты 4-го порядка для решения задачи Коши на подинтервалах.

Ключевые слова: интегро-дифференциальные уравнения с частными производными параболического типа, задача с параметром, аппроксимация, численно приближенный метод, алгоритм.

Information about authors:

Assanova Anar Turmaganbetkyzy - themember of Editorial Board of journal "News of the NAS RK. Physico-Mathematical Series", Institute of Mathematics and Mathematical Modeling, Institute of Information and Computational Technologies, chief scientific researcher, Doctor of Physical and Mathematical Sciences, professor, assanova@math.kz; <https://orcid.org/0000-0001-8697-8920>;

Bakirova Elmira Ainabekovna - Institute of Mathematics and Mathematical Modeling, Institute of Information and Computational Technologies, leading scientific researcher, Candidate of Physical and Mathematical Sciences, ass. professor, Kazakh National Women's Teacher Training University, professor, bakirova1974@mail.ru; <https://orcid.org/0000-0002-3820-5373>;

Kadirbayeva Zhazira Muratbekovna - Institute of Mathematics and Mathematical Modeling, Institute of Information and Computational Technologies, leading scientific researcher, Candidate of Physical and Mathematical Sciences, International Information Technology University, assistant professor, zh.kadirbayeva@gmail.com; <https://orcid.org/0000-0001-8861-4100>

REFERENCES

- [1] Sloan I. and Thomée V. Time discretization of an integro-differential equation of parabolic type // *SIAM Journal on Numerical Analysis*, 23(1986), 1052–1061.
- [2] Cannon J. and Lin Y. Non-classical H1 projection and Galerkin methods for non-linear parabolic integro-differential equations // *Calcolo*, 25(1988), 187–201.
- [3] Thomée V. and Zhang N.-Y. Error estimates for semidiscrete finite element methods for parabolic integro-differential equations // *Mathematics of Computation*, 53(1989), 121–139.
- [4] Lin Y., Thomée V., and Wahlbin L. Ritz-Volterra projections to finite element spaces and applications to integro-differential and related equations // *SIAM Journal on Numerical Analysis*, 28(1991), 1047–1070.
- [5] Pani A. and Peterson E. Finite element methods with numerical quadrature for parabolic integrodifferential equations // *SIAM Journal on Numerical Analysis*, 33(1996), 1084–1105.
- [6] Lin Y. Semi-discrete finite element approximations for linear parabolic integrodifferential equations with integrable kernels // *Journal of Integral Equations and Applications*, 10(1998), 51–83.
- [7] Pani A. and Sinha R. Finite element approximation with quadrature to a time dependent parabolic integro-differential equation with nonsmooth initial data // *Journal of Integral Equations and Applications*, 13(2001), 35–72.
- [8] Pani A. and Fairweather G. H1-Galerkin mixed finite element methods for parabolic partial integro-differential equations // *IMA Journal of Numerical Analysis*, 22(2002), 231–252.
- [9] McLean W., Sloan I. and Thomée V. Time discretization via Laplace transformation of an integro-differential equation of parabolic type // *IMA Journal of Numerical Analysis*, 24(2004), 439–463.
- [10] Matache A.-M., Schwab C. and Wihler T. Fast numerical solution of parabolic integrodifferential equations with applications in finance // *SIAM Journal on Scientific Computing*, 27(2005), 369–393.
- [11] Sinha R., Ewing R. and Lazarov R. Some new error estimates of a semidiscrete finite volume element method for a parabolic integro-differential equation with nonsmooth initial data // *SIAM Journal on Numerical Analysis*, 43(2006), 2320–2344.
- [12] Zhang N. On fully discrete Galerkin approximations for partial integrodifferential equations of parabolic type // *Mathematics of Computation*, 60(1993), 133–166.
- [13] Dolejší V. and Vlasák M. Analysis of a BDF-DGFE scheme for nonlinear convection-diffusion problems // *NumerischeMathematik*, 110(2008), 405–447.
- [14] Volpert V. *Elliptic partial differential equations Vol. 2: Reaction-Diffusing Equations*, Birkhauser Springer, Basel etc., 2014.
- [15] Douglas J. and Jones B. Numerical methods for integro-differential equations of parabolic and hyperbolic types // *NumerischeMathematik*, 4(1962), 96–102.
- [16] Yanik E. and Fairweather G. Finite element methods for parabolic and hyperbolic partial integro-differential equations // *Nonlinear Analysis*, 12(1988), 785–809.
- [17] Pani A., Thomée V., and Wahlbin L. Numerical methods for hyperbolic and parabolic integro-differential equations // *Journal of Integral Equations and Applications*, 4(1992), 533–584.
- [18] Dzhumabayev D.S. Criteria for the unique solvability of a linear boundary-value problem for an ordinary differential equation // *U.S.S.R. Computational Mathematics and Mathematical Physics*. 1989. Vol. 29, No. 1. P. 34-46.
- [19] Dzhumabaev D.S. A method for solving the linear boundary value problem for an integro-differential equation // *Computational Mathematics and Mathematical Physics*. 2010. Vol. 50, No. 7. P. 1150-1161.
- [20] Dzhumabaev D.S. An algorithm for solving a linear two-point boundary value problem for an integro-differential equation // *Computational Mathematics and Mathematical Physics*. 2013. Vol. 53, No. 6. P. 736-758.
- [21] Dzhumabaev, D.S., Bakirova, E.A. Criteria for the unique solvability of a linear two-point boundary value problem for systems of integro-differential equations // *Differential Equations*. 2013. Vol. 49, No. 9. P. 1087-1102.
- [22] Dzhumabaev D.S. Necessary and sufficient conditions for the solvability of linear boundary-value problems for the Fredholm integro-differential equations // *Ukrainian Mathematical Journal*. 2015. Vol. 66, No. 8. P. 1200-1219.
- [23] Dzhumabaev D.S. Solvability of a linear boundary value problem for a Fredholm integro-differential equation with impulsive inputs // *Differential Equations*. 2015. Vol. 51, No. 9. P. 1180-1196.
- [24] Dzhumabaev D.S. On one approach to solve the linear boundary value problems for Fredholm integro-differential equations // *Journal of Computational and Applied Mathematics*. 2016. Vol. 294, P. 342-357.
- [25] Dzhumabaev D.S. Bakirova, E.A., Kadirbayeva Zh.M. An algorithm for solving a control problem for a differential equation with a parameter // *News of the NAS RK. Phys.-Math. Series*. 2018. Vol. 5, No. 321. P.25-32. DOI: <https://doi.org/10.32014/2018.2518-1726.4>
- [26] Assanova A.T., Bakirova E.A., Kadirbayeva Zh.M. Numerical implementation of solving a boundary value problem for a system of loaded differential equations with parameter // *News of the NAS RK. Phys.-Math. Series*. 2019. Vol. 3, No. 325. -P. 77-84. DOI: <https://doi.org/10.32014/2019.2518-1726.27>

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы *М. С. Ахметова, Т. А. Апендиев, Д. С. Аленов*
Верстка на компьютере *А. М. Кульгинбаевой*

Подписано в печать 12.12.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
9,6 п.л. Тираж 300. Заказ 6.