

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Әль-фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
Әль-Фараби атындағы
Қазақ ұлттық университетінің

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-farabi kazakh
national university

SERIES
PHYSICO-MATHEMATICAL

6(328)

NOVEMBER – DECEMBER 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Бас редакторы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі **Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев У.У. проф. корр.-мүшесі(Қазақстан)
Жүсіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Д. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қырғыстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорус)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик(Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы»РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18, <http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жангаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф., чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)
Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.
Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

Editor in chief
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d :

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskiy I.N. prof., academician(Ukraine)
Kovalev A.M. prof., academician(Ukraine)
Mikhalevich A.A. prof., academician(Belarus)
Pashayev A. prof., academician(Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician(Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.76>

Volume 6, Number 328 (2019), 90 – 105

UDC 517.929

A.Sh. Shaldanbayev¹, M.B. Ivanova², A.N.Urmatova³, A.A. Shaldanbayeva⁴

¹Silkway International University, Shymkent, Kazakhstan;

²South Kazakhstan medical Academy, Shymkent, Kazakhstan;

³South Kazakhstan State University M.O.Auezov, Shymkent, Kazakhstan;

⁴Regional social-innovative University, Shymkent, Kazakhstan.

shaldanbaev51@mail.ru, marina-iv@mail.ru, urmatova81@bk.ru, altima_a@mail.ru

SPECTRAL DECOMPOSITION OF A FIRST ORDER FUNCTIONAL DIFFERENTIAL OPERATOR

Abstract. In this paper we study spectral properties of a boundary value problem of a first-order differential equation with constant coefficients and a deviating argument. By spectral properties we mean completeness and basis property of the system of eigenfunctions and associated functions of the boundary value problem, as well as the Volterra property.

Keywords: equation with deviating argument, completeness, basis property, Volterra property, Sturm-Liouville operator, Riesz basis.

1. Introduction

In applications the following eigenvalue problem is often occurred in a more general form [1, p. 520]:

$$Au = \lambda Su.$$

If the operator S is determined by the equality $(Sf)(x) = f(-x)$, let us say, for all $f \in L_2$, then the following generalized spectral problem occurs:

$$Au = \lambda u(-x).$$

In the case when A is a differential operator, we get a differential equation with deviating argument.

Theory of differential equations with deviating argument is the subject of a huge number of works, among which we note only monographs by A.D. Myshkis [2], L.E. Elsgolts and S.B. Norkin [3], which provides an extensive bibliography. Study of the Sturm - Liouville type boundary value problems for an equation with deviating argument is the subject of a monograph by S.B. Norkin [4]. In this and other works, the spectral questions of differential equations with deviations in higher terms, where there is no spectral parameter, are investigated. Only a few works are devoted to the case when the deviating argument is contained in a term with a spectral parameter. In this regard, we note the works of T.Sh. Kalmenov, S.T. Akhmetova and A.Sh. Shaldanbaev [5], A.M. Ibraimkulov [6], S.T. Akhmetova [7].

Apparently, main theorems of the theory of solvability of differential equations with deviating (delayed) arguments were formulated in the monograph by A.D. Myshkis [2].

T.Sh. Kalmenov [5] laid the foundation to a systematic study of spectral questions for a first-order differential equation with a deviating argument of the indicated form. These ideas were developed in [7].

Apparently, generalized spectral problems of the type

$$Au = \lambda u(-x),$$

where A is some differential operator of the first order, were first investigated, at the initiative of T.Sh. Kalmenova, in [5]. In [7], various spectral properties of the generalized spectral problem were studied:

$$\begin{aligned}y'(x) &= \lambda y(1-x), \quad 0 < x < 1, \\ \alpha y(0) + \beta y(1) &= 0,\end{aligned}$$

including conditions of self-adjointness, Volterra property and basis property.

Theorem 1.7.7 of the above paper [7] states the Riesz basis property of the system of root functions of the considered generalized spectral problem for

$$(|\alpha| + |\beta|)(\alpha^4 - \beta^4) \neq 0.$$

Results of this work were transferred to the interval $[-1, 1]$, by using another method, in [8], where the final solution to basis property questions of root functions of the generalized spectral problem is given

$$\begin{aligned}u'(-x) &= \lambda u(x), \quad -1 < x < 1, \\ u(-1) &= \alpha u(1).\end{aligned}$$

From results of this work it follows that any such correct boundary-value problem is either Volterra or the system of its root functions forms a Riesz basis.

The method of [5] was generalized in [9], in particular, in this paper two abstract theorems on eigenvalues and root vectors of the operators A and A^2 , which may be of independent interest, are proved. We give their statement.

Consider a linear operator A in the Hilbert space H . We suppose that domain $D(A)$ of the operator A is dense in H . Then there exists an operator A^* , which is conjugate to the operator A . Let spectrum of the operator A be discrete. The following proposition holds.

Theorem 1.1. Let number λ_0^2 be an eigenvalue of the operator A^2 . If the number λ_0 is not eigenvalue of the operator A , then λ_0 is an eigenvalue of the operator A .

In the next theorem we consider the case of root vectors.

Theorem 1.2. Let λ_0^2 be an eigenvalue of the operator A^2 . If λ_0 is not eigenvalue of the operator A , then any root vector u_1 of the operator A^2 (if, ofcourse, it exists), corresponding to the eigenvalue λ_0^2 , is an associated element of the operator A , corresponding to the eigenvalue λ_0 .

In the work of A.M. Ibraimkulov [6] completeness of the root vectors of a second-order equation was studied. The studies of this author were continued in [10] - [14].

Among recent studies we can note works of W.Watkins [15–16], in which questions on solvability of one-dimensional differential equations with involution were considered, A.P. Khromov and his followers [17-18], which considered solvability of integral and differential equations in partial derivatives with involution.

The variable separation method for solving partial differential equations is based on the spectral theory of one-dimensional differential operators. Spectral theory of self-adjoint and non-self-adjoint ordinary differential operators, which originated in the bowels of mathematical physics equations and began with the classical works of Sturm, Liouville, Steklov and others, has received quite complete development over the past century. The spectral theory of self-adjoint ordinary differential operators is almost complete. In the field of the spectral theory of non-self-adjoint ordinary differential operators, significant results on completeness and basis property of eigenfunctions and associated functions were obtained in the works of M.V. Keldysh [19], V.A. Ilyin [20-25], M. Otelbaev [26], A.A. Shkalikov [27], Radziewsky [28] and many other mathematicians.

Theory of basis property of systems of eigenfunctions and associated functions of non-self-adjoint ordinary differential operators, proposed by V.A. Ilyin, got rapid development. A fairly complete idea about development of the theory of basis property by V.A. Ilyin was given in review articles [29-30].

Compared with the spectral theory of ordinary differential operators, the spectral theory of one-dimensional differential operators with involution is in its infancy. Apparently, the first works on the spectral theory of one-dimensional differential operators with involution were carried out by initiative of T.Sh. Kalmenov [31-35] in the last decade of the present century. These studies were continued in the cycle of works by M.A. Sadybekov and A.M. Sarsenby [36-41]. Over the past decade, researchers' interest

in differential equations with deviating arguments has grown markedly, as evidenced by publications [42–58]. Theory of bases is described in detail in [60–63].

In this work, we continue the studies begun in [5].

Problem Formulation. Find a spectral decomposition of the operator

$$Au = u'(1 - x), \quad x \in (0,1), \quad (1)$$

$$D(A) = \{u(x) \in C^1(0,1) \cap C[0,1]: \alpha u(0) + \beta u(1) = 0\}, \quad (2)$$

where α, β are arbitrary complex numbers, satisfying the condition:

$$|\alpha| + |\beta| \neq 0. \quad (3)$$

2. Research Method.

The method is based on the following theorem of N.K. Bari [58].

Theorem. If the sequence $\{\psi_j\}$ is complete in a Hilbert space H , it corresponds to the complete biorthogonal sequence $\{\varphi_j\}$ for any $f \in H$

$$\sum_{j=1}^{\infty} |(f, \psi_j)|^2 < \infty, \quad \sum_{j=1}^{\infty} |(f, \varphi_j)|^2 < \infty, \quad (4)$$

then the system $\{\psi_j\}$ forms a Riesz basis of the Hilbert space H .

Therefore, we first show completeness of the system of eigenfunctions of the operator (1) - (2); then we find complete biorthogonal systems of functions and prove inequalities (4).

2.1. On spectrum of the operator.

Consider the following boundary value problem:

$$y'(1 - x) = \lambda y(x); \quad x \in (0,1), \quad (5)$$

$$\alpha y(0) + \beta y(1) = 0 \quad (6)$$

where α, β are arbitrary complex numbers, satisfying the condition (3), and λ is a spectral parameter.

It is easy to note that a general solution of the equation (5) has the form

$$y(x, \lambda) = A \left[\cos \lambda \left(\frac{1}{2} - x \right) - \sin \lambda \left(\frac{1}{2} - x \right) \right], \quad (7)$$

where $A \neq 0$ is an arbitrary nonzero constant.

Putting (7) into the boundary condition (2), we have

$$\begin{aligned} \alpha u(0) + \beta u(1) &= \alpha A \left(\cos \frac{\lambda}{2} - \sin \frac{\lambda}{2} \right) + \beta A \left(\cos \frac{\lambda}{2} + \sin \frac{\lambda}{2} \right) = \\ &= A \left[(\alpha + \beta) \cos \frac{\lambda}{2} - (\alpha - \beta) \sin \frac{\lambda}{2} \right] = A \Delta(\lambda) = 0, \end{aligned}$$

since $A \neq 0$, therefore

$$\Delta(\lambda) = (\alpha + \beta) \cos \frac{\lambda}{2} - (\alpha - \beta) \sin \frac{\lambda}{2} = 0.$$

Assuming, that $\alpha^2 - \beta^2 \neq 0$, we get

$$\begin{aligned} \alpha - \beta \neq 0, \quad \alpha + \beta \neq 0, \Rightarrow \\ \operatorname{tg} \frac{\lambda}{2} = \frac{\alpha + \beta}{\alpha - \beta}, \quad \frac{\lambda_n}{2} = n\pi + \operatorname{arctg} \frac{\alpha + \beta}{\alpha - \beta}, \Rightarrow \\ \lambda_n = 2n\pi + 2 \operatorname{arctg} \frac{\alpha + \beta}{\alpha - \beta}. \end{aligned}$$

Lemma 2.1. If

$$\alpha^4 - \beta^4 \neq 0, \quad (8)$$

then the boundary value problem

$$y'(1-x) = \lambda y(x); x \in (0,1), \quad (5)$$

$$\alpha y(0) + \beta y(1) = 0 \quad (6)$$

has infinite set of eigenvalues

$$\lambda_n = 2n\pi + 2 \operatorname{arctg} \frac{\alpha + \beta}{\alpha - \beta}, \quad n = 0, \pm 1, \pm 2, \dots \quad (9)$$

and the eigenfunctions corresponding to them

$$y_n(x) = A_n \left[\cos \lambda_n \left(\frac{1}{2} - x \right) - \sin \lambda_n \left(\frac{1}{2} - x \right) \right], \quad n = 0, \pm 1, \pm 2, \dots \quad (10)$$

where A_n are arbitrary constants.

All eigenvalues are simple, i.e. if λ_n is an eigenvalue, then $\Delta(\lambda_n) = 0$ and $\dot{\Delta}(\lambda_n) \neq 0$, where the icon $(\dot{})$ means the derivative with respect to the spectral parameter λ .

There are no associated functions.

Proof. From the condition of Lemma it follows that $\alpha + \beta \neq 0$, $\alpha - \beta \neq 0$, then from the equation

$$\Delta(\lambda) = (\alpha + \beta) \cos \frac{\lambda}{2} - (\alpha - \beta) \sin \frac{\lambda}{2} = 0,$$

we have

$$\operatorname{tg} \frac{\lambda}{2} = \frac{\alpha + \beta}{\alpha - \beta}.$$

This equation does not have any roots only in two cases:

$$\frac{\alpha + \beta}{\alpha - \beta} = \pm i.$$

This condition holds only when $\alpha = \pm i\beta$, i.e. $\alpha^2 + \beta^2 = 0$, which is possible due to the condition of Lemma 3.1.

In all other cases our equation has roots, which are given by the formulas

$$\lambda_n = 2n\pi + 2 \operatorname{arctg} \frac{\alpha + \beta}{\alpha - \beta}, \quad n = 0, \pm 1, \pm 2, \dots$$

By (7) we find the corresponding eigenfunctions:

$$y_n(x) = A_n \left[\cos \lambda_n \left(\frac{1}{2} - x \right) - \sin \lambda_n \left(\frac{1}{2} - x \right) \right], \quad (10)$$

where A_n are arbitrary constants.

If λ is a multiple root of the equation $\Delta(\lambda) = 0$, then the system of equations

$$\begin{cases} \Delta(\lambda) = 0, \\ \dot{\Delta}(\lambda) \neq 0 \end{cases}$$

implies that $\alpha^2 + \beta^2 = 0$, that is also impossible.

Indeed,

$$\begin{aligned} \Delta(\lambda) &= (\alpha + \beta) \cos \frac{\lambda}{2} - (\alpha - \beta) \sin \frac{\lambda}{2} = 0, \\ \dot{\Delta}(\lambda) &= \frac{1}{2} \left[-(\alpha + \beta) \sin \frac{\lambda}{2} - (\alpha - \beta) \cos \frac{\lambda}{2} \right] = 0, \Rightarrow \\ &\begin{cases} (\alpha + \beta) \cos \frac{\lambda}{2} - (\alpha - \beta) \sin \frac{\lambda}{2} = 0, \\ (\alpha - \beta) \cos \frac{\lambda}{2} + (\alpha + \beta) \sin \frac{\lambda}{2} = 0. \end{cases} \end{aligned}$$

Since this system of equations has a nontrivial solution, then its determinant will vanish, i.e.

$$\Delta = \begin{vmatrix} \alpha + \beta & -(\alpha - \beta) \\ \alpha - \beta & \alpha + \beta \end{vmatrix} = (\alpha + \beta)^2 + (\alpha - \beta)^2 = 2(\alpha^2 + \beta^2) = 0.$$

Definition 2.1. If $u(x)$ is an eigenfunction of the boundary value problem

$$\begin{aligned} Au &= u'(1-x) = \lambda u(x); x \in (0,1), \\ \alpha u(0) + \beta u(1) &= 0, \end{aligned}$$

then solution of the boundary value problem

$$\begin{aligned} Bv &= v'(1-x) - \lambda v(x) = u(x), \\ \alpha v(0) + \beta v(1) &= 0 \end{aligned}$$

is called associated function of this boundary value problem.

Now we show that if the condition $\alpha^4 - \beta^4 \neq 0$ holds, our boundary value problem (5) - (6) does not have associated functions.

Let $u(x)$ be an eigenfunction of the boundary value problem (5) - (6), and $v(x)$ is its corresponding associated function. Then differentiating (5) - (6) by the spectral parameter λ , we have

$$\begin{aligned} \dot{u}'(1-x) &= \lambda \dot{u} + u(x), \quad \alpha \dot{u}(0) + \beta \dot{u}(1) = 0, \quad => \\ \dot{u}'(1-x) - \lambda \dot{u} &= u(x), \quad \alpha \dot{u}(0) + \beta \dot{u}(1) = 0. \end{aligned}$$

Consequently, the difference $z(x) = \dot{u}(x) - v(x)$ is an eigenfunction of our boundary value problem (5) - (6). Then, obviously, the function

$$\dot{u}(x) = v(x) + z(x)$$

is an associated function of our boundary value problem. We prove that it is not possible. Indeed,

$$\begin{aligned} \alpha u(0) + \beta u(1) = 0 &= \left| u(x) = A \left[\cos \lambda \left(\frac{1}{2} - x \right) - \sin \lambda \left(\frac{1}{2} - x \right) \right] \right| = \\ &= \alpha A \left(\cos \frac{\lambda}{2} - \sin \frac{\lambda}{2} \right) + \beta A \left(\cos \frac{\lambda}{2} - \sin \frac{\lambda}{2} \right) = \\ &= A \left[(\alpha + \beta) \cos \frac{\lambda}{2} - (\alpha - \beta) \sin \frac{\lambda}{2} \right] = A \cdot \Delta(\lambda). \end{aligned}$$

Differentiating this formula by the spectral parameter λ , we get

$$\begin{aligned} \frac{d}{d\lambda} [\alpha u(0) + \beta u(1)] &= A \cdot \dot{\Delta}(\lambda), \\ \alpha \dot{u}(0) + \beta \dot{u}(1) &= A \cdot \dot{\Delta}(\lambda) \neq 0, \end{aligned}$$

where λ is an eigenvalue of the boundary value problem (5) - (6).

2.2. On completeness.

Lemma 2.2. If

$$\alpha^4 - \beta^4 \neq 0, \tag{8}$$

then eigenfunctions of the boundary value problem

$$y'(x) = \lambda y(1-x); x \in (0,1), \tag{5}$$

$$\alpha y(0) + \beta y(1) = 0 \tag{6}$$

form a complete system in the space $L^2(0,1)$.

Proof. Let $\{y_n\}, n = 0, \pm 1, \pm 2, \dots$ be a system of eigenfunctions of the boundary value problem (5) - (6). We assume that

$$\int_0^1 \overline{f(x)} y_n(x) dx = 0, \quad \int_0^1 \overline{f(x)} y_{-n}(x) dx = 0, n = 0, \pm 1, \pm 2, \dots$$

then

$$\int_0^1 \overline{f(x)} \left[\cos(2n\pi + 2\varphi) \left(\frac{1}{2} - x \right) - \sin(2n\pi + 2\varphi) \left(\frac{1}{2} - x \right) \right] dx = 0,$$

$$\int_0^1 \overline{f(x)} \left[\cos(2n\pi - 2\varphi) \left(\frac{1}{2} - x \right) + \sin(2n\pi - 2\varphi) \left(\frac{1}{2} - x \right) \right] dx = 0,$$

where

$$\varphi = \operatorname{arctg} \frac{\alpha + \beta}{\alpha - \beta}.$$

Supposing $t = \frac{1}{2} - x$, from the first formula we have

$$\begin{aligned} x &= \frac{1}{2} - t, & dx &= -dt, \Rightarrow \\ & \int_{-1/2}^{1/2} \overline{f} \left(\frac{1}{2} - t \right) [\cos(2n\pi + 2\varphi)t - \sin(2n\pi + 2\varphi)t] (-dt) = \\ & = \int_{-1/2}^{1/2} \overline{f} \left(\frac{1}{2} - t \right) [\cos(2n\pi + 2\varphi)t - \sin(2n\pi + 2\varphi)t] dt = 0; \end{aligned} \quad (12)$$

Similarly, we obtain

$$= \int_{-1/2}^{1/2} \overline{f} \left(\frac{1}{2} - t \right) [\cos(2n\pi - 2\varphi)t + \sin(2n\pi - 2\varphi)t] dt = 0. \quad (13)$$

Summing up equalities (12) and (13), we get

$$\begin{aligned} & \int_{-1/2}^{1/2} \overline{f} \left(\frac{1}{2} - t \right) [2 \cos 2n\pi t \cos 2\varphi t - 2 \cos 2n\pi t \sin 2\varphi t] dt = 0, \\ & \int_{-1/2}^{1/2} \overline{f} \left(\frac{1}{2} - t \right) (\cos 2\varphi t - \sin 2\varphi t) \cos 2n\pi t dt = 0, n = 0, 1, 2, \dots \end{aligned}$$

In this formula supposing $2\pi t = x$, we have

$$\begin{aligned} t &= \frac{x}{2\pi}, & dt &= \frac{dx}{2\pi}, \\ \int_{-\pi}^{+\pi} \overline{f} \left(\frac{1}{2} - \frac{x}{2\pi} \right) \left(\cos \frac{\varphi x}{\pi} - \sin \frac{\varphi x}{\pi} \right) \cos nx dx &= 0, n = 0, 1, 2, \dots \end{aligned} \quad (14)$$

Further, subtracting the formula (13) from (12), we get

$$\begin{aligned} & \int_{-1/2}^{1/2} \overline{f} \left(\frac{1}{2} - t \right) [-2 \sin 2n\pi \cdot \sin 2\varphi t - 2 \sin 2n\pi t \cdot \cos 2\varphi t] dt = 0, \Rightarrow \\ & \int_{-1/2}^{1/2} \overline{f} \left(\frac{1}{2} - t \right) [\cos 2\varphi t + \sin 2\varphi t] \sin 2n\pi t dt = 0, \quad n = 1, 2, \dots \end{aligned} \quad (15)$$

Now from (14) we have

$$\int_{-\pi}^{+\pi} \overline{f} \left(\frac{1}{2} - \frac{x}{2\pi} \right) \left(\cos \frac{\varphi x}{\pi} - \sin \frac{\varphi x}{\pi} \right) \cos nx dx = 0. \Rightarrow$$

In this formula assuming , we obtain

$$-x = \xi, dx = -d\xi, =>$$

$$-\int_{-\pi}^{+\pi} \bar{f}\left(\frac{1}{2} + \frac{\xi}{2\pi}\right) \left(\cos \frac{\varphi\xi}{\pi} + \sin \frac{\varphi\xi}{\pi}\right) \cos n\xi d\xi = 0, =>$$

$$\int_{-\pi}^{+\pi} \bar{f}\left(\frac{1}{2} + \frac{\xi}{2\pi}\right) \left(\cos \frac{\varphi\xi}{\pi} + \sin \frac{\varphi\xi}{\pi}\right) \cos n\xi d\xi = 0,$$

or denoting ξ by x , we get

$$\int_{-\pi}^{+\pi} \bar{f}\left(\frac{1}{2} + \frac{\varphi}{2\pi}\right) \left(\cos \frac{\varphi x}{\pi} + \sin \frac{\varphi x}{\pi}\right) \cos nx dx = 0, n = 0,1,2, \dots \quad (16)$$

Adding (14) to (16), we receive

$$\int_{-\pi}^{+\pi} \left[\frac{\bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right) + \bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right)}{2} \cos \frac{\varphi x}{\pi} + \frac{\bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right) - \bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right)}{2} \sin \frac{\varphi x}{\pi} \right] \cdot \cos nx dx = 0. \quad (17)$$

The function in the integral in (17) is even, therefore

$$\int_0^{\pi} \left(P\bar{f} \cos \frac{\varphi x}{\pi} + Q\bar{f} \sin \frac{\varphi x}{\pi} \right) \cos nx dx = 0, \quad n = 0,1,2, \dots,$$

where

$$P\bar{f}(x) = \frac{\bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right) + \bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right)}{2}, Q\bar{f}(x) = \frac{\bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right) - \bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right)}{2}.$$

Due to completeness of the system of functions $\{\cos nx\}, n = 0,1,2, \dots$ in the space $L^2(0, \pi)$, we have

$$P\bar{f} \cos \frac{\varphi x}{\pi} + Q\bar{f} \sin \frac{\varphi x}{\pi} = 0$$

Now we transform the formula (15). Assuming $t = -x$, we get

$$\int_{-1/2}^{1/2} \bar{f}\left(\frac{1}{2} + x\right) (\cos 2\varphi x - \sin 2\varphi x) (-dx) =$$

$$\int_{-1/2}^{1/2} \bar{f}\left(\frac{1}{2} + x\right) (\cos 2\varphi x - \sin 2\varphi x) dx = 0.$$

Consequently, we have a pair of the formulas

$$\int_{-1/2}^{1/2} \bar{f}\left(\frac{1}{2} - x\right) (\cos 2\varphi x + \sin 2\varphi x) \sin 2n\pi x dx = 0, \quad (18)$$

$$\int_{-1/2}^{1/2} \bar{f}\left(\frac{1}{2} + x\right) (\cos 2\varphi x - \sin 2\varphi x) \sin 2n\pi x dx = 0. \quad (19)$$

Subtracting the formula (18) from the formula (19), then dividing the result by 2, we obtain

$$\int_{-1/2}^{1/2} \left[\frac{\bar{f}\left(\frac{1}{2} + x\right) - \bar{f}\left(\frac{1}{2} - x\right)}{2} \cos 2\varphi x - \frac{\bar{f}\left(\frac{1}{2} + x\right) + \bar{f}\left(\frac{1}{2} - x\right)}{2} \sin 2\varphi x \right] \cdot \sin 2n\pi x dx = 0, \quad n = 1, 2, \dots \tag{20}$$

The function in the integral in (20) is even, thus

$$\int_0^{1/2} (Q\bar{f} \cos 2\varphi x - P\bar{f} \sin 2\varphi x) \sin 2n\pi x dx = 0, \tag{21}$$

where

$$Q\bar{f}(x) = \frac{\bar{f}\left(\frac{1}{2} + x\right) - \bar{f}\left(\frac{1}{2} - x\right)}{2}, \quad P\bar{f}(x) = \frac{\bar{f}\left(\frac{1}{2} + x\right) + \bar{f}\left(\frac{1}{2} - x\right)}{2}.$$

Assuming $t = 2\pi x$, we change the variable in the formula (21), then

$$x = \frac{t}{2\pi}, \quad dx = \frac{dt}{2\pi},$$

$$\int_0^\pi \left[\frac{\bar{f}\left(\frac{1}{2} + \frac{t}{2\pi}\right) - \bar{f}\left(\frac{1}{2} - \frac{t}{2\pi}\right)}{2} \cos \frac{\varphi t}{\pi} - \frac{\bar{f}\left(\frac{1}{2} + \frac{t}{2\pi}\right) + \bar{f}\left(\frac{1}{2} - \frac{t}{2\pi}\right)}{2} \sin \frac{\varphi t}{\pi} \right] \cdot \sin n t dt = 0, \quad n = 1, 2, \dots$$

Due to completeness of the system of functions in the space $L^2(0, \pi)$, we have

$$\frac{\bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right) - \bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right)}{2} \cos \frac{\varphi x}{\pi} - \frac{\bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right) + \bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right)}{2} \sin \frac{\varphi x}{\pi} = 0.$$

Consequently,

$$\begin{cases} \frac{\bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right) + \bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right)}{2} \cos \frac{\varphi x}{\pi} + \frac{\bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right) - \bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right)}{2} \sin \frac{\varphi x}{\pi} = 0, \\ \frac{\bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right) + \bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right)}{2} \sin \frac{\varphi x}{\pi} - \frac{\bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right) - \bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right)}{2} \cos \frac{\varphi x}{\pi} = 0. \end{cases}$$

Since the determinant of this system

$$\Delta = \begin{vmatrix} \cos \frac{\varphi x}{\pi} & \sin \frac{\varphi x}{\pi} \\ \sin \frac{\varphi x}{\pi} & -\cos \frac{\varphi x}{\pi} \end{vmatrix} = -\cos^2 \frac{\varphi x}{\pi} - \sin^2 \frac{\varphi x}{\pi} = -1 \neq 0,$$

then

$$\frac{\bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right) + \bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right)}{2} = 0, \quad \frac{\bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right) - \bar{f}\left(\frac{1}{2} - \frac{x}{2\pi}\right)}{2} = 0.$$

Summing up these formulas, we get $\bar{f}\left(\frac{1}{2} + \frac{x}{2\pi}\right) = 0$ almost for all $x \in [0, \pi]$, consequently, almost for all $x \in [0, 1]$ we have the equality $\bar{f} = 0$, that is required to prove.

2.3. On the conjugate boundary value problem.

We find conjugate boundary value problem to our boundary value problem

$$Ay = y'(1 - x) = \lambda y(x), \quad x \in (0, 1); \tag{5}$$

$$\alpha y(0) + \beta y(1) = 0. \tag{6}$$

Let $z(x) \in D(L^+)$, i.e. belong to the domain of the conjugate problem, then we have the formula

$$(Ay, z) = (y, A^+z), \quad \forall y \in D(A), \quad z \in D(A^+).$$

Expanding this formula and integrating by parts, we find $D(A^+)$.

$$\begin{aligned} (Ay, z) &= \int_0^1 Sy'(x)\overline{z(x)} dx = (Sy', z) = (y', Sz) = |Sz(x) = z(1-x)| = \\ &= \int_0^1 \overline{Sz} dy = Sz \cdot y|_0^1 - \int_0^1 (\overline{Sz})' y(x) dx = \overline{z(1-x)} \cdot y(x)|_0^1 - \int_0^1 (\overline{Sz})' y(x) dx = \\ &= \overline{z(0)}y(1) - \overline{z(1)}y(0) + \int_0^1 \overline{z'(1-x)} y(x) dx. \end{aligned}$$

Equating to zero, outside the integral term, we compose a system of equations:

$$\begin{cases} \alpha y(0) + \beta y(1) = 0 \\ \overline{z(1)}y(0) - \overline{z(0)}y(1) = 0. \end{cases}$$

Since the system of equations has a non-trivial solution, then the determinant vanishes, i.e.

$$-\alpha\overline{z(0)} - \beta\overline{z(1)} = 0, \Rightarrow \overline{z(0)} + \beta\overline{z(1)} = 0, \Rightarrow \overline{\alpha}z(0) + \overline{\beta}z(1) = 0.$$

From the equality

$$(y, A^+z) = \int_0^1 y(x)\overline{z'(1-x)} dx$$

we have

$$A^+z = z'(1-x)$$

consequently, conjugate boundary value problem has the following form:

$$A^+z = z'(1-x) = \mu z(x), \quad x \in (0,1); \quad (5)^+$$

$$\overline{\alpha}z(0) + \overline{\beta}z(1) = 0. \quad (6)^+$$

It is easy to note that this problem similar to the boundary value problem (5) - (6). Since $(\overline{\alpha})^4 - (\overline{\beta})^4 = 0 \Leftrightarrow \alpha^4 - \beta^4 = 0$, then the conditions on its solvability are also similar. In particular, Lemma 2.1 yields the following Lemma 2.3.

Lemma 2.3. If

$$(\overline{\alpha})^4 - (\overline{\beta})^4 \neq 0 \quad (8)^+$$

then the boundary value problem (5)⁺ - (6)⁺ has infinite set of eigenvalues:

$$\mu_m = 2m\pi + 2\arctg \frac{\overline{\alpha} + \overline{\beta}}{\overline{\alpha} - \overline{\beta}}, \quad m = 0, \pm 1, \pm 2, \dots \quad (9)^+$$

and their corresponding eigenfunctions:

$$z_m(x) = B_m \left[\cos \mu_m \left(\frac{1}{2} - x \right) - \sin \mu_m \left(\frac{1}{2} - x \right) \right] \quad m = 0, \pm 1, \pm 2, \dots \quad (10)^+$$

where B_m are arbitrary constants.

All eigenvalues μ_m are simple.

There are no associated functions.

Lemma 2.4. If

$$(\overline{\alpha})^4 - (\overline{\beta})^4 \neq 0 \quad (8)^+$$

then eigenfunctions $\{z_n\}$ of the boundary value problem

$$A^+z = z'(1-x) = \mu z(x), \quad x \in (0,1); \quad (5)^+$$

$$\overline{\alpha}z(0) + \overline{\beta}z(1) = 0 \quad (6)^+$$

form a complete system in the space $L^2(0,1)$.

2.4. On the biorthogonal system

Lemma 2.5. If the functions

$$y_n(x) = A_n \left[\cos \lambda_n \left(\frac{1}{2} - x \right) - \sin \lambda_n \left(\frac{1}{2} - x \right) \right], \quad n = 0, \pm 1, \pm 2, \dots$$

are eigenfunctions of the boundary value problem

$$Ay = y'(1-x) = \lambda y(x), \quad x \in (0,1); \quad (5)$$

$$\alpha y(0) + \beta y(1) = 0, \quad (6)$$

then the functions

$$z_n(x) = \frac{1}{A_n} \left[\cos \bar{\lambda}_n \left(\frac{1}{2} - x \right) - \sin \bar{\lambda}_n \left(\frac{1}{2} - x \right) \right]$$

are eigenfunctions of the conjugate boundary value problem

$$A^+z = z'(1-x) = \mu z(x), \quad x \in (0,1); \quad (5)^+$$

$$\bar{\alpha} z(0) + \bar{\beta} z(1) = 0; \quad (6)^+$$

moreover, we have the formula

$$(y_n, z_m) = \delta_{nm},$$

where δ_{nm} is the Kronecker symbol.

Proof.

$$\begin{aligned} (y_n, z_m) &= \\ &= \int_0^1 \left[\cos \lambda_n \left(\frac{1}{2} - x \right) - \sin \lambda_n \left(\frac{1}{2} - x \right) \right] \cdot \overline{\left[\cos \bar{\lambda}_m \left(\frac{1}{2} - x \right) - \sin \bar{\lambda}_m \left(\frac{1}{2} - x \right) \right]} dx \\ &= \int_0^1 \left[\cos \lambda_n \left(\frac{1}{2} - x \right) - \sin \lambda_n \left(\frac{1}{2} - x \right) \right] \left[\cos \lambda_m \left(\frac{1}{2} - x \right) - \sin \lambda_m \left(\frac{1}{2} - x \right) \right] dx = \\ &= \int_0^1 \left\{ \left[\cos \lambda_n \left(\frac{1}{2} - x \right) \cos \lambda_m \left(\frac{1}{2} - x \right) + \sin \lambda_n \left(\frac{1}{2} - x \right) \sin \lambda_m \left(\frac{1}{2} - x \right) \right] - \right. \\ &\quad \left. - \left[\cos \lambda_n \left(\frac{1}{2} - x \right) \sin \lambda_m \left(\frac{1}{2} - x \right) \right] - \left[\cos \lambda_n \left(\frac{1}{2} - x \right) \sin \lambda_m \left(\frac{1}{2} - x \right) + \right. \right. \\ &\quad \left. \left. + \sin \lambda_n \left(\frac{1}{2} - x \right) \cos \lambda_m \left(\frac{1}{2} - x \right) \right] \right\} dx = \\ &= \int_0^1 \left[\cos(\lambda_n - \lambda_m) \left(\frac{1}{2} - x \right) - \sin(\lambda_n + \lambda_m) \left(\frac{1}{2} - x \right) \right] dx = \\ &= \left[-\frac{\sin(\lambda_n - \lambda_m) \left(\frac{1}{2} - x \right)}{\lambda_n - \lambda_m} \right]_0^1 - \left[\frac{\cos(\lambda_n + \lambda_m) \left(\frac{1}{2} - x \right)}{\lambda_n + \lambda_m} \right]_0^1 = \\ &= \frac{2 \sin \frac{\lambda_n - \lambda_m}{2}}{\lambda_n - \lambda_m} = \frac{\sin \frac{\lambda_n - \lambda_m}{2}}{\frac{\lambda_n - \lambda_m}{2}} = \left| \frac{\frac{\lambda_n}{2} = n\pi + \operatorname{arctg} \frac{\alpha + \beta}{\alpha - \beta}}{\frac{\lambda_n - \lambda_m}{2} = (n - m)\pi} \right| = \\ &= \frac{\sin(n - m\pi)}{(n - m)\pi} = 0, \end{aligned}$$

when $n \neq m$.

If $n = m$, then

$$\begin{aligned} (y_n, y_m) &= \int_0^1 \left[\cos \lambda_n \left(\frac{1}{2} - x \right) - \sin \lambda_n \left(\frac{1}{2} - x \right) \right]^2 dx = \\ &= \int_0^1 \left[1 - 2 \sin \lambda_n \left(\frac{1}{2} - x \right) \cos \lambda_n \left(\frac{1}{2} - x \right) \right] dx = \\ &= \int_0^1 [1 - \sin \lambda_n (1 - 2x)] dx = \left[x - \frac{\cos \lambda_n (1 - 2x)}{2\lambda_n} \right]_0^1 = 1 \end{aligned}$$

2.5. On basis property

Definition 2.1. Sequence $\{\varphi_j\}$ of vectors of the Banach space B is called basis of this space, if each vector $x \in B$ is expended uniquely in a series

$$x = \sum_{j=1}^{\infty} c_j \times \psi_j$$

converging by the norm of the space B .

Any bounded invertible operator transforms any orthonormal basis into some other basis of the space H . The basis $\{\psi_j\}$ of the Hilbert space H , obtained from the orthonormal basis by using such transformation, is called basis equivalent to orthonormal or the Riesz basis.

Theorem (N.K. Bari). If the sequence $\{\psi_j\}$ is complete in the Hilbert space H , it corresponds to a complete biorthogonal sequence $\{\psi_j\}$ and for all $f \in H$

$$\sum_{j=1}^{\infty} |(f, \psi_j)|^2 < \infty, \sum_{j=1}^{\infty} |(f, \varphi_j)|^2 < \infty \quad (4)$$

then the system $\{\psi_j\}$ forms Riesz basis in the space H .

Using this theorem, we prove basis property of the system of eigenfunctions of our boundary value problem

$$Ay = y'(1 - x) = \lambda y(x), \quad x \in (0, 1); \quad (5)$$

$$\alpha y(0) + \beta y(1) = 0, \quad (6)$$

where α, β are arbitrary constants, satisfying the condition

$$\alpha^4 - \beta^4 \neq 0. \quad (8)$$

and λ is a spectral parameter.

Our sequence is complete (see Lemma 2.2) in the space $H = L^2(0, 1)$, and it corresponds to a biorthogonal sequence $\{z_n\}$ (see Lemma 2.5), which is also complete in H (see Lemma 2.4), therefore, it only remains for us to prove inequalities (4):

Let

$$a_n = (f, y_n) = \int_0^1 f \cdot \bar{y}_n(x) dx, \quad b_n = (f, z_n) = \int_0^1 f(x) \bar{z}_n(x) dx. \quad (22)$$

In our case

$$y_n(x) = (-1)^n \left[\cos \lambda_n \left(\frac{1}{2} - x \right) - \sin \lambda_n \left(\frac{1}{2} - x \right) \right],$$

$$z_n(x) = (-1)^n \left[\cos \bar{\lambda}_n \left(\frac{1}{2} - x \right) - \sin \bar{\lambda}_n \left(\frac{1}{2} - x \right) \right].$$

First we transform the integrals (6).

$$\begin{aligned} a_n &= (-1)^n \int_0^1 f(x) \left[\cos \bar{\lambda}_n \left(\frac{1}{2} - x \right) - \sin \bar{\lambda}_n \left(\frac{1}{2} - x \right) \right] dx = \\ &= (-1)^n \cdot \int_0^1 f(x) \cos \bar{\lambda}_n \left(\frac{1}{2} - x \right) dx - (-1)^n \int_0^1 f(x) \sin \bar{\lambda}_n \left(\frac{1}{2} - x \right) dx; \\ \int_0^1 f(x) \cos \bar{\lambda}_n \left(\frac{1}{2} - x \right) dx &= \left| \begin{array}{l} t = \frac{1}{2} - x \\ dt = -dx \end{array} \right| = \int_{\frac{1}{2}}^{\frac{1}{2}} f \left(\frac{1}{2} - t \right) \cos \bar{\lambda}_n t (-dt) = \end{aligned}$$

$$\begin{aligned}
&= \int_{-\frac{1}{2}}^{\frac{1}{2}} f\left(\frac{1}{2} - t\right) [\cos 2n\pi t \cos 2\varphi t - \sin 2n\pi t \sin 2\varphi t] dt = \\
&= \left| x = 2\pi t, \quad t = \frac{x}{2\pi}, \quad dt = \frac{dx}{2\pi} \right| = \\
&= \int_{-\pi}^{+\pi} f\left(\frac{1}{2} - \frac{x}{2\pi}\right) \left[\cos nx \cdot \cos \frac{\varphi x}{\pi} - \sin nx \cdot \sin \frac{\varphi x}{\pi} \right] \frac{dx}{2\pi} = \\
&= \frac{1}{2\pi} \int_{-\pi}^{+\pi} f\left(\frac{1}{2} - \frac{x}{2\pi}\right) \cos \frac{\varphi x}{\pi} \cos nx \, dx - \\
&\quad - \frac{1}{2\pi} \int_{-\pi}^{+\pi} f\left(\frac{1}{2} - \frac{x}{2\pi}\right) \sin \frac{\varphi x}{\pi} \cdot \sin nx \, dx;
\end{aligned}$$

The system

$$\frac{1}{\sqrt{2\pi}}, \quad \frac{1}{\sqrt{\pi}} \sin x, \dots, \frac{\cos nx}{\sqrt{\pi}}, \quad \frac{\sin nx}{\sqrt{\pi}}, \dots$$

forms orthonormal basis of the space $H = L^2(0,1)$. Based on this fact, we estimate the coefficients a_n and b_n (see (22)).

$$\begin{aligned}
a_n &= \frac{1}{2\sqrt{\pi}} \int_{-\pi}^{+\pi} f\left(\frac{1}{2} - \frac{x}{2\pi}\right) \cos \frac{\varphi x}{\pi} \times \frac{\cos nx}{\sqrt{\pi}} - \\
&\quad - \frac{1}{2\sqrt{\pi}} \int_{-\pi}^{+\pi} f\left(\frac{1}{2} - \frac{x}{2\pi}\right) \sin \frac{\varphi x}{\pi} \times \frac{\sin nx}{\sqrt{\pi}} \, dx, \\
Rea_n &= \frac{1}{2\sqrt{\pi}} \int_{-\pi}^{+\pi} Re f\left(\frac{1}{2} - \frac{x}{2\pi}\right) \cos \frac{\varphi x}{\pi} \times \frac{\cos nx}{\sqrt{\pi}} \, dx - \\
&\quad - \frac{1}{2\sqrt{\pi}} \int_{-\pi}^{+\pi} Ref\left(\frac{1}{2} - \frac{x}{2\pi}\right) \sin \frac{\varphi x}{\pi} \times \frac{\sin nx}{\sqrt{\pi}} \, dx, \quad n \neq 0
\end{aligned}$$

Let

$$u(x) = \frac{Ref\left(\frac{1}{2} - \frac{x}{2\pi}\right) \cos \frac{\varphi x}{\pi}}{2\sqrt{\pi}}, \quad v(x) = \frac{1}{2\sqrt{\pi}} Ref\left(\frac{1}{2} - \frac{x}{2\pi}\right) \sin \frac{\varphi x}{\pi}.$$

Then

$$Rea_n = \frac{1}{\sqrt{\pi}} \int_{-\pi}^{+\pi} u(x) \cos nx \, dx - \frac{1}{\sqrt{\pi}} \int_{-\pi}^{+\pi} v(x) \sin nx \, dx = \alpha_n - \beta_n, \quad n \neq 0$$

If $n = 0$, we have

$$\begin{aligned}
a_0 &= \frac{1}{2\pi} \int_{-\pi}^{+\pi} f\left(\frac{1}{2} - \frac{x}{2\pi}\right) \cos \frac{\varphi x}{\pi} \, dx, \\
Rea_0 &= \frac{1}{2\pi} \int_{-\pi}^{+\pi} Ref\left(\frac{1}{2} - \frac{x}{2\pi}\right) \cos \frac{\varphi x}{\pi} \, dx, \\
\alpha_0 &= \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{+\pi} u(x) \, dx = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{+\pi} \frac{Ref\left(\frac{1}{2} - \frac{x}{2\pi}\right) \cos \frac{\varphi x}{\pi}}{2\sqrt{\pi}} \, dx = \\
&= \frac{1}{\sqrt{2}} \times \frac{1}{2\pi} \int_{-\pi}^{+\pi} Ref\left(\frac{1}{2} - \frac{x}{2\pi}\right) \cos \frac{\varphi x}{\pi} \, dx = \frac{Rea_0}{\sqrt{2}}, \Rightarrow Rea_0 = \alpha_0 \sqrt{2}.
\end{aligned}$$

Further,

$$\begin{aligned} |Rea_n| &\leq |\alpha_n| + |\beta_n|, \Rightarrow \\ |Rea_n|^2 &\leq (|\alpha_n| + |\beta_n|)^2 \leq |\alpha_n|^2 + 2|\alpha_n| \times |\beta_n| + |\beta_n|^2 \leq \\ &\leq |\alpha_n|^2 + |\beta_n|^2 + |\alpha_n|^2 + |\beta_n|^2 \leq 2(|\alpha_n|^2 + |\beta_n|^2), \\ |Rea_0| &\leq \sqrt{2} |\alpha_0|, \quad |Rea_0|^2 \leq 2|\alpha_0|^2. \end{aligned}$$

Consequently,

$$\sum_{n=0}^{\infty} |Rea_n|^2 \leq 2 \left(\sum_{n=0}^{\infty} |\alpha_n|^2 + \sum_{n=1}^{\infty} |\beta_n|^2 \right) \leq 2(\|u\|^2 + \|v\|^2) < \infty.$$

Similarly, we have

$$\sum_{n=0}^{\infty} |Ima_n|^2 < \infty.$$

Therefore,

$$\sum_{n=0}^{\infty} |a_n|^2 = \sum_{n=0}^{\infty} |Ima_n|^2 + |Rea_n|^2 < \infty.$$

Estimation of the series $\sum_{n=0}^{\infty} |b_n|^2$ is carried out similarly.

We have proved the main Theorem 3.1., and Theorem 3.2 is its corollary.

3. Research Results.

Theorem 3.1. Suppose that

$$\alpha^4 - \beta^4 \neq 0, \tag{8}$$

then the system of eigenfunctions of the boundary value problem

$$y'(1-x) = \lambda y(x); x \in (0,1), \tag{9}$$

$$\alpha y(0) + \beta y(1) = 0 \tag{6}$$

forms Riesz basis in the space $L^2(0,1)$, i.e. we have

$$f(x) = \sum_{-\infty}^{+\infty} (f, z_n) y_n(x),$$

Converging in the space $L^2(0,1)$, where

$$y_n(x) = (-1)^n \left[\cos \lambda_n \left(\frac{1}{2} - x \right) - \sin \lambda_n \left(\frac{1}{2} - x \right) \right],$$

$$z_n(x) = (-1)^n \left[\cos \bar{\lambda}_n \left(\frac{1}{2} - x \right) - \sin \bar{\lambda}_n \left(\frac{1}{2} - x \right) \right],$$

$$\lambda_n = 2n\pi + 2 \operatorname{arctg} \frac{\alpha + \beta}{\alpha - \beta}, \quad n = 0, \pm 1, \pm 2, \dots \tag{9}$$

and $f(x)$ is an arbitrary element in the space $L^2(0,1)$.

Theorem 3.2. If $\alpha^4 - \beta^4 \neq 0$, then for any element $u(x) \in D(A)$ we have the spectral expansion

$$Au = \sum_{-\infty}^{+\infty} \lambda_n (u, z_n) y_n(x) \tag{11}$$

converging in the space $L^2(0,1)$, where

$$\lambda_n = 2n\pi + 2 \operatorname{arctg} \frac{\alpha + \beta}{\alpha - \beta}, \quad n = 0, \pm 1, \pm 2, \dots \tag{9}$$

$$z_n(x) = (-1)^n \left[\cos \bar{\lambda}_n \left(\frac{1}{2} - x \right) - \sin \bar{\lambda}_n \left(\frac{1}{2} - x \right) \right], \tag{10}^+$$

$$y_n(x) = (-1)^n \left[\cos \lambda_n \left(\frac{1}{2} - x \right) - \sin \lambda_n \left(\frac{1}{2} - x \right) \right]. \tag{10}$$

Theorem 3.2. is the main result of this work.

4. Discussion.

Formula (11) is not possible if there are associated vectors, the well-known Kesselman - Mikhailov theorem [61- [62] states that, not the system of eigenvectors, the system of root vectors is basic, and this is significance of the results of this work. Formula (11) can find application in electrical engineering, information theory, crystallography, and signal transmission theory. It can be useful in study various boundary value problems by the method of variables separation.

5. Conclusion.

1) Operator (1) - (2) is not semi-bounded;

2) There is an alternative: either the boundary value problem (5) - (6) is Volterra i.e. has no eigenvalues, or the system of its eigenvectors forms a Riesz basis of the space $L^2(0,1)$.

ОӘЖ 517.929

А.Ш. Шалданбаев¹, М.Б. Иванова², А.Н. Урматова³, А.А. Шалданбаева⁴

¹Халықаралық Silkway университеті, Шымкент қ., Қазақстан;

²Оңтүстік Қазақстан медициналық академиясы, Шымкент қ., Қазақстан;

³М.О.Ауезов атындағы Оңтүстік Қазақстан мемлекеттік университеті, Шымкент қ., Қазақстан

⁴Аймақтық әлеуметтік-инновациялық университеті, Шымкент қ., Қазақстан;

БІРІНШІ РЕТТІ ФУНКЦИОНАЛ-ДИФФЕРЕНЦИАЛ ОПЕРАТОРДЫҢ СПЕКТРӘЛДІ ТАРАЛЫМЫ

Аннотация. Бұл еңдекте аргументі ауытқыған бірінші ретті дифференциалдық теңдеудің спектрәлдік қасиеттері зерттелді. Шекаралық есептің спектрәлдік қасиеттері ретінде біз оның меншікті және олармен еншілес функцияларының системасының толымдылығы мен базистігін, сондайақ, оның вольтерлігін танымыз.

Түйін сөздер: аргументі ауытқыған теңдеу, толымдылық, базистік, вольтерлік, Штурм -Лиувиллдің операторы, Рисстің базисі.

УДК 517.929

А.Ш.Шалданбаев¹, М.Б.Иванова², А.Н.Урматова³, А.А.Шалданбаева⁴

¹Международный университет Silkway, г. Шымкент, Казахстан;

²Южно-Казахстанская медицинская академия, г. Шымкент, Казахстан;

³ЮКГУ имени М.Ауезова, г. Шымкент, Казахстан;

⁴Региональный социально-инновационный университет, г. Шымкент, Казахстан.

СПЕКТРАЛЬНОЕ РАЗЛОЖЕНИЕ ФУНКЦИОНАЛЬНО-ДИФФЕРЕНЦИАЛЬНОГО ОПЕРАТОРА ПЕРВОГО ПОРЯДКА

Аннотация. В настоящей работе получено спектральное разложение функционально-дифференциального оператора первого порядка, с помощью прямого доказательства полноты системы собственных функций и теоремы Н.К.Бари.

Ключевые слова: уравнение с отклоняющимся аргументом, полнота, базисность, вольтерровость, операторы Штурма-Лиувилля, базис Рисса.

Information about authors:

Shaldanbayev A.Sh. – doctor of physico-mathematical Sciences, associate Professor, head of the center for mathematical modeling, «Silkway» International University, Shymkent; <http://orcid.org/0000-0002-7577-8402>

Ivanova M.B. - South Kazakhstan medical Academy, Shymkent; <https://orcid.org/0000-0001-6548-0699>

Urmatova A.N. - SKSU named After M. Auezov, senior lecturer of the Department of mathematics, <https://orcid.org/0000-0002-1323-1639>

Shaldanbayeva A.A. - "Regional Social-Innovative University", Shymkent; <https://orcid.org/0000-0003-2667-3097>

REFERENCES

- [1] Kato T. Perturbation Theory of Linear Operators M. Mir. 1972. 740p.
- [2] Myshkis A.D. Linear Differential Equations with delayed argument, M. 1972. 352 p.
- [3] Elsgolts L.E. and Norkin S.B. Introduction to theory of differential equations with deviating argument, M. 1971. 296 p.
- [4] Norkin S.B. Differential Equations with Delayed Argument, M. 1965. 356 p.
- [5] Kalmenov T.Sh., Akhmetova S. and Shaldanbaev A.Sh. To spectral theory of equations with deviating argument // Mathematical Journal, Almaty. 2004. V. 4, No. 3. P. 41-48.

- [6] Ibraimkulov A.M. On spectral properties of boundary value problem for an equation with a deviating argument // News of AS.Kaz.SSR, ser.phys.-math. 1988. No. 3. Pp. 22-25.
- [7] Akhmetova S.T. Spectral properties of boundary value problems for one class of differential equations with deviating argument: diss. ... cand. physical and mathematical sciences: 01.01.02. Shymkent, 2004. 96 p.
- [8] Sadybekov M. A. and Sarsenbi A. M. Solutions of basic spectral problems of all boundary value problems for one first-order differential equation with a deviating argument. Uzbek Mathematical Journal. 2007. No. 3. Pp. 49-53.
- [9] Sarsenbi A.M. On eigenvalues of degrees of linear operators. Vestnik L. N. Gumilyov ENU. 2006. No. 4 (50).- Pp. 20-24.
- [10] Sadybekov M. A. and Sarsenbi A.M. On notion of regularity of boundary value problems for a second-order differential equation with a deviating argument// Mathematical Journal. 2007. v. 7, No. 1 (18). Pp. 82-88.
- [11] Sarsenbi A.M. On asymptotic behavior of eigenvalues of boundary value problems for a second-order differential equation with a deviating argument// Izvestiya NAS RK, Ser. Physical and Math.2007. No. 1. Pp. 49-52.
- [12] Sadybekov M. A. and Sarsenbi A. M. Unconditional convergence of spectral expansions associated with a second-order differential equation with a deviating argument// Vestnik KazNU. 2006. No. 2 (49). Pp. 48-54.
- [13] Sarsenbi A.M. On Reisz basis of a system of eigenfunctions of one boundary value problem for the equation $-u''(-x) = \lambda u(x)$ // Izvestiya NAS RK. 2007. Ser. Phys.-Math. No. 3. Pp. 32-34.
- [14] Sarsenbi A. M. Irregular boundary value conditions in the case of differential equations with deviating argument //Reports of NAS RK. 2007. No. 1. Pp. 12-16.
- [15] W.Watkins. Modified Wiener equations. Int. J. Math. Math. Sci.Vol-27. p: 347-356. 2001y.
- [16] W.T.Watkins. Asymptotic properties of differential equations with involutions. Int. J.Pure Appl. Math. Vol 44. p: 485-492, 2008 y.
- [17] Burlutskaya M. Sh. and Khromov A. P. Foundation of the Fourier method in mixed problems with involution. // Izv. Sarat. Univ. New ser. Ser. Of Math. Mech. and Inf.2011. Vol. 11, No. 4. pp. 3-12.
- [18] Burlutskaya M. Sh. and Khromov A. P. Fourier Method in a mixed problem for a first-order partial differential equation with involution.// J. Comput. Math. and Math. Phys. 2011. v. 51, No. 12. Pp. 2233-2246.
- [19] Keldysh M. V. On completeness of eigenfunctions of certain classes of nonselfadjoint linear operators // Uspekhi Math. Nauk. - 1971. - Vol. 26, No. 4 (160). pp. 15-41.
- [20] Il'in V. A. Existence of a reduced system of eigenand associated functions for a nonselfadjoint ordinary differential operator // Number theory, mathematical analysis and their applications, Trudy Mat. Inst. Steklov. 1976. 142. P. 148-155.
- [21] Il'in V. A. Necessary and sufficient conditions of basis property and equivalent convergence with trigonometric series for spectral expansions, 1 // Differential Equations. 1980. Vol. 16, No. 5. Pp. 771-794.
- [22] Il'in V. A. Necessary and sufficient conditions of basis property and equivalent convergence with trigonometric series for spectral expansions, 2 // Differential Equations. 1980. Vol. 16, No. 6. Pp. 980-1009.
- [23] Il'in V. A. Necessary and sufficient conditions of basis property in L_2 and equivalent convergence with trigonometric series for spectral expansions and expansions by exponent systems// DAN SSSR. 1983. v. 273, No. 4. Pp. 784-793
- [24] Il'in V. A. On unconditional basis property on a closed interval of systems of eigenfunctions and associated functions of the second order differential operator// DAS SSSR. 1983. Vol. 273, No. 5. Pp. 1048-1053.
- [25] Il'in V. A. On relationship between the type of boundary value conditions and properties of basicity and equivalent convergence with a trigonometric series for expansions on root functions of a non-self-adjoint differential operator// Differential Equations. 1994. Vol. 30, No. 9. Pp. 1516-1529.
- [26] Otelbaev M. Estimates of S-numbers and conditions of completeness of a root vector system of a non-self-adjoint Sturm-Liouville operator, Math. Notes, 1979, Vol. 25, No. 3, pp. 409-418.
- [27] Shkalikov A. A. Boundary value problems for ordinary differential equations with parameter in boundary conditions// Trudy sem.by I. G. Petrovsky. 1983. Vol. 9. Pp. 190-229.
- [28] Gomilko A.M. and Radzievsky G. V. Basis properties of eigenfunctions of a regular boundary value problem for a vector functional-differential equation// Differential Equations. 1991. Vol. 27, No. 3. Pp. 385-395.
- [29] Il'in V.A., Kritskov L. V. Properties of spectral expansions corresponding to nonselfadjoint differential operators // J. Math. Sci. (NY). – 2003. 116, N 5. P. 3489-3550.
- [30] Il'in V. A. and Kritskov L. V. Properties of spectral expansions corresponding to non-self-adjoint operators, Functional analysis. Results of science and technology. Ser.: Res.Math. and its App. Them. review., M.: VINITI, 2006, Vol. 96, pp. 5-105.
- [31] Shaldanbaev A. Sh. and Akhmetova S. T. On completeness of eigenvectors of the Cauchy problem. Republican scientific journal "Science and Education SK" № 27, 2002. pp. 58-62.
- [32] Akhmetova S. T., On Bitsadze - Samarsky problem for wave equation, Mathematical Journal, Almaty. 2003, Vol. 3. No. 2 (8), pp. 15-18.
- [33] Shaldanbayev A. Sh. Criteria of Volterra property for differential operator of the first order with deviating argument, Vestnik of Karaganda University, series "Mathematics", № 3(47)/2007, pp. 39-43.
- [34] Sadybekov M. A. and Sarsenbi A. M. Unconditional convergence of spectral expansions associated with a second-order differential equation with deviating argument. // Vestnik KAZNU. 2006. No. 2 (49). Pp. 48-54.
- [35] Sadybekov M. A. and Sarsenbi A. M. Solution of main spectral problems of all boundary value problems for a first-order one-dimensional differential equation with deviating argument. // Uzbek Mathematical Journal - 2007.- No. 3.- pp. 88-94.
- [36] Sadybekov M. A. and Sarsenbi A.M. On notion of regularity of boundary value problems for a second-order differential equation with deviating argument. // Mathematical Journal. 2007. V. 7, No. 1(18). Pp. 82-88.
- [37] Sadybekov M. A. and Sarsenbi A. M. Criterion on basis property of system of eigenfunctions of multiple differentiation operator with involution.// Differential Equation. 2012. v. 48, No. 8. Pp. 1126-1132.
- [38] Kopzhasarova A.A, Sarsenbi A.M. Basis Properties of Eigenfunctions of Second-Order Differential Operators with Involution//Abstract and Applied Analysis. 2012. Volume 2012, Article ID 576843. 6 pages doi:10.1155/2012/576843.

- [39] Kopzhassarova A. A., Lukashov A. L., Sarsenbi A. M. Spectral properties of non-self-adjoint perturbations for a spectral problem with involution // *Abstr. Appl. Anal.* 2012. Pp. 1–5.
- [40] Sarsenbi A.M.; Tengaeva A.A. [On the basis properties of root functions of two generalized eigenvalue problems//](#) *Differential Equations.* 2012. vol 48, no. 2. pp. 306–308.
- [41] Imanbaeva A. B., Kopzhassarova A. A. and Shaldanbaev A. Sh. Asymptotic expansion of solution of singularly perturbed Cauchy problem for a system of ordinary differential equations with constant coefficients. "Izvestiya NAS RK. Series phys. and Math.", 2017, №5, 112 -127.
- [42] Kopzhassarova A. A., Shaldanbaev A. sh. and Imanbaeva A. B. Solution of singularly perturbed Cauchy problem by similarity method. "Izvestiya NAS RK. Series Phys.-Math.", 2017, №5, 127-134.
- [43] Amir Sh. Shaldanbayev, and Manat T. Shomanbayeva, Solution of singularly perturbed Cauchy problem for ordinary differential equation of second order with constant coefficients by Fourier method, Citation: *AIP Conference Proceedings* 1880, 040017 (2017); doi: 10.1063/1.5000633 View online: <http://dx.doi.org/10.1063/1.5000633> View Table of Contents: <http://aip.scitation.org/toc/apc/1880/1> Published by the American Institute of Physics.
- [44] Akylbayev M. I., Beisebaeva A. and Shaldanbayev A. Sh. On periodic solution of the Goursat problem for a wave equation of a special form with variable coefficients (in English), *Izvestiya NAS RK. Series Phys. and Math.*, No. 1. 2018, p.34-50
- [45] A. A. Shaldanbaeva, M.I. Akylbayev, A. Sh. Shaldanbaev, A.Zh. Beisebaeva, The spectral decomposition of cauchy problem's solution for Laplace equation// *News of the National Academy of Sciences of the Republic of Kazakhstan, Physico-mathematical Series*, ISSN 1991-346X Volume 5, Number 321 (2018), 75 – 87. <https://doi.org/10.32014/2018.2518-1726.10>
- [46] A.Sh. Shaldanbayev, A.A. Shaldanbayeva, B.A. Shaldanbay, On projectional orthogonal basis of a linear non-self-adjoint operator, *News of the national academy of sciences of the republic of Kazakhstan, Physico-mathematical series*, ISSN 1991-346X <https://doi.org/10.32014/2019.2518-1726.15>, Volume 2, Number 324 (2019), 79 – 89.
- [47] M.I.Akylbayev, A.Sh.Shaldanbayev, I.Orazov, A.Beysebayeva, About single operator method of solution of a singularly perturbed Cauchy problem for an ordinary differential equation n – order, // *News of the national academy of sciences of the republic of Kazakhstan, Physico-mathematical series*, ISSN 1991-346X Volume 2, Number 324 (2019), 17 – 36. <https://doi.org/10.32014/2019.2518-1726.8>
- [48] Ashyralyev A., Sarsenbi A. M. Well-posedness of an elliptic equations with an involution // *EJDE.* 2015. 284. P. 1–8.
- [49] Kritskov L. V., Sarsenbi A. M. Spectral properties of a nonlocal problem for the differential equation with involution // *Differ. Equ.* 2015. 51, N 8. P. 984– 990.
- [50] Kurdyumov V. P. On Riesz bases of eigenfunction of 2-nd order differential operator with involution and integral boundary conditions // *Izv. Saratov Univ. (N.S.), Ser. Math. Mech. Inform.* 2015. 15, N 4. P. 392–405.
- [51] Kirane M., Al-Salti N. Inverse problems for a nonlocal wave equation with an involution perturbation // *J. Nonlinear Sci. Appl.* 2016. 9. P. 1243–1251.
- [52] Ya. O. Baranetskiy, L. I. Kolyasa, Boundary value problem for second order differential operator equation with involution // *Вісник національного університету, "Львівська політехніка", "Фізико-математичні науки"*, № 871, 2017, pp. 20–26.
- [53] [Kritskov L.V., Sarsenbi A.M., Equiconvergence property for spectral expansions related to perturbations of the operator \$S-u\(x\)\$ with initial data // *Filomat*, 2018, vol. 32, № 3, pp. 1069-1078 .](#)
- [54] [Kritskov L.V., Sadybekov M.A. and Sarsenbi A.M., Nonlocal spectral problem for a second-order differential operator with an involution// *BULLETIN OF THE KARAGANDA UNIVERSITY-MATHEMATICS*, 2018 № 3 \(91\), pp. 53-60.](#)
- [55] [Kritskov L.V., Sadybekov M.A., Sarsenbi A.M., Properties in \$L_p\$ of root functions for a nonlocal problem with involution, *Turkish Journal of Mathematics, Scientific and Technical research Council of Turkey - TUBITAK/Turkiye Bilimsel ve Teknik Arastirma Kurumu* \(Turkey\), 2019, vol. 43, pp. 393-401.](#)
- [56] Vladykina V. E. and Shkalikov A. A., Spectral properties of ordinary differential operators with involution// *Reports of the Academy of Sciences*, 2019, vol. 484, No. 1, pp. 12-17.
- [57] Shaldanbaev A. Sh. and Kulazhanov G. S. on spectral properties of one model operator with deviating argument// *International conference "VEKUA-100"*, Novosibirsk 2007.
- [58] Bari N. K. Biorthogonal systems and bases in Hilbert space, // *Uch. Zap. MSU-* 1951. Vol. 4, ser.148. pp. 69-106.
- [59] Gokhberg I. S. and Krane M. G. Introduction to theory of linear non-self-adjoint operators in Hilbert space, M.: Nauka, 1965, 448 p.
- [60] Naimark M. A. *Linear Differential Operators.* M.: Nauka, 1969. 528 p.
- [61] Kesselman G. M. On unconditional convergence of expansions by eigenfunctions of some differential operators // *Izv. Mathematics.* - 1964.- No. 2 (39). - pp. 82-93.
- [62] Mikhailov V. P. On Riesz basis in $L_2(0;1)$ // *DAN USSR.* 1962. Vol. 144, No. 5. Pp. 981-984.
- [63] Lidsky V. B. Non-self-adjoint operators having trace. *DAN USSR*, 1959, vol. 125. No. 3, pp. 485-488.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы *М. С. Ахметова, Т. А. Апендиев, Д. С. Аленов*
Верстка на компьютере *А. М. Кульгинбаевой*

Подписано в печать 12.12.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
9,6 п.л. Тираж 300. Заказ 6.