

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
әл-Фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
Әл-Фараби атындағы
Қазақ ұлттық университетінің

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-Farabi Kazakh
National University

SERIES
PHYSICO-MATHEMATICAL

1(329)

JANUARY – FEBRUARY 2020

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р ы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі
Ғ.М. Мұтанов

Р е д а к ц и я а л қ а с ы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев У.У. проф. корр.-мүшесі (Қазақстан)
Жүсіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Д. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қырғыстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорусь)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19; 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2020

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Главный редактор
д.ф.-м.н., проф. академик НАН РК
Г.М. Мутанов

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жантаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф., чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физика-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219, 220; тел.: 272-13-19; 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2020

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

Editor in chief

doctor of physics and mathematics, professor, academician of NAS RK

G.M. Mutanov

Editorial board:

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskiy I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19; 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2020.2518-1726.8>

Volume 1, Number 329 (2020), 62 – 70

UDC 517.2, 517.958, 537.6, 539.3, 624.19

L.A. Alexeyeva¹, E.B. Kurmanov^{1,2}

¹Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan;

²Al-Farabi Kazakh National University, Almaty, Kazakhstan

alexeeva@math.kz, kurmanovergali@gmail.com

**GREEN TENSOR OF MOTION EQUATIONS
OF TWO COMPONENTS
BIOT'S MEDIUM BY STATIONARY VIBRATIONS**

Abstract. Here processes of wave propagation in a two-component Biot's medium are considered which are generated by periodic forces actions. By use Fourier transformation of generalized functions, the Green tensor - a fundamental solutions of oscillation equations of this medium has been constructed. This tensor describes the process of propagation of harmonic waves of a fixed frequency in spaces of dimension $N = 1, 2, 3$ under the action of power sources concentrated at the coordinates origin, described by a singular delta -function. Based on it, generalized solutions of these equations are constructed under the action of various sources of periodic perturbations, which are described by both regular and singular generalized functions. For regular acting forces, integral representations of solutions are given that can be used to calculate the stress-strain state of a porous water-saturated medium.

Key words: Biot's medium, solid and liquid components, fundamental solution, generalized direct and inverse Fourier transform, regularization.

Various mathematical models of deformable solids mechanics are used in the study of seismic processes in the earth's crust. The processes of waves propagation are most studied in elastic media. But these models do not take into account many real properties of the ambient array. These are, for example, the presence of groundwater, which complicates the construction and operation of surface and underground structures, affect the magnitude and distribution of stresses. Models, which take into account the water saturation form the earth's crust structures, the presence of gas bubbles, etc., are multi-component medium. A variety of multicomponent media, the complexity of the processes associated with their deformation, lead to a large difference in the methods of analysis and modelling used in the solution of wave problems.

Porous medium saturated with liquid or gas, from the point of view of continuum mechanics, is essentially a two-phase continuous medium, one phase of which is particles of liquid (gas), other solid particles is its elastic skeleton. There are various mathematical models of such media, developed by various authors. The most famous of them are the models of M. Biot, V.N. Nikolaevsky, L.P. Horoshun [1-7]. However, the class of solved tasks to them is very limited and mainly associated with the construction and study of particular solutions of these equations based on the methods of full and partial separation of variables and theory of special functions in the works of Rakhmatullin, H. A., Saatov Ya. U., Filippov I. G., Artykov T. U. [6,7], Erzhanov Zh. S, Ataliev Sh.M., Alexeyeva L.A., Shershnev V.V. [8,9] etc. In this regard, it is important to develop effective methods of solution of boundary value problems for such media with use of modern mathematical methods.

Periodic on time processes are very widespread in practice. By this cause here we consider the process of wave propagation in the Biot's medium, posed by the periodic forces of different types. Based on Fourier transformation of generalized functions we constructed fundamental solutions of oscillation equations of Biot's medium. It is Green's tensor, which describes the process of propagation of harmonic waves at a fixed frequency in the space-time of dimension $N=1,2,3$, under the action of concentrated at

the coordinates origin. By use this tensor we construct generalized solutions of these equations for arbitrary sources of periodic disturbances, which can be described both regular and singular distributions. They can be used to calculate the stress-strain state of a porous water-saturated medium by seismic waves propagation.

1 The parameters and motion equations of a two-components M. Biot medium

The equations of motion of a homogeneous isotropic two-component M. Biot medium are described by the following system of second-order hyperbolic equations [1-3]:

$$\begin{aligned} (\lambda + \mu)\text{grad div}u_s + \mu\Delta u_s + Q\text{grad div}u_f + F^s(x,t) &= \rho_{11}\ddot{u}_s + \rho_{12}\ddot{u}_f \\ Q\text{grad div}u_s + R\text{grad div}u_f + F^f(x,t) &= \rho_{12}\ddot{u}_s + \rho_{22}\ddot{u}_f \end{aligned} \quad (1)$$

$(x,t) \in R^N \times [0, \infty)$. Here N is the dimension of the space. At a plane deformation $N=2$, the total spatial deformation corresponds to $N=3$, at $N=1$ the equations describe the dynamics of a porous liquid-saturated rod. We denote $u_s = u_{sj}(x,t)e_j$ a displacements vector of the elastic skeleton, $u_f = u_{fj}(x,t)e_j$ is the displacements vector of a liquid, e_j ($j = 1, \dots, N$) are the basic orts of the Lagrangian Cartesian coordinate system (everywhere by repeating indices there is summation from 1 to N).

Constants $\rho_{11}, \rho_{12}, \rho_{22}$ have the dimension of mass density and are associated with the density of the masses of the particles, composing a skeleton ρ_s and a fluid ρ_f , by relationships:

$$\rho_{11} = (1 - m)\rho_s - \rho_{12}, \quad \rho_{22} = m\rho_f - \rho_{12},$$

where m is a porosity of medium. The constant of the attached density ρ_{12} is related to the dispersion of the deviation of the micro-velocities of the fluid particles in the pores from the average velocity of the fluid flow and depends on the geometry of the pores. Elastic constants λ, μ are the Lamé parameters of an isotropic elastic skeleton, and Q, R characterize the interaction of the skeleton with a liquid on the basis of *Biot law for stresses*:

$$\begin{aligned} \sigma_{ij} &= (\lambda\partial_k u_k + Q\partial_k U_k)\delta_{ij} + \mu(\partial_i u_j + \partial_j u_i) \\ \sigma &= -mp = R\partial_k U_k + Q\partial_k u_k \end{aligned} \quad (2)$$

Here $\sigma_{ij}(x,t)$ are the stress tensor in the skeleton, $p(x,t)$ is a pressure in the fluid. Further we use the notations for partial derivatives: $\partial_k = \frac{\partial}{\partial x_k}$, $u_{j,k} = \partial_k u_j$, $\Delta = \partial_k \partial_k$ is Laplace operator. The external mass forces acting on the skeleton $F^s = F_j^s(x,t)e_j$ and on the liquid component $F^f = F_j^f(x,t)e_j$.

There are three sound speeds in this medium:

$$c_1^2 = \frac{\alpha_1 + \sqrt{\alpha_1^2 - 4\alpha_2\alpha_3}}{2\alpha_2}, \quad c_2^2 = \frac{\alpha_1 - \sqrt{\alpha_1^2 - 4\alpha_2\alpha_3}}{2\alpha_2}, \quad c_3 = \sqrt{\frac{\rho_{22}\mu}{\alpha_2}} \quad (3)$$

where the next constants were introduced:

$$\alpha_1 = (\lambda + 2\mu)\rho_{22} + R\rho_{11} - 2Q\rho_{12}, \quad \alpha_2 = \rho_{11}\rho_{22} - (\rho_{12})^2, \quad \alpha_3 = (\lambda + 2\mu)R - Q^2$$

The first two speeds c_1, c_2 ($c_1 > c_2$) describe the velocity of propagation of two types of *dilatational waves*. The second slower dilatation wave is called *the repackaging wave*. A third velocity C_3 corresponds to *shear waves* and at $\rho_{12} = 0$ coincides with velocity of transverse waves propagation in an elastic skeleton ($C_3 < C_1$).

We introduce also two velocities of propagation of dilatational waves in corresponding elastic body and in an ideal compressible fluid:

$$c_s = \sqrt{\frac{\lambda + 2\mu}{\rho_{11}}}, \quad c_f = \sqrt{\frac{R}{\rho_{22}}}$$

2 Problems of periodic oscillations of the Biot's medium

Construction of motion equation solutions by periodic oscillations is very important for practice since existing power sources of disturbances are often periodic in time and therefore can be decomposed into a finite or infinite Fourier series in the form:

$$F^s(x, t) = \sum_n F_n^s(x) e^{-i\omega_n t}, \quad F^f(x, t) = \sum_n F_n^f(x) e^{-i\omega_n t} \quad (4)$$

where periods of oscillation of each harmonic $T_n = 2\pi/\omega_n$ are multiple to the general period T of oscillation. Therefore, it is enough to consider the case of stationary oscillations, when the acting forces are periodic on time with an oscillation frequency ω :

$$F^s(x, t) = F^s(x) e^{-i\omega t}, \quad F^f(x, t) = F^f(x) e^{-i\omega t} \quad (5)$$

The solution of the equations (1) can be represented in the similar form:

$$u_s(x, t) = u_s(x) e^{-i\omega t}, \quad u_f(x, t) = u_f(x) e^{-i\omega t} \quad (6)$$

where the complex amplitudes of the displacements $u_s(x)$, $u_f(x)$ must be determined. If the solution has been known for any frequency ω , then we get similar decomposition for the displacements of the medium:

$$u^s(x, t) = \sum_n u_n^s(x) e^{-i\omega_n t}, \quad u^f(x, t) = \sum_n u_n^f(x) e^{-i\omega_n t} \quad (7)$$

which give us the solution of problem for forces (4).

We get equations for complex amplitudes by stationary oscillations, substituting (6) into the system (1):

$$\begin{aligned} (\lambda + \mu) \text{grad div } u^s + \mu \Delta u^s + Q \text{grad div } u^f + F^s(x) &= -\rho_{11} \omega^2 u^s - \rho_{12} \omega^2 u^f \\ Q \text{grad div } u^s + R \text{grad div } u^f + F^f(x) &= -\rho_{12} \omega^2 u^s - \rho_{22} \omega^2 u^f \end{aligned} \quad (8)$$

To construct the solutions of this system we define Green tensor of it.

3 Green tensor of Biot equations by stationary oscillations

Let's construct fundamental solutions of the system (1) in the form:

$$\begin{pmatrix} F^s \\ F^f \end{pmatrix} = \begin{pmatrix} \delta_k^{[j]} e_k \\ \delta_{k+N}^{[j]} e_k \end{pmatrix} \delta(x) e^{-i\omega t}, \quad k = 1, \dots, N, j = 1, \dots, 2N \quad (9)$$

where δ_k^j is Kronecker symbol. They describe the motion of Biot medium at the action of sources of stationary oscillations, concentrated in the point $x=0$. The upper index of this tensor fixes the current concentrated force and its direction. The lower index corresponds to component of the movement of the skeleton and fluid, respectively $k = 1, \dots, N$ and $k = N + 1, \dots, 2N$.

Their complex amplitudes $U_m^j(x, \omega)$ ($j, m = 1, \dots, 2N$) satisfy to the next system of equation:

$$\begin{aligned}
 (\lambda + \mu)U_{j,ji}^k + \mu U_{i,jj}^k + \omega^2 \rho_{11}U_i^k + QU_{j,ji}^{k+N} - \omega^2 \rho_{12}U_i^{k+N} + \delta(x)\delta_j^k &= 0 \\
 QU_{j,ji}^k + \rho_{12}\omega^2 U_i^k + RU_{j,ji}^{k+N} + \rho_{22}\omega^2 U_i^{k+N} + \delta(x)\delta_{j+N}^k &= 0 \\
 j = 1, \dots, 2N, \quad k = 1, \dots, 2N
 \end{aligned}
 \tag{10}$$

Since fundamental solutions are not unique, we'll construct such, which tend to zero at infinity:

$$U_i^j(x, \omega) \rightarrow 0 \quad \text{at} \quad \|x\| \rightarrow \infty \tag{11}$$

and satisfy to radiation condition of type of Somerfield radiation conditions [10]. Matrix of such fundamental equations is names *Green tensor* of Eq. (8).

4 Fourie transform of fundamental solutions

To construct $U_m^j(x, \omega)$ we use the Fourier transformation, which for regular functions has the form:

$$\begin{aligned}
 F[\varphi(x)] &= \bar{\varphi}(\xi) = \int_{R^N} \varphi(x) e^{i(\xi, x)} dx_1 \dots dx_N \\
 F^{-1}[\bar{\varphi}(\xi)] &= \varphi(x) = \frac{1}{(2\pi)^N} \int_{R^N} \bar{\varphi}(\xi) e^{-i(\xi, x)} d\xi_1 \dots d\xi_N
 \end{aligned}$$

where $\xi = (\xi_1, \dots, \xi_N)$ are Fourier variables. Let's apply Fourier transformation to Eqs (10), and use property of Fourier transform of derivatives [10]:

$$\frac{\partial}{\partial x_j} \leftrightarrow -i\xi_j \tag{12}$$

Then we get the system of 2N linear algebraic equations for the Fourier components of this tensor:

$$\begin{aligned}
 -(\lambda + \mu)\xi_j \xi_j \bar{U}_j^k - \mu \|\xi\|^2 \bar{U}_j^k - Q\xi_j \xi_j \bar{U}_{j+N}^k + \rho_{11}\omega^2 \bar{U}_j^k + \rho_{12}\omega^2 \bar{U}_{j+N}^k + \delta_j^k &= 0, \\
 -Q\xi_j \xi_j \bar{U}_j^k - R\xi_j \xi_j \bar{U}_{j+N}^k + \rho_{12}\omega^2 \bar{U}_j^k + \rho_{22}\omega^2 \bar{U}_{j+N}^k + \delta_{j+N}^k &= 0, \\
 j = 1, \dots, N, \quad k = N + 1, \dots, 2N
 \end{aligned}
 \tag{13}$$

By use gradient-divergence method this system has been solved by us. For this the next basic function were introduced

$$f_{0k}(\xi, \omega) = \frac{1}{c_k^2 \|\xi\|^2 - \omega^2}, \quad f_{jk}(\xi, \omega) = \frac{f_{(j-1)k}(\xi, \omega)}{-i\omega}, \quad j = 1, 2; \tag{14}$$

and the next theorem was proved [11,12].

Theorem 1. *Components of Fourier transform of fundamental solutions have the form*

$$j = \overline{1, N}, \quad k = \overline{1, N},$$

$$\overline{U}_j^k = (-i\xi_j)(-i\xi_k)[\beta_1 f_{21} + \beta_2 f_{22} + \beta_3 f_{23}] + \frac{1}{\alpha_2}(\rho_{12}\delta_{j+N}^k - \rho_{22}\delta_j^k) f_{03}$$

$$\overline{U}_{j+N}^k = (-i\xi_j)(-i\xi_k)[\gamma_1 f_{21} + \gamma_2 f_{22} + \gamma_3 f_{23}] - \frac{\mu}{\alpha_2} \delta_{j+N}^k \|\xi\|^2 f_{23} - \frac{1}{\alpha_2}(\rho_{11}\delta_{j+N}^k + \rho_{12}\delta_j^k) f_{03}$$

$$j = 1, \dots, N \quad k = N + 1, \dots, 2N$$

$$\overline{U}_j^k = (-i\xi_j)(-i\xi_{k-N})[\eta_1 f_{21} + \eta_2 f_{22} + \eta_3 f_{23}] + \frac{1}{\alpha_2}(\rho_{12}\delta_{j+N}^k - \rho_{22}\delta_j^k) f_{03}$$

$$\overline{U}_{j+N}^k = (-i\xi_j)(-i\xi_{k-N})[\varsigma_1 f_{21} + \varsigma_2 f_{22} + \varsigma_3 f_{23}] - \frac{\mu}{\alpha_2} \delta_{j+N}^k \|\xi\|^2 f_{23} - \frac{1}{\alpha_2}(\rho_{11}\delta_{j+N}^k + \rho_{12}\delta_j^k) f_{03}$$

where the next constants have been introduced:

$$D_1 = \frac{1}{\alpha_2 \nu_{12}}, \quad \nu_{lm} = c_l^2 - c_m^2, \quad q_1 = Q\rho_{12} - (\lambda + \mu)\rho_{12}, \quad q_2 = \rho_{11}R - Q\rho_{12},$$

$$d_1 = (\lambda + \mu)\rho_{22} - Q\rho_{12}, \quad d_2 = Q\rho_{22} - R\rho_{12}, \quad d_{3j} = \rho_{12}c_j^2 - Q \quad (j=1,2)$$

$$\beta_j = (-1)^{(j+1)} \frac{D_1 c_j^2}{\alpha_2 \nu_{3j}} (d_1 b_{sj} + d_2 d_{3j}), \quad \beta_3 = -\frac{c_3^2}{\alpha_2 \nu_{31} \nu_{32}} (d_1 b_{s3} + d_2 d_{33});$$

$$\gamma_j = (-1)^{j+1} \frac{D_1 c_j^2}{\alpha_2 \nu_{3j}} (q_1 b_{fj} + q_2 d_{3j}), \quad \gamma_3 = -\frac{D_1 c_3^2 \nu_{12}}{\alpha_2 \nu_{31} \nu_{32}} (q_1 b_{f3} + q_2 d_{33});$$

$$\eta_j = (-1)^{j+1} \frac{D_1 c_j^2}{\alpha_2 \nu_{3j}} (d_j d_{3j} + d_2 b_{js}), \quad \eta_3 = -\frac{c_3^2 \nu_{12}}{\alpha_2 \nu_{31} \nu_{32}} (d_1 d_{33} + d_2 b_{3s});$$

$$\varsigma_j = (-1)^{j+1} \frac{D_1 c_j^2}{\alpha_2 \nu_{3j}} (q_1 d_{3j} + q_2 b_{(4-j)s}), \quad \varsigma_3 = -\frac{c_3^2 \nu_{12}}{\alpha_2 \nu_{31} \nu_{32}} (q_1 d_{33} + q_2 b_{3s}) \quad b_{ff} = \rho_{22} \nu_{ff}, b_{sj} = \rho_{11} \nu_{js}.$$

This form is very convenient for constructing originals of Green tensor.

5 Green tensor construction. Radiation conditions

At first let's construct the originals of basic function:

$$\Phi_{0m}(x, \omega) = F^{-1}[f_{0m}(\xi, \omega)],$$

which, in accordance to its definition (14), satisfy to the equation

$$(c_m^2 \|\xi\|^2 - \omega^2) f_{0m} = 1. \quad (15)$$

Using the property (12) for derivatives from here we get Helmholtz equation for fundamental solution (accurate within a factor c_m^{-2}):

$$(\Delta + k_m^2)\Phi_{0m} + c_m^{-2}\delta(x) = 0, \quad k_m = \frac{\omega}{c_m} \tag{16}$$

Fundamental solutions of Helmholtz equation which satisfy to Sommerfeld conditions of radiation: at $r \rightarrow \infty$

$$\begin{aligned} \Phi'_{0m}(r) - ik_m\Phi_{0m}(r) &= O(r^{-1}), & N = 3, \\ \Phi'_{0m}(r) - ik_m\Phi_{0m}(r) &= O(r^{-1/2}), & N = 2. \end{aligned}$$

are well known [10]. They are unique. Using them we obtain:

$$\text{for } N = 3 \quad \Phi_{0m} = \frac{1}{4\pi r c^2} e^{ik_m r}, \quad k_m = \frac{\omega}{c_m};$$

$$\text{for } N = 2 \quad \Phi_{0m} = \frac{i}{4c^2} H_0^{(1)}(k_m r),$$

where $H_j^{(1)}(k_m r)$ is cylindrical Hankel function of the first kind;

$$\text{for } N = 1 \quad \Phi_{0m} = \frac{\sin k_m |x|}{2k_m c_m^2}.$$

These functions (subject to factor $e^{-i\alpha t}$) describe harmonic waves which move from $x=0$ to infinity and decay at infinity. Last property is true only for $N=2,3$. In the case $N=1$ all fundamental solutions of Eq. (16):

$$\left(\frac{d^2}{dx^2} + k_m^2 \right) \Phi_{0m} + c_m^{-2}\delta(x) = 0,$$

don't decay at infinity.

From theorem 1 the next theorem follows.

Theorem 2. The components of Green tensor of Biot's equations at stationary oscillations with frequency ω , satisfying the conditions of radiation, have the form:

$$\text{for } j = \overline{1, N}, \quad k = \overline{1, N},$$

$$U_j^k = -\omega^{-2} \sum_{m=1}^3 \beta_m \frac{\partial^2 \Phi_{0m}}{\partial x_j \partial x_k} + \frac{1}{\alpha_2} (\rho_{12} \delta_{j+N}^k - \rho_{22} \delta_j^k) \Phi_{03},$$

$$U_{j+N}^k = -\omega^{-2} \sum_{m=1}^3 \gamma_m \frac{\partial^2 \Phi_{0m}}{\partial x_j \partial x_k} + \frac{\mu}{\alpha_2 \omega^2} (c_3^{-2} \delta(x) + k_3^2 \Phi_{0m}) \delta_{j+N}^k - \frac{1}{\alpha_2} (\rho_{11} \delta_{j+N}^k + \rho_{12} \delta_j^k) \Phi_{03};$$

for $j = 1, \dots, N \quad k = N + 1, \dots, 2N$

$$U_j^k = -\omega^{-2} \sum_{m=1}^3 \eta_m \frac{\partial^2 \Phi_{0m}}{\partial x_j \partial x_k} + \frac{1}{\alpha_2} (\rho_{12} \delta_{j+N}^k - \rho_{22} \delta_j^k) \Phi_{03}$$

$$U_{j+N}^k = -\omega^{-2} \sum_{m=1}^3 \zeta_m \frac{\partial^2 \Phi_{0m}}{\partial x_j \partial x_k} + \frac{\mu}{\alpha_2 \omega^2} (c_3^{-2} \delta(x) + k_3^2 \Phi_{0m}) \delta_{j+N}^k - \frac{1}{\alpha_2} (\rho_{11} \delta_{j+N}^k + \rho_{12} \delta_j^k) \Phi_{03}$$

where

$$\frac{d^2\Phi_{0m}}{dx^2} = \frac{1}{2c_m^2 k_m} \left(k_m^2 (\sin k_m |x|) - 2k_m \delta(x) \right) \quad \text{for } N = 1,$$

$$\frac{\partial^2\Phi_{0m}}{\partial x_j \partial x_k} = -\frac{i}{4c_m^2} \left(0.5k_m^2 (H_0(k_m r) - H_2(k_m r)) r_{,j} r_{,k} + k H_1^1(k_m r) r_{,jk} \right) \quad \text{for } N = 2,$$

$$\frac{\partial\Phi_{0m}}{\partial x_j \partial x_k} = \frac{1}{4\pi r c_m^2} e^{ikr} \left\{ r_{,j} r_{,k} \left(\left(ik_m - \frac{1}{r} \right)^2 + \frac{1}{r^2} \right) + r_{,jk} \left(ik_m - \frac{1}{r} \right) \right\} \quad \text{for } N = 3;$$

$$k_m = \frac{\omega}{c_m}, \quad r = \|\mathbf{x}\|, \quad r_{,j} = \frac{x_j}{r}, \quad r_{,ij} = \frac{1}{r} \left(\delta_{ij} - \frac{x_i x_j}{r^2} \right).$$

Proof. By use originals of basic functions, property (12) of derivatives, we can obtain from formulas for \overline{U}_j^k in theorem 1 the originals of all addends, beside that which contain factor $\|\xi\|^2$. But using (16) we have

$$-\Delta\Phi = c_m^{-2} \delta(x) + k_m^2 \Phi_{0m} \quad \leftrightarrow \quad \|\xi\|^2 f_{0m} = c_m^{-2} + k_m^2 f_{0m}$$

Then formulas of Theorem 2 follow from formulas of Theorem 1.

Conclusion. Under the action of arbitrary mass forces with frequency ω in the Biot's medium, the solution for complex amplitudes has the form of a tensor-functional convolution:

$$u_j(x, t) = U_j^k(x, \omega) * F_k(x) e^{-i\omega t}, \quad j, k = \overline{1, 2N} \quad (17)$$

Note that mass forces may be different from the space of generalized vector-function, singular and regular. Since Green tensor is singular, contains delta-functions, this convolution are calculated on the rule of convolution in generalized function space. If a support of acting forces are bounded (contained in a ball of finite radius), then all convolutions exist. If supports are not bounded, then the existence condition (17) require some limitations on behavior of forces at infinity which depend on the type of mass forces.

The obtained solutions allow us to study the dynamics of porous water and gas-saturated media at the action of periodic sources of disturbances of a sufficiently arbitrary form. In particular, under the action of certain forces on surfaces, for example cracks, in porous media that can be simulated by simple and double layers on the crack surface.

There is another interesting feature of the Green tensor of the Biot equations, which contains, as one of the terms, the delta function what complicates the application of this tensor for solving boundary value problems based on the boundary element method or boundary untegral equations method [13,14]. Here, when constructing the model, the viscosity of the liquid is not taken into account, which, apparently, leads to the presence of such terms, and requires improvement of this model taking into account the viscosity.

Acknowledgments

This work was supported by the Grant of Ministry of Education and Science of the Republic of Kazakhstan through the program No. AP05132272 "Boundary value problems of the dynamics of deformable solid and electro-magnetic media and their solving"

Л.А. Алексеева¹, Е.Б. Курманов^{1,2}

¹Математика және математикалық модельдеу институты ҚР БҒМ, Алматы, Қазақстан;

²әл-Фараби атындағы Қазақ Ұлттық Университеті, Алматы, Қазақстан

СТАЦИОНАР ТЕРБЕЛІСТЕР КЕЗІНДЕГІ ЕКІ КОМПОНЕНТТІ М. БИО ОРТАСЫНЫҢ ҚОЗҒАЛЫС ТЕҢДЕУІНІҢ ГРИН ТЕНЗОРЫ

Аннотация. Қатты серпімді және идеалды сұйықтан тұратын екі компонентті Био ортаны қарастырамыз. Мұндай ортаның қозғалысы қатты және сұйық компоненттердің орын ауыстыруы үшін гиперболалық типтегі екінші ретті дифференциалдық теңдеулердің біріккен жүйесімен сипатталады. Әртүрлі типтегі периодты күштер тудыратын био ортадағы толқындардың таралу процестері зерттелген.

Бұл жүйенің стационарлы шешімдері $N = 1, 2, 3$ өлшемдер кеңістігіндегі гармоникалық тербелістерді сипаттайды. Бұл жағдайда күрделі тербеліс амплитудасы үшін теңдеулер жүйесі эллиптикалық түрде болады. Жалпыланған функциялардың және оның қасиеттерінің Фурье түрлендіруі негізінде ортаға тербеліс теңдеулерінің іргелі шешімі – сингулярлы дельта функциясымен сипатталған шығу көзіне шоғырланған күш көздерінің әсерінен тұрақты жиіліктегі уақыттық-гармоникалық толқындардың таралуын сипаттайтын Био орта – Грин тензоры тұрғызылды.

Тұрақты және сингулярлы жалпыланған функциялармен сипатталатын периодтық бұзулардың әртүрлі көздерінің әсері кезінде, осы теңдеулердің жалпыланған шешімдері тұрғызылды. Алынған нәтижелерді газ және сұйық қаныққан кеуекті ортадағы толқындық процестерді зерттеу үшін қолдануға болады.

Түйін сөздер: Био ортасы, қатты және сұйық компоненттер, іргелі шешім, жалпыланған тура және кері Фурье түрлендірулері, регуляризация.

Л.А. Алексеева¹, Е.Б. Курманов^{1,2}

¹Институт математики и математического моделирования МОН РК, Алматы, Казахстан;

²Казахский национальный университет имени аль-Фараби, Алматы, Казахстан

ТЕНЗОР ГРИНА УРАВНЕНИЙ ДВИЖЕНИЯ ДВУХКОМПОНЕНТНОЙ СРЕДЫ М. БИО ПРИ СТАЦИОНАРНЫХ КОЛЕБАНИЯХ

Аннотация. Рассматривается двухкомпонентная среда Био, содержащая твердую упругую компоненту и идеальную жидкую. Движение такой среды описывается связанной системой дифференциальных уравнений в частных производных второго порядка гиперболического типа для перемещений твердой и жидкой компоненты. Исследуются процессы распространения волн в среде Био, порождаемые действующими периодическими силами различного типа.

Строятся стационарные решения этой системы, описывающие гармонические колебания в пространствах размерности $N = 1, 2, 3$. В этом случае система уравнений для комплексных амплитуд колебаний является эллиптической. На основе преобразования Фурье обобщенных функций и его свойств построено фундаментальное решение уравнений колебаний среды Био - тензор Грина, который описывает процесс распространения гармонических по времени волн фиксированной частоты при действии сосредоточенных в начале координат силовых источников, описываемых сингулярной дельта-функцией.

На его основе построены обобщенные решения этих уравнений при действии разнообразных источников периодических возмущений, которые описываются как регулярными, так и сингулярными обобщенными функциями. Полученные решения можно использовать для исследования волновых процессов в газо- и жидконасыщенных пористых средах.

Ключевые слова: среда Био, твердая и жидкая компоненты, фундаментальное решение, обобщенное прямое и обратное преобразование Фурье, регуляризация.

Information about authors:

Alexeyeva L.A. - Doctor of physical-mathematical science, professor, Chief Researcher, Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan, E-mail: alexeeva@math.kz, <https://orcid.org/0000-0002-7131-4635>

Kurmanov E.B. – junior researcher, Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan, PhD student at Faculty of Physics and Technology, Al-Farabi Kazakh National University. E-mail: kurmanovergali@gmail.com, <https://orcid.org/0000-0003-3695-0166>

REFERENCES

- [1] Biot M.A. Theory of propagation of elastic waves in a fluid-saturated porous solid. 1. Low frequency range. J. Acoust. Soc. Amer. V.28. N 2. 1956. P.168-178.
- [2] Biot M.A. Theory of propagation of elastic waves in a fluid-saturated porous solid. 2. Higher frequency range. J. Acoust. Soc. Amer., V.28, 2, 1956, P.178-191.
- [3] Biot M.A. Mechanics of deformation and propagation of acoustic waves in a porous medium // - Mechanics. Sat 1963. № 6 (28). P.103-104.
- [4] Nikolayevsky V.N. Mechanics of porous and fractured media // M: Nauka, 1984. P. 232.
- [5] Horoshun L.P. To the theory of saturated porous media. // PM. 1976. V. 12, № 12.
- [6] Rakhmatullin Kh.A., Saatov Ya.U., Filippov I.G. and Artykov T.U. Waves in two-component media Science, Tashkent: Nauka, 1974. P. 266.
- [7] Saatov.Ya.U. Plane problems of mechanics of elastic-porous media. Tashkent: «Nauka», 1972. C. 228-239.
- [8] Yerzhanov Zh.S., Aitaliev Sh.M. and Alexeyeva L.A., Dynamics of tunnels and underground pipelines. Alma-Ata, Science Kaz. SSR, 1989. P. 240.
- [9] Alexeyeva L.A., V.V. Shershnev. Fundamental solutions of the equations of motion of the Biot's medium // Reports of NAS RK. 1994. № 1. P. 3–6.
- [10] Vladimirov V.S. Generalized functions in mathematical physics. M.: Mir. 1978.
- [11] Alexeyeva L.A., Kurmanov E.B. “Fundamental and generalized solutions of the two-component medium M. Bio 1. The Fourier transform of fundamental solutions and their regularization”, Math. J., 17, No. 2, 2017, 13–30.
- [12] Alexeyeva L.A., Kurmanov E. B. Fourier transform of fundamental solution for the motion equations of two-component Biot's media // AIP Conference Proceedings, 1880:(1) <https://doi.org/10.1063/1.5000675> (2017).
- [13] Alexeyeva L.A. Stationary boundary value problems of the dynamics of thermoelastic rods News NAS RK, series of physical and mathematical, 2014. No. 3. P.144-152.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы: *М. С. Ахметова, Г. Б. Халидуллаева, Д. С. Аленов*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 04.02.2020.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.
4,6 п.л. Тираж 300. Заказ 1.

Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19