

ISSN 2518-1726 (Online),  
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ  
әл-Фараби атындағы Қазақ ұлттық университетінің

# Х А Б А Р Л А Р Ы

---

---

## ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН  
Қазақстан Республикасының  
Ғылым Академиясының  
Әл-Фараби атындағы  
Қазақ ұлттық университетінің

## NEWS

OF THE NATIONAL ACADEMY OF SCIENCES  
OF THE REPUBLIC OF KAZAKHSTAN  
Al-Farabi Kazakh  
National University

**SERIES**  
**PHYSICO-MATHEMATICAL**

**2(330)**

**MARCH – APRIL 2020**

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р ы  
ф.-м.ғ.д., проф., ҚР ҰҒА академигі  
**Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

**Жұмаділдаев А.С.** проф., академик (Қазақстан)  
**Кальменов Т.Ш.** проф., академик (Қазақстан)  
**Жантаев Ж.Ш.** проф., корр.-мүшесі (Қазақстан)  
**Өмірбаев У.У.** проф., корр.-мүшесі (Қазақстан)  
**Жүсіпов М.А.** проф. (Қазақстан)  
**Жұмабаев Д.С.** проф. (Қазақстан)  
**Асанова А.Т.** проф. (Қазақстан)  
**Бошкаев К.А.** PhD докторы (Қазақстан)  
**Сұраған Д.** корр.-мүшесі (Қазақстан)  
**Quevedo Hernando** проф. (Мексика),  
**Джунушалиев В.Д.** проф. (Қырғыстан)  
**Вишневский И.Н.** проф., академик (Украина)  
**Ковалев А.М.** проф., академик (Украина)  
**Михалевич А.А.** проф., академик (Белорусь)  
**Пашаев А.** проф., академик (Әзірбайжан)  
**Такибаев Н.Ж.** проф., академик (Қазақстан), бас ред. орынбасары  
**Тигиняну И.** проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде  
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19; 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2020

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Главный редактор  
д.ф.-м.н., проф. академик НАН РК  
**Г.М. Мутанов**

Редакционная коллегия:

**Джумадильдаев А.С.** проф., академик (Казахстан)  
**Кальменов Т.Ш.** проф., академик (Казахстан)  
**Жангаев Ж.Ш.** проф., чл.-корр. (Казахстан)  
**Умирбаев У.У.** проф., чл.-корр. (Казахстан)  
**Жусупов М.А.** проф. (Казахстан)  
**Джумабаев Д.С.** проф. (Казахстан)  
**Асанова А.Т.** проф. (Казахстан)  
**Бошкаев К.А.** доктор PhD (Казахстан)  
**Сураган Д.** чл.-корр. (Казахстан)  
**Quevedo Hernando** проф. (Мексика),  
**Джунушалиев В.Д.** проф. (Кыргызстан)  
**Вишневский И.Н.** проф., академик (Украина)  
**Ковалев А.М.** проф., академик (Украина)  
**Михалевич А.А.** проф., академик (Беларусь)  
**Пашаев А.** проф., академик (Азербайджан)  
**Такибаев Н.Ж.** проф., академик (Казахстан), зам. гл. ред.  
**Тигиняну И.** проф., академик (Молдова)

**«Известия НАН РК. Серия физика-математическая».**

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов  
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219, 220; тел.: 272-13-19; 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© Национальная академия наук Республики Казахстан, 2020

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

Editor in chief

doctor of physics and mathematics, professor, academician of NAS RK

**G.M. Mutanov**

Editorial board:

**Dzhumadildayev A.S.** prof., academician (Kazakhstan)  
**Kalmenov T.Sh.** prof., academician (Kazakhstan)  
**Zhantayev Zh.Sh.** prof., corr. member. (Kazakhstan)  
**Umirbayev U.U.** prof., corr. member. (Kazakhstan)  
**Zhusupov M.A.** prof. (Kazakhstan)  
**Dzhumabayev D.S.** prof. (Kazakhstan)  
**Asanova A.T.** prof. (Kazakhstan)  
**Boshkayev K.A.** PhD (Kazakhstan)  
**Suragan D.** corr. member. (Kazakhstan)  
**Quevedo Hernando** prof. (Mexico),  
**Dzhunushaliyev V.D.** prof. (Kyrgyzstan)  
**Vishnevskiy I.N.** prof., academician (Ukraine)  
**Kovalev A.M.** prof., academician (Ukraine)  
**Mikhalevich A.A.** prof., academician (Belarus)  
**Pashayev A.** prof., academician (Azerbaijan)  
**Takibayev N.Zh.** prof., academician (Kazakhstan), deputy editor in chief.  
**Tiginyanu I.** prof., academician (Moldova)

**News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.**

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006.

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19; 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

---

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN  
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2020.2518-1726.23>

Volume 2, Number 330 (2020), 120 – 126

D.N. Nurgabyl<sup>1,2</sup>, T.M. Seitova<sup>2</sup>

<sup>1</sup>Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan;  
<sup>2</sup>Zhetysay State University named after I.Zhansugurov, Taldykorgan, Kazakhstan.  
E-mail: [kebek.kz@mail.ru](mailto:kebek.kz@mail.ru), [terasytoltganai@mail.ru](mailto:terasytoltganai@mail.ru)

INTRODUCING METHOD OF GENERALIZED DERIVATIVE  
CONCEPT IN MATHEMATICS

**Abstract.** In this paper, we consider a technique for introducing the concept of a generalized derivative function of one variable. Some assertions of combinatorics are given and proved; the concept of the derivative of the natural order is introduced using the limit of the sequence. Using the above statements, the concept of a fractional derivative is introduced. The basic properties of the fractional derivative are formulated and proved. Examples are given

**Key words:** mathematical analysis, derivative, combination, limit transition, fractional derivative.

**Introduction**

Mathematical analysis is a field of mathematics related to the concepts of function, derivative and integral.

The great English physicist, astronomer and mathematician Isaac Newton and the German mathematician and philosopher Gottfried Leibniz completed the construction of differential and integral calculi by the end of the 17th century. The discovery of differential and integral calculus was the beginning of a period of rapid development of mathematics.

Mathematics continues to develop rapidly. Various generalizations of the concepts of function, derivatives, and integral have a particular interest. For example, mathematicians, including Leibniz, Euler, Liouville, and Riemann, dealt with generalization of the concept of a derivative. Generalized functions and their derivatives find various applications in real processes of the economy and production [1-4].

Our goal is to develop a methodology for introducing various definitions of a generalized derivative function of one variable.

To achieve this goal, we first give some statements of combinatorics, introduce the concept of a derivative of the natural order using the limit of the sequence. Using these statements, we introduce the concept of a fractional derivative

**1. Auxiliary definitions and formulas**

As it is known [5], the number of combinations of  $n$  elements by  $k$  is equal

$$C_n^k = \frac{n!}{k!(n-k)!}, \quad n \geq k. \tag{1}$$

Here it is convenient to assume that  $0! = 1$ . The following well-known equalities are justly:

$$C_n^k + C_n^{k+1} = C_{n+1}^{k+1}, \tag{2}$$

$$C_n^0 \cdot C_m^k + C_n^1 \cdot C_m^{k-1} + \dots + C_n^{k-1} \cdot C_m^1 + C_n^k \cdot C_m^0 = C_{n+m}^k, \quad m, n \geq k. \tag{3}$$

In particular, in  $n = m$  formula (3) takes the form:

$$\sum_{i=0}^k C_n^i C_n^{k-i} = C_{2n}^k, \quad n > k \tag{4}$$

Obviously, that the formula (1) one can be written in the form of

$$C_n^k = \frac{n(n-1)\dots(n-k+1)}{k!}. \quad (5)$$

Now, we will consider a generalization of formulas (2), (3), (5). We notice that the right-hand side of formula (5) is defined for any real values  $n$ . Thus, by definition, we assume that

$$C_\tau^k = \frac{(\tau-0)(\tau-1)\dots(\tau-k+1)}{k!}, \quad (6)$$

where  $\tau$  is a real number. For example:

$$C_{-2}^k = \frac{(-2-0)(-2-1)(-2-2)\dots(-2-(k-1))}{k!} = (-1)^k (k+1), \quad C_{-1}^k = (-1)^k$$

*Lemma 1.* For any real values of  $\tau$ , the equality

$$C_\tau^k + C_{\tau+1}^k = C_{\tau+1}^{k+1} \quad (7)$$

*Proof:* Using (4), we have

$$\begin{aligned} C_\tau^k + C_{\tau+1}^{k+1} &= \frac{\tau(\tau-1)\dots(\tau-(k-1))}{k!} + \frac{\tau(\tau-1)\dots(\tau-(k-1))(\tau+1)}{(k+1)!} = \\ &= \frac{\tau(\tau-1)\dots(\tau-(k-1))}{k!} \cdot \left(1 + \frac{\tau-k}{k+1}\right) = \frac{\tau(\tau-1)\dots(\tau-(k-1))(\tau+1)}{(k+1)!} = \\ &= \frac{(\tau+1-0)(\tau+1-1)(\tau+1-2)\dots(\tau+1-k)}{(k+1)!} = C_{\tau+1}^{k+1}. \end{aligned}$$

Lemma 1 is proved.

*Lemma 2.* For any real values of  $\tau$ , the equality

$$C_\tau^0 C_\tau^k + C_\tau^1 C_\tau^{k-1} + \dots + C_\tau^{k-1} C_\tau^1 + C_\tau^k C_\tau^0 = C_{2\tau}^k \quad (8)$$

*Proof:* We use the following statement [6].

If the polynomials  $P(x)$  and  $Q(x)$ , whose degrees do not exceed  $n$ , have equal values for more than  $n$  different of values unknowns, then  $P(x) = Q(x)$ .

We write the equalities  $P(\tau) = \sum_{i=0}^k C_\tau^i C_\tau^{k-i}$ ,  $Q(\tau) = C_{2\tau}^k$ , where  $P(\tau)$  and  $Q(\tau)$  are polynomials of degree  $k$ .

By virtue of formula (4), the polynomials  $P(\tau)$  and  $Q(\tau)$  have equal values at  $\tau = n > k$ . Then, by based on the above statement, it is easy to verify the validity of formula (8) for any real values  $\tau$ . Lemma 2 is proved.

## 2. About a definition of a natural order derivative

For a function of one variable, we define a derivative of the natural order in a slightly different way. Let the function  $f(x)$  be defined and continuous on the interval  $[a, b]$ ,  $a < 0$ ,  $b > 0$ , at that  $f(x) = 0$  for  $x < 0$ , where  $x$  is a fixed point of this interval.

We divide the segment  $[0, x]$  into equal  $n$  parts by points  $0, \frac{x}{n}, \frac{2x}{n}, \dots, \frac{nx}{n}$ . We define the increment of the function  $y = f(x)$  at the point  $x$  in the form  $\Delta y = f(x) - f(x - \frac{x}{n})$ . As it is known, if the ratio

$$\frac{\Delta y}{\frac{x}{n}} = \frac{f(x) - f\left(x - \frac{x}{n}\right)}{\frac{x}{n}}$$

has a limit is at  $n \rightarrow \infty$ , then this limit is called the derivative of the function  $f(x)$  at the point  $x$  and is denoted by:

$$f'(x) = \lim_{n \rightarrow \infty} \frac{f(x) - f\left(x - \frac{x}{n}\right)}{\frac{x}{n}}.$$

Now we define the increment  $\Delta^2 y$ :

$$\begin{aligned} \Delta^2 y &= f(x) - f\left(x - \frac{x}{n}\right) - \left[ f\left(x - \frac{x}{n}\right) - f\left(x - \frac{2x}{n}\right) \right] = \\ &= f(x) - 2f\left(x - \frac{x}{n}\right) + f\left(x - \frac{2x}{n}\right) = C_2^0 f(x) + C_2^1 f\left(x - \frac{x}{n}\right) + C_2^2 f\left(x - \frac{2x}{n}\right) = \\ &= \sum_{k=0}^2 (-1)^k \cdot C_2^k f\left(x - \frac{kx}{n}\right), \end{aligned}$$

etc. continuing this process, we obtain:

$$\Delta^m y = \sum_{k=0}^m (-1)^k C_m^k f\left(x - \frac{kx}{n}\right). \quad (9)$$

Let  $m$  be a fixed positive integer. We choose a positive integer  $n$  so that  $n > m$ . Then the formula (9) will take the form:

$$\Delta^m y = \sum_{k=1}^n (-1)^k C_m^k \cdot f\left(x - \frac{kx}{n}\right), \quad (10)$$

as  $C_m^k = 0$ ;  $k > m$ . Using (10), we obtain

$$f^{(m)}(x) = \lim_{n \rightarrow \infty} \frac{\Delta^m y}{\left(\frac{x}{n}\right)^m} = \lim_{n \rightarrow \infty} \left(\frac{x}{n}\right)^{-m} \sum_{k=0}^n (-1)^k C_m^k \cdot f\left(x - \frac{kx}{n}\right). \quad (11)$$

As an example, we consider the function  $f(x) = x^2$ :

$$(x^2)'' = \lim_{n \rightarrow \infty} \left(\frac{x}{n}\right)^{-2} \left[ x^2 - 2\left(x - \frac{x}{n}\right)^2 + \left(x - \frac{2x}{n}\right)^2 \right] = \lim_{n \rightarrow \infty} \frac{2 \frac{x^2}{n^2}}{\frac{x^2}{n^2}} = 2, \text{ где } C_2^k = 0, k = \overline{3, n}.$$

### 3. Definition of a fractional order derivative. Properties

Based on the fact that the expression  $C_\tau^k$  according to (6) is defined for any real values of  $\tau$ , then the right-hand side of formula (10) is determined for any real values of  $\tau$ .

Now, using [7], for any real number  $\tau$  we define the derivative of the  $\tau$  th order:

$$f^{(\tau)}(x) = \lim_{n \rightarrow \infty} \frac{1}{\left(\frac{x}{n}\right)^\tau} \sum_{k=0}^n (-1)^k c_\tau^k f\left(x - \frac{kx}{n}\right). \tag{12}$$

Since any real number is an infinite decimal fraction, we call the derivative (14) fractional.

For example, given that  $C_{-1}^k = (-1)^k$ , from (12) the formula we find:

$$f^{(-1)}(x) = \lim_{n \rightarrow \infty} \frac{1}{\left(\frac{x}{n}\right)^{-1}} \sum_{k=0}^n f\left(x - \frac{kx}{n}\right) = \lim_{n \rightarrow \infty} \sum_{k=0}^n f\left(x - \frac{kx}{n}\right) \cdot \left(\frac{x}{n}\right) \tag{13}$$

Example: We define the (-1)st order derivative for function  $f(x) = x$ . From (13) we get:

$$\begin{aligned} f^{(-1)}(x) &= \lim_{n \rightarrow \infty} \frac{x}{n} \sum_{k=0}^n \left(x - \frac{kx}{n}\right) = \lim_{n \rightarrow \infty} \frac{x}{n} \left[ (x-0) + \left(x - \frac{x}{n}\right) + \left(x - \frac{2x}{n}\right) + \dots + \left(x - \frac{kx}{n}\right) \right] = \\ &= \lim_{n \rightarrow \infty} \frac{x}{n} \left[ (x-1)x - \frac{x}{n} \cdot \frac{(n+1)n}{2} \right] = \lim_{n \rightarrow \infty} \left[ \frac{x^2(n+1)}{n} - \frac{x^2(n+1)n}{2n^2} \right] = x^2 - \frac{x^2}{2} = \frac{x^2}{2}. \end{aligned}$$

In this way,  $(x)^{-1} = \frac{x^2}{2}$ .

Now we give the main properties of the  $\tau$  th derivative for any real of values of  $\tau$ .

Theorem 1. Right

$$\begin{aligned} (\alpha \cdot f(x) + \beta \cdot g(x))^{(\tau)} &= \alpha \cdot f^{(\tau)}(x) + \beta \cdot g^{(\tau)}(x), \\ \left[ f^{(\tau_1)}(x) \right]^{(\tau_2)} &= f^{(\tau_1 + \tau_2)}(x). \end{aligned} \tag{14}$$

To establish relations (14), it suffices to use representation (12) of the derivative of an arbitrary function.

From formula (15), we note that the sum  $\sum_{k=0}^m f\left(x - \frac{kx}{n}\right) \cdot \frac{x}{n}$  represents the integral sum of the function  $f(x)$  for a given partition  $\left\{ \frac{kx}{n} \right\}$ ,  $k = \overline{1, n}$  of the segment  $[0, x]$ ,  $x \in [0, x]$ .

Since the function  $f(x)$ , being continuous on the segment  $[0, x]$ , is integrable on  $[0, x]$ , therefore, the limit (13) gives us a definite integral  $\lim_{n \rightarrow \infty} \sum_{k=0}^n f\left(x - \frac{kx}{n}\right) \cdot \frac{x}{n} = \int_0^x f(t) dt$  and thus, we obtain

$$f^{(-1)}(x) = \int_0^x f(t) dt. \tag{15}$$

Theorem 2. For the natural value m, the derivative of the (-m)th order function is determined by the formula:

$$f^{(-m)}(x) = \frac{1}{(m-1)!} \int_0^x f(t)(x-t)^{m-1} dt. \tag{16}$$



Proof: By hypothesis, the function  $f(x)$  is continuous at any point  $x \in [a, b]$ . Therefore, the function  $f^{(-1)}(x) = g(x) = \int_0^x f(t)dt$  is also continuous in the same  $x \in [a, b]$ . Therefore, for the point  $x \in [0, x]$  we can find

$$f^{(-2)}(x) = \int_0^x f(t)(x-t)dt.$$

Next, by induction, we establish that

$$f^{(-k)}(x) = \frac{1}{(k-1)!} \int_0^x f(t)(x-t)^{k-1} dt. \quad (17)$$

Theorem 2 is proved.

### Conclusions

Thus, in this paper, we propose a technique for introducing the concept of a fractional derivative. Using the limit of the sequence, the notion of a derivative of the natural order is introduced, the definition of a fractional derivative is given for any real values of  $x$ . The basic properties of a fractional derivative are proved. Examples are given.

Practice has shown that this approach of introducing the concept of a generalized derivative contributes to the effective assimilation by students of various definitions of generalized functions.

Д.Н. Нұрғабыл<sup>1,2</sup>, Т.М. Сеитова<sup>2</sup>

<sup>1</sup> Математика және математикалық модельдеу институты, Алматы, Қазақстан;

<sup>2</sup> І. Жансүгіров атындағы Жетісу мемлекеттік университеті, Талдықорған, Қазақстан

### МАТЕМАТИКА КУРСЫНДА ЖАЛПЫЛАНҒАН ТУЫНДЫ ҰҒЫМЫН БЕРУ ӘДІСТЕМЕСІ

Қазіргі математикада функция, туынды және интеграл ұғымдарының әртүрлі жалпылама тұжырымдары ерекше қызығушылық тудырады. Мысалы, туынды ұғымның жалпылауымен математиктер, оның ішінде Лейбниц, Эйлер, Лиувиль және Риман айналысқан. Жалпыланған функциялар мен олардың туындылары экономика мен өндірістің нақты процестерінде әртүрлі қолданыстарын табууда. Бұл жұмыстың мақсаты бір айнымалы функцияның жалпыланған туындысының әр түрлі анықтамаларын енгізу әдістемесін жасау болып табылады. Аталған мақсатқа жету үшін бұл мақалада алдымен комбинаториканың тұжырымдарын, тізбек шегі ұғымын пайдалана отырып, натурал ретті туынды ұғымы енгізілген. Осы тұжырымдарды қолдана отырып, бөлшек туынды деген ұғым енгізіледі.

Комбинаторикада  $n$  элементтен  $k$  бойынша алынған терулер саны

$$C_n^k = \frac{n(n-1)\dots(n-k+1)}{k!} \quad (1)$$

формуласымен анықталатыны белгілі. (1) формуланың оң жағы кезкелген  $n = \tau$  нақты саны үшін де анықталатынын байқаймыз. Онда анықтама бойынша

$$C_\tau^k = \frac{(\tau-0)(\tau-1)\dots(\tau-k+1)}{k!}. \quad (2)$$

формуласын қабылдауымызға болады. Осы формуланы пайдалана отырып бір айнымалы функцияның натурал ретті туындысын сәл басқаша енгізілік.  $f(x)$  функциясы  $[a, b]$ ,  $a < 0$ ,  $b > 0$  аралығында анықталсын және осы интервалдың кезкелген нақты  $x$  нүктесі үшін  $f(x) = 0$ , мұнда  $x < 0$ .  $0, \frac{x}{n}, \frac{2x}{n}, \dots, \frac{nx}{n}$

нүктелерімен  $[0, x]$  кесіндісін  $n$  бірдей бөліктерге бөлелік. Сонда

$$\frac{\Delta y}{\frac{x}{n}} = \frac{f(x) - f(x - \frac{x}{n})}{\frac{x}{n}}$$

===== 124 =====

катысының  $n \rightarrow \infty$  -да шегі бар болса, онда бұл шекті  $f(x)$  функциясының  $x$  нүктесіндегі туындысы деп атаймыз. Сонымен:

$$f'(x) = \lim_{n \rightarrow \infty} \frac{f(x) - f\left(x - \frac{x}{n}\right)}{\frac{x}{n}}; \quad f^{(m)}(x) = \lim_{n \rightarrow \infty} \frac{\Delta^m y}{\left(\frac{x}{n}\right)^m} = \lim_{n \rightarrow \infty} \left(\frac{x}{n}\right)^{-m} \sum_{k=0}^n (-1)^k C_m^k \cdot f\left(x - \frac{kx}{n}\right). \quad (3)$$

Сонда (2) формулаға сәйкес  $C_{\tau}^k$  өрнегі кезкелген  $\tau$  нақты саны үшін анықталатын болғандықтан (3) формуланың оң жақтары кезкелген  $\tau$  нақты саны үшін де анықталатын болады. Олай болса, кезкелген  $\tau$  нақты саны үшін  $f(x)$  функциясының  $x$  нүктесіндегі  $\tau$ -ші ретті туындысы

$$f^{(\tau)}(x) = \lim_{n \rightarrow \infty} \frac{1}{\left(\frac{x}{n}\right)^{\tau}} \sum_{k=0}^n (-1)^k c_{\tau}^k f\left(x - \frac{kx}{n}\right). \quad (4)$$

Формуласымен анықталады. Ал кезкелген нақты сан шексіз ондық бөлшек болғандықтан, (4) –ші туындыны бөлшек ретті туынды деп атаймыз.

Теорема.  $m$  натурал саны үшін  $f(x)$  функциясының  $(-m)$ -ші ретті туындысы:

$$f^{(-m)}(x) = \frac{1}{(m-1)!} \int_0^x f(t)(x-t)^{m-1} dt.$$

формуласымен анықталады.

**Түйін сөздер:** математикалық анализ, туынды, теру, шекке көшу, бөлшек ретті туынды

Д.Н. Нургабыл<sup>1,2</sup>, Т.М. Сеитова<sup>2</sup>

<sup>1</sup>Институт математики и математического моделирования, Алматы, Казахстан;

<sup>2</sup>Жетысуский государственный университет им. И. Жансугурова, Талдыкорган, Казахстан

### МЕТОДИКА ВВЕДЕНИЯ ПОНЯТИЯ ОБОБЩЕННОЙ ПРОИЗВОДНОЙ В КУРСЕ МАТЕМАТИКИ

**Аннотация.** В современной математике особый интерес представляют различные обобщения понятий функции, производных и интеграла. Например, вопросами об обобщении понятия производной занимались математики, в том числе Лейбниц, Эйлер, Лиувилл и Риман. Обобщенные функции и их производные находят различные применения в реальных процессах экономики и производства.

Цель настоящей работы – разработка методики введения различных определений обобщенной производной функции одной переменной.

Для достижения этой цели вначале приведены некоторые утверждения комбинаторики, введены понятие производной натурального порядка с помощью предела последовательности. Используя эти утверждения, введены понятие производной дробного порядка.

В комбинаторике число сочетаний из  $n$  элементов по  $k$  определяется по формуле

$$C_n^k = \frac{n(n-1)\dots(n-k+1)}{k!}. \quad (1)$$

Заметим, что правая часть формулы (1) определена при любых вещественных значениях  $n = \tau$ . Тогда по определению можем положить, что

$$C_{\tau}^k = \frac{(\tau-0)(\tau-1)\dots(\tau-k+1)}{k!}. \quad (2)$$

Используя эту формулу определим для функции одной переменной производную натурального порядка несколько иным образом.

Пусть функция  $f(x)$  определена и непрерывна на промежутке  $[a, b]$ ,  $a < 0$ ,  $b > 0$ , причем  $f(x) = 0$  при  $x < 0$ , где  $x$ -фиксированная точка этого интервала.

Отрезок  $[0, x]$  разобьем на  $n$  равных частей точками  $0, \frac{x}{n}, \frac{2x}{n}, \dots, \frac{nx}{n}$ . Тогда, если отношение

$$\frac{\Delta y}{\frac{x}{n}} = \frac{f(x) - f(x - \frac{x}{n})}{\frac{x}{n}}$$

имеет предел при  $n \rightarrow \infty$ , то этот предел называется производной функции  $f(x)$  в точке  $x$  и обозначается:

$$f'(x) = \lim_{n \rightarrow \infty} \frac{f(x) - f(x - \frac{x}{n})}{\frac{x}{n}}, \quad f^{(m)}(x) = \lim_{n \rightarrow \infty} \frac{\Delta^m y}{\left(\frac{x}{n}\right)^m} = \lim_{n \rightarrow \infty} \left(\frac{x}{n}\right)^{-m} \sum_{k=0}^n (-1)^k C_m^k \cdot f\left(x - \frac{kx}{n}\right) \quad (3)$$

На основании того, что выражение  $C_{\tau}^k$  согласно (2) определено при любых вещественных значениях  $\tau$ , то правая часть формулы (3) определяется при любых вещественных значениях  $\tau$ .

Следовательно, для любого вещественного числа  $\tau$  производная  $\tau$ -го порядка функции  $f(x)$  в точке  $x$  определяется в виде:

$$f^{(\tau)}(x) = \lim_{n \rightarrow \infty} \frac{1}{\left(\frac{x}{n}\right)^{\tau}} \sum_{k=0}^n (-1)^k c_{\tau}^k f\left(x - \frac{kx}{n}\right). \quad (4)$$

Так как любое вещественное число это бесконечная десятичная дробь, то производную (4) назовем дробным.

Теорема. Для натурального значения  $m$  производная функции  $f(x)$   $(-m)$ -го порядка определяется по формуле:

$$f^{(-m)}(x) = \frac{1}{(m-1)!} \int_0^x f(t)(x-t)^{m-1} dt.$$

**Ключевые слова:** математический анализ, производная, сочетание, предельный переход, дробная производная.

#### Information about the authors:

Nurgabyl D. N. - doctor of physic-mathematical Sciences, Professor: Zhetysu State University named after I.Zhansugurov, Taldykorgan and Kazakh National Women's Teacher Training University, Almaty; Leading Researcher: Institute of Mathematics and Mathematical Modeling, Almaty, <https://orcid.org/0000-0002-8111-862X>

Seitova T.M. - Master student of Zhetysu State University named after I. Zhansugurov, Taldykorgan, <https://orcid.org/0000-0003-0175-7067>

#### REFERENCES

- [1] Serikbaev D., Tokmagambetov N. An inverse Problem for the Pseudo-Parabolic equation for a Sturm-Liouville operator // *News of the National Academy of sciences of the Republic of Kazakhstan*, Volume 4, Number 326(2019), pp. 122-128. ISSN 1991-346X. <https://doi.org/10.32014/2019.2518-1726.50>
- [2] Assanova A.T., Bakirova E.A., Kadirbayeva Zh.M. Numerical implementation of solving a Boundary Value Problem for a System of loaded differential equations with parameter. *News of the National Academy of sciences of the Republic of Kazakhstan*, Volume 3, Number 325 (2019), pp. 77-84. ISSN 1991-346X. <https://doi.org/10.32014/2019.2518-1726.27>
- [3] M.K. Dauylbayev, N. Atakhan, A.E. Mirzakulova, Asymptotic Expansion of solution of general BVP with initial jumps for higher-order Singularly Perturbed integro-differential Equation. *News of the National Academy of sciences of the Republic of Kazakhstan*, Volume 6, Number 322 (2018), pp. 28-36. ISSN 1991-346X. <https://doi.org/10.32014/2018.2518-1726.14>
- [4] M.N.Kalimoldayev, A.A.Abdildayeva, M.A.Akhmetzhanov, F.M.Galiyeva, Mathematical modeling of the Problem of Optimal Control of Electric Power Systems, *News of the National Academy of sciences of the Republic of Kazakhstan*, Volume 5, Number 321 (2018), pp. 62-67. ISSN 1991-346X. <https://doi.org/10.32014/2018.2518-1726.8>
- [5] Saveliev L.Ya. Combinatory and probability, Novosibirsk: Nauka, 1975 (in Russian).
- [6] Kurosh A.G. Course of Higher Algebra, M: Nauka, 1975(in Russian).
- [7] Rovinsky M. How to differentiate a function a fractional number of times // *Quantum*. M: Nauka, No. 8, 1986(in Russian).

## **Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

**ISSN 2518-1726 (Online), ISSN 1991-346X (Print)**

Редакторы: *М. С. Ахметова, Г. Б. Халидуллаева, Д. С. Аленов*  
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 05.04.2020.  
Формат 60x881/8. Бумага офсетная. Печать – ризограф.  
11 п.л. Тираж 300. Заказ 2.

---

*Национальная академия наук РК*  
*050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19*