

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
әл-Фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының Ғылым
Академиясының
Қазақ ұлттық университетінің
им. аль-Фараби

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-Farabi Kazakh
National University

SERIES
PHYSICO-MATHEMATICAL

2(330)

MARCH – APRIL 2020

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р ы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі
Ғ.М. Мұтанов

Р е д а к ц и я а л қ а с ы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев У.У. проф., корр.-мүшесі (Қазақстан)
Жүсіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Д. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қырғыстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорусь)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19; 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2020

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Главный редактор
д.ф.-м.н., проф. академик НАН РК
Г.М. Мутанов

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жангаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф., чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физика-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219, 220; тел.: 272-13-19; 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2020

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

Editor in chief

doctor of physics and mathematics, professor, academician of NAS RK

G.M. Mutanov

Editorial board:

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof., corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskiy I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006.

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19; 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2020.2518-1726.24>

Volume 2, Number 330 (2020), 127 – 132

УДК 550.383

A.F. Yakovets¹, A. Jahanshir², G.I. Gordienko¹, B.T. Zhumabayev¹, Yu.G. Litvinov¹

¹«Institute of Ionosphere» JSC «NCSRT», Almaty, Kazakhstan;

²Buein Zahra Technical University, Iran.

E-mail: artyak40@mail.ru, assiya173@mail.ru

DIURNAL DEPENDENCE OF THE FREQUENCY OF OCCURRENCE OF TRAVELING IONOSPHERIC DISTURBANCES OVER ALMATY

Abstract. We studied the diurnal dependence of the frequency of occurrence of large-scale traveling ionospheric disturbances (LSTID's) at mid-latitudes, which are a manifestation of atmospheric gravity waves (AGWs) generated in polar regions during geomagnetic disturbances. A significant amount of vertical sounding data of the ionosphere obtained at the Institute of the Ionosphere (Almaty 76 ° 55'E, 43 ° 15'N) in 2000-2007 was analyzed on a digital PARUS ionosonde connected to a computer designed to collect, store and process ionograms in digital form.

Further processing included the calculation of the altitude distributions of electron density

($N(h)$ profiles) by the Titterton method and deriving from them variations of a number of F -region parameters (electron density at fixed heights $N_h(t)$; density at the maximum of the layer $N_mF(t)$; the heights of the maximum of the layer $N_mF(t)$, etc.). The ionosonde provided a readout accuracy of

$h'(t) \sim 2.5$ km and a readout accuracy of $f_oF \sim 0.05$ MHz. It was shown that from session to session the number of waves observed during the night changed. In constructing the diurnal dependence of the frequency of occurrence of traveling ionospheric disturbances, all waves were taken into account.

A diurnal dependence of the frequency of occurrence of traveling ionospheric disturbances over Almaty was constructed from the measurement data. For this, a visual control of the behavior of a number of parameters of the F -region was carried out, identification of the LSTID's, determination of the time of their onset and duration.

It is shown that the predominant beginning of the development of the LSTID's is close to the moment of local midnight. 87% of traveling ionospheric disturbances were observed in the interval 20:00-04:00 LT. The distribution of the frequency of occurrence of the LSTID's in the time of day coincides with the dependence of the substorm frequency on world time, explained on the basis of diurnal variations in the angle of inclination of the Earth's magnetic axis to the Sun-Earth line. Ion drag and the dependence of the level of auroral activity, and, consequently, the intensity of the generation of LSTID's, on world time determines that the most favorable conditions for the distribution of LSTID's are created over Almaty at night.

Key words: ionosphere, vertical sounding, diurnal dependence of the frequency of occurrence of large-scale traveling ionospheric disturbances.

Introduction

Large-scale traveling ionospheric disturbances (LSTIDs) are caused by atmospheric gravity waves (AGWs), generated in the polar regions during geomagnetic storms [1], when the fast enhancement of auroral electrojets leads to the heating of the atmosphere. The process of rapid expansion and further compression of the atmosphere generates AGWs that propagate towards the equator and originates an LSTIDs on the way of their propagation. For many years, the propagation of AGWs in the neutral atmosphere and their ionospheric manifestation have been studied both experimentally and theoretically. The results of these studies are presented in the review works [1, 2]. The typical parameters of LSTIDs in the F region of the ionosphere are represented by the following values: the periods lie within the range from 40 min to 3 h; the horizontal wave lengths are 1000–3000 km; and the phase velocities are 400–1000 m/s. It is believed that generation and propagation of AGWs play an important role in the transfer of energy

from the magnetosphere to the low latitude ionosphere. Although the LSTIDs have been studied for several decades, some fundamental problems remain little studied. Such problems include the question of the diurnal dependence of the frequency of their appearance at mid-latitudes.

Description of the equipment, and observation results

Nighttime observations of LSTIDs in the F region of the ionosphere are carried out at the Institute of Ionosphere (Almaty 76°55' E, 43°15' N) with a PARUS digital ionosonde connected to a computer for the collection, storage, and processing of ionograms in digital form. The information required for calculating various parameters of LSTIDs is read from ionograms using the semiautomatic method. Ionospheric sounding is conducted every 5 min. Nighttime measurement sessions last ~8–12 h, depending on the season. Ionograms provide the values of virtual heights of radio signal reflection $h'(t)$ at several fixed operating frequencies of sounding and the values of critical frequencies f_oF . The further processing involves the estimation of the height distributions of electron densities ($N(h)$ profiles) by the Titheridge method [3] and obtaining the variations in several parameters of the F region based on them (the electron density at fixed heights $N_h(t)$; the density at layer maximum $N_mF(t)$; the heights of layer maximum $h_mF(t)$, etc.).

The ionosonde ensures the reading accuracies $h'(t) \sim 2.5$ km and $f_oF \sim 0.05$ MHz. Nighttime was chosen for observations because LSTIDs with big amplitudes of variations in ionospheric parameters at midlatitudes are usually observed at night [4]. For the period 2000–2007 we carried out 1166 night observations, while 581 nights were characterized by wave activity [5]. The variations in $N_h(t)$ at a series of heights made it possible to determine the form of a height profile of amplitudes $A(h)$ with the A_m maximum absolute amplitude. For analysis, we selected observation sessions that recorded LSTIDs with a relative amplitude (δh) exceeding 25% at a height corresponding to A_m . Here, $\delta h = A(h)/N(h)$, where $A(h)$ is the absolute amplitude of a wave at height h and $N(h)$ is the value of the undisturbed electron density at a given height.

Figure 1 shows a typical time behavior of a number of parameters after the onset of large magnetic disturbances. From the figure it follows that the beginning of the LSTIDs falls on the time interval 20:00–22:00, and the end on the interval 02:00–04:00. The lower curve corresponds to the height of the base of the layer ($h = 150$ km). The upper (bold) curve corresponds to the variations of $N_mF(t)$ at the maximum of the h_mF layer. Variations in the electron density shown in the figures demonstrate a feature characteristic of most of the sessions in which LSTIDs were observed. The peculiarity lies in the fact that the LSTIDs in the $N_mF(t)$ variations are manifested much weaker than in the $N_h(t)$ variations at fixed heights located below the height of the layer maximum. The reasons for this altitudinal dependence of the ionosphere response to AGW passage were considered in [6].

It should be noted that the number of waves observed during the night changed from session to session. In the example shown in figure 1, four waves are present. In constructing the diurnal dependence of the frequency of occurrence of traveling ionospheric disturbances, we took into account all the waves.

Earlier [7], a good correlation of LS TIDs with auroral substorms was proved; therefore, good similarity of their diurnal dependences should be expected. The long-term component in auroral disturbances, representing the dependence of the intensity and number of substorms on world time, was noted in [8, 9]. It became obvious that the beginning of the auroral substorms, as follows from the behavior of the auroral electrode jet index (AL -index), was the most frequent between 13:00 and 19:00 UT. More than 30% of the reported peaks in the AL -index accounted for a relatively narrow time domain of 13:00–16:00 UT [9]. The dependence of the substorm frequency on world time was explained on the basis of diurnal variations in the angle of inclination of the Earth's magnetic axis to the Sun-Earth line.

We analyzed the diurnal dependence of the frequency of occurrence of traveling ionospheric disturbances over Almaty according to measurements for 2000–2007. For this, a visual control of the behavior of a number of parameters of the F-region was carried out, identification of the LS TIDs, determination of their start time and duration. It is necessary to notice that high probabilities of the formation of nighttime enhancements in NmF2 and the passage of LS TIDs mean a high probability of their simultaneous presence over the observation point, leading to a necessity to distinguish these two events. We performed selection LS TIDs as made it in our work [10].




Figure 1 - Variations of the *F*-region parameters during the night with high magnetic activity (August 31 – September 1, 2005): top panel — electron density $N(t)$ at a series of heights with a distance between adjacent heights of 10 km; the second panel - the heights of the maximum of the *F*-region $h_m F$ and its base $h_{bot} F$; the third and fourth from the top of the panel are $h_m F$ and $h_{bot} F$ with an exclusive trend




Figure 2 - diurnal dependence of the frequency of occurrence of large-scale traveling ionospheric disturbances over Almaty

According to measurements for 2000-2007. we have constructed a diurnal dependence of the frequency of occurrence of traveling ionospheric disturbances (figure 2). From the figure it follows that the predominant beginning of the development of the LS TIDs is close to the moment of local midnight. 87% of the traveling ionospheric disturbances were observed in the interval 20:00-04:00 LT. Taking into account the difference between world time and local Almaty time is 5 hours, we find that the overwhelming majority of the LS TIDs was observed in the interval 15:00-23:00 UT.

Let us consider why the LS TIDs are local in time near the midnight. In [11], the response of the ionosphere to a large magnetic storm was studied using data from the global network of vertical sounding stations and the Cosmos-900 satellite. It has been shown that the ionospheric response, which is the LS TIDs with a quasiperiod of ~ 3-4 hours, is observed throughout the globe, while the amplitudes of the LS TIDs in the night observation sector are several times higher than the amplitudes observed in the illuminated half of the Earth.

The planetary nature of the propagation of large-scale LS TIDs was also studied in [7] for a period of high solar activity, where it was shown that the probability of observing LS TIDs in the Australian-Asian longitude sector is significantly higher than the probability for the European and American sectors. This longitudinal effect is explained on the basis of the mechanism of ion drag of AGW, which takes maximum values at the illuminated time of day, and the diurnal dependence of auroral activity with a maximum of 13–19 UT. While neutral particles move freely across the line of the geomagnetic field, ions rotate around the field line and have difficulty crossing the field lines. This difference between the mobility of neutral particles and the mobility of ions limits the movement of neutral particles in the AGW in the collision of neutral particles and ions, leading to the attenuation of the AGW. This mechanism is called the ion drag effect. The magnitude of ion drag depends on the frequency of collisions of neutrals with ions, which is linearly related to the densities of neutrals and ions. Since the density of ions in the daytime is much higher than the nighttime density, the AGW attenuation on the daytime side of the Earth noticeably exceeds the attenuation on the nighttime side.

Ion drag and the dependence of the level of auroral activity, and, consequently, on the intensity of the generation of LS TIDs, on world time determines that the most favorable conditions for the distribution of LS TIDs are created over Almaty at night.

Conclusion

On the basis of a graphical representation of the reaction of the parameters of the F2 layer in the 23rd cycle of solar activity to the passage of large-scale traveling ionospheric disturbances (LS TIDs), the diurnal dependence of the frequency of the appearance of LS TIDs over Almaty was studied. It is shown that the frequency has pronounced maxima in the interval 20: 00-04: 00 local time. The distribution of the frequency of occurrence of the LS TIDs in time of day coincides with the dependence of the substorm frequency on world time, which is explained on the basis of diurnal variations in the angle of inclination of the Earth's magnetic axis to the Sun-Earth line.

The work was carried out in accordance with RBP-008 "Development of space technologies for monitoring and forecasting natural resources, technogenic environmental changes, creation of space technology and ground-based space infrastructure, research of long-distance and near-space objects" under the theme "Creation of a system of diagnostics and forecast of space weather for analysis and forecasting of conditions of functioning of space vehicles, navigation and communication systems (2018-2020), registration number (RN) 0118PK00800.

А.Ф. Яковец¹, А. Жаханшир², Г.И. Гордиенко¹, Б.Т. Жумабаев¹, Ю.Г. Литвинов¹

¹ЕЖШС «Ионосфера институты» «ҰҒЗТО» АҚ, Алматы, Қазақстан;

²Буин Захра техникалық университеті, Иран

АЛМАТЫ ҚАЛАСЫНЫҢ ҮСТІНЕН ҚОЗҒАЛАТЫН ИОНОСФЕРАЛЫҚ АУЫТҚУЛАРДЫҢ ПАЙДА БОЛУ ЖИЛІГІНІҢ ТӘУЛІКТІК ТӘУЕЛДІЛІГІ

Аннотация. Орташа ендіктегі ірі масштабты қозғалатын ионосфералық ұйтқулардың (ІМҚИҰ) пайда болу жылігінің, геомагниттік ауытқулар кезінде полярлық аудандарда генерацияланатын атмосфералық гравитациялық толқындардың (АГТ) көріністері болып табылатын тәуліктік тәуелділігі зерттелді.

Ионосфераны тік зондтау деректерінің 2000-2007 жылдары алынған елеулі көлемі талданды. Ионосфера институтында (Алматы $76^{\circ}55' E$, $43^{\circ}15' N$) сандық ионозонд "ПАРУС" компьютермен ұштасқан, ионограммаларды сандық түрде жинауға, сақтауға және өңдеуге арналған. ІМҚИҰ әр түрлі параметрлерін есептеу үшін қажетті ақпарат ионограммен жартылай автоматты әдіспен есептелді. Ионосфераны зондтау әрбір 5 мин. жүргізілді. Түнгі өлшеу сеанстарының ұзындығы маусымға байланысты өзгерді және $\sim 8-12$ сағ құрады. Ионограммалардан маңызы бар қаланың қолданыстағы биіктерге көрсету $h'(t)$ радиосигналды бірқатар белгіленген жұмыс жиілік зондтау және маңызы бар сыни жиілік f_oF есептелінді. Одан әрі өңдеу Титеридж әдісімен электронды тығыздықтың $N(h)$ -профильдердің биіктік таралуын есептеуді және олардың негізінде F-аймақтың бірқатар параметрлерінің вариацияларын алуды ($N_h(t)$ тіркелген биіктіктердегі электрондық тығыздықты; $N_mF(t)$ қабатының максимумындағы тығыздықты; $N_mF(t)$ қабатының максимум биіктігін және т.б.) қамтиды. Ионозонд $h'(t) \sim 2.5$ км оқу дәлдігін және $f_oF \sim 0.05$ МГц оқу дәлдігін қамтамасыз етті.

Сеанстан сеансқа түнгі уақытта байқалатын толқындардың саны өзгергені көрсетілген.

Қозғалатын ионосфералық ауытқулардың пайда болу жиілігінің тәуліктік тәуелділігін құру кезінде барлық толқындар ескерілді. Өлшеу деректері бойынша Алматы үстінен қозғалатын ионосфералық ұйтқулардың пайда болу жиілігінің тәуліктік тәуелділігі салынды. Ол үшін F-облыстың бірқатар параметрлерінің мінез-құлқына визуалды бақылау, ІМҚИҰ сәйкестендіру, олардың басталу уақыты мен ұзақтығын анықтау жүзеге асырылды. ІМҚИҰ дамуының басым бастауы жергілікті түн жаруға жақын екендігі көрсетілген. 87% қозғалатын ионосфералық 20:00-04:00 LT аралығында қалыптан байқалды. ІМҚИҰ пайда болу жиілігінің тәулік уақытында таралуы жер магниттік осінің Күн-Жер сызығына көлбеу бұрышының тәуліктік вариациясы негізінде түсіндірілетін қосалқы ағын жиілігінің әлемдік уақыттан тәуелділігіне сәйкес келеді. Иондық тежелу және авроральді белсенділік деңгейінің тәуелділігі, демек, ІМҚИҰ генерациясының қарқындылығы әлемдік уақыттан бастап ІМҚИҰ тарату үшін аса қолайлы жағдайлар Алматының үстінен түнгі уақытта құрылатынын анықтайды.

А.Ф. Яковец¹, А. Жаханшир², Г.И. Гордиенко¹, Б.Т. Жумабаев¹, Ю.Г. Литвинов¹

¹ДТОО «Институт ионосферы» АО «НЦКИТ», Алматы, Казахстан;

²Буин Захра Технический Университет, Иран

СУТОЧНАЯ ЗАВИСИМОСТЬ ЧАСТОТЫ ПОЯВЛЕНИЯ ПЕРЕМЕЩАЮЩИХСЯ ИОНОСФЕРНЫХ ВОЗМУЩЕНИЙ НАД АЛМАТЫ

Аннотация. Изучена суточная зависимость частоты появления крупномасштабных перемещающихся ионосферных возмущений (КМ ПИВ) на средних широтах, являющихся проявлением атмосферных гравитационных волн (АГВ), генерируемых в полярных районах во время геомагнитных возмущений. Проанализирован значительный объем данных вертикального зондирования ионосферы, полученных в 2000-2007 гг. в Институте ионосферы (Алматы $76^{\circ}55'E$, $43^{\circ}15'N$) на цифровом ионозонде «ПАРУС», сопряженным с компьютером, предназначенным для сбора, хранения и обработки ионограмм в цифровом виде. Информация, необходимая для расчетов разнообразных параметров КМ ПИВ, считывалась с ионограмм полуавтоматическим методом. Зондирование ионосферы проводилось каждые 5 мин. Длина ночных сеансов измерений изменялась в зависимости от сезона и составляла $\sim 8-12$ ч. С ионограмм считывались значения действующих высот отражения $h'(t)$ радиосигнала на ряде фиксированных рабочих частот зондирования и значения критических частот f_oF . Дальнейшая обработка включала в себя расчет высотных распределений электронной плотности ($N(h)$ -профилей) методом Титериджа и получение на их основе вариаций ряда параметров F-области (электронной плотности на фиксированных высотах $N_h(t)$; плотности в максимуме слоя $N_mF(t)$; высоты максимума слоя $N_mF(t)$ и др.). Ионозонд обеспечивал точность считывания $h'(t) \sim 2.5$ км и точность считывания $f_oF \sim 0.05$ МГц. Показано, что от сеанса к сеансу менялось количество волн, наблюдаемых в течение ночи. При построении суточной зависимости частоты появления перемещающихся ионосферных возмущений учитывались все волны.

Была построена суточная зависимость частоты появления перемещающихся ионосферных возмущений над Алматы по данным измерений. Для этого осуществлялся визуальный контроль поведения ряда параметров F-области, идентификация КМПИВ, определение времени их начала и продолжительности.

Показано, что преобладающее начало развития КМПИВ близко к моменту местной полуночи. 87% перемещающихся ионосферных возмущений наблюдалось в интервале 20:00-04:00 LT. Распределение частоты появления КМПИВ во времени суток совпадает с зависимостью частоты суббурь от мирового времени, объясняемой на основе суточных вариаций угла наклона земной магнитной оси к линии Солнце-

Земля. Ионное торможение и зависимость уровня авроральной активности, а, следовательно, и интенсивности генерации КМ ПИВ, от мирового времени определяет то, что наиболее благоприятные условия для распространения КМ ПИВ создаются над Алматы в ночные часы.

Information about authors:

Yakovets Arthur Fedorovich, Ph.D. mat. Sciences, VNS, Institute of the Ionosphere, JSC "NTSKIT", Almaty, artyak40@mail.ru. <https://orcid.org/0000-0002-3374-9507>;

Arezu Jahanshir, PhD in Teoretical Physics and Mathematical, Assistant Professor & Faculty Member, Department of Physics & Engineering Sciences, Buein Zahra Technical University, Iran. jahanshir@bzte.ac.ir, <https://orcid.org/0000-0001-7923-2186>;

Gordienko Galina Ivanovna, Ph.D. mat. Sciences, ANS, Institute of the Ionosphere, JSC "NCSCIT", Almaty, ggordienko@mail.ru. ggordienko@mail.ru, <https://orcid.org/0000-0002-9337-206X>;

Zhumabayev Beibit Tenelovich, Ph.D. mat. Sciences, ANS, Institute of the Ionosphere, JSC "NCCIT", Almaty, beibit.zhu@mail.ru. beibit.zhu@mail.ru, <https://orcid.org/0000-0001-7882-0277>;

Litvinov Yuri Georgievich, Ph.D. mat. Sci., SSS, Institute of the Ionosphere, NCSCIT JSC, Almaty, yurii-litvinov@mail.ru. yurii-litvinov@mail.ru, <https://orcid.org/0000-0002-0600-0517>

REFERENCES

[1] Hunsucker R. D. Atmospheric gravity waves generated in the high-latitude ionosphere: A review // *Rev. Geophys.* V. 20. P. 293–315. 1982.

[2] Hocke K., Schlegel K. A review of atmospheric gravity waves and traveling ionospheric disturbances: 1982–1995 // *Ann. Geophysicae.* V. 14. P. 917–940. 1996.

[3] Titheridge J.E. Ionogram analysis with the generalized program Polan. Boulder. CO USA: National Geophys. Data Center. 189 p. 1985.

[4] Hajkowicz L.A. Global study of large scale travelling ionospheric disturbances (TIDs) following a step like onset of auroral substorm in both hemispheres // *Planet. Space Sci.* V. 38. P. 913–923. 1990.

[5] Yakovets A.F., Vodyannikov V.V., Andreev A.B., Gordienko G.I., and Litvinov Yu.G. Features of Statistical Distributions of Large-Scale Traveling Ionospheric Disturbances over Almaty. *Geomagnetism and Aeronomy*, 2011, Vol. 51, No. 5, pp. 640–645. (Translated).

[6] Yakovets A.F., Vodyannikov V.V., Gordienko G.I., Ashkaliev Ya.F., Litvinov Yu.G. and Akasov S.B. Response of the Nighttime Midlatitude Ionosphere to the Passage of an Atmospheric Gravity Wave. *Geomagnetism and Aeronomy*, 2008, Vol. 48, No. 4, pp. 511–517. (Translated).

[7] Hajkowicz, L. A. Ionospheric response to auroral substorms during sunspot maximum (1980–82) // *Ann. Geophysicae*, 13, 95–104, 1995.

[8] Hajkowicz, L. A. Universal time effect in the occurrence of large-scale ionospheric disturbances // *Planet. Space Sci.*, 40, 1093–1099, 1992.

[9] Hajkowicz, L. A. Equatorial ionospheric response to isolated auroral substorms over a solar cycle (1980–85): evidence of longitudinal anomaly // *Ann. Geophysicae*, 14, 906–916, 1996.

[10] Yakovets A.F., Gordienko G.I., Zhumabayev B.T., Litvinov Yu.G. Comparison of altitude profiles of amplitude of two types of F2-layer ionospheric disturbances // *News of the National Academy of Sciences of the Republic of Kazakhstan, Physico-Mathematical Series.* Volume 4, Number 320 (2018), 79 – 84.

[11] Karpachev A.T., Deminova G.F. Planetary picture of the effects of large-scale AGWs in the ionosphere during a storm on March 22, 1979 // *Geomagnetism and Aeronomy*, 2008, Vol. 44, No. 6, pp. 799-812. 2004. (Translated).

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы: *М. С. Ахметова, Г. Б. Халидуллаева, Д. С. Аленов*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 05.04.2020.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
11 п.л. Тираж 300. Заказ 2.

Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19