

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
әл-Фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
Әл-Фараби атындағы
Қазақ ұлттық университетінің

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-Farabi Kazakh
National University

SERIES
PHYSICO-MATHEMATICAL

3 (331)

MAY – JUNE 2020

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р ы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі
Ғ.М. Мұтанов

Р е д а к ц и я а л қ а с ы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев У.У. проф., корр.-мүшесі (Қазақстан)
Жүсіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Д. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қырғыстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорусь)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19; 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2020

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Главный редактор
д.ф.-м.н., проф. академик НАН РК
Г.М. Мутанов

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жангаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф., чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физика-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219, 220; тел.: 272-13-19; 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2020

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

Editor in chief

doctor of physics and mathematics, professor, academician of NAS RK

G.M. Mutanov

Editorial board:

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof., corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskiy I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006.

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19; 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2020.2518-1726.55>

Volume 3, Number 331 (2020), 209 – 216

УДК 550.383

G.I. Gordiyenko, A.F. Yakovets, Yu.G. Litvinov«Institute of Ionosphere» JSC «NCSRT», Almaty, Kazakhstan,
ggordiyenko@mail.ru, artyak40@mail.ru, yurii-litvinov@mail.ru**LONG - TERM VARIATIONS
IN THE F- AND D-REGIONS OF IONOSPHERE**

Abstract. The long-term changes in the near-noon, diurnal, and midnight critical frequencies of the ionosphere F2 layer (foF2) and near-noon minimum reflection frequencies (fmin) were studied using ground-based vertical radio sounding of the ionosphere at Alma-Ata station [43.25N, 76.92E] for the period 1957-2017. The data on solar and geomagnetic activity were used as factors affecting the state of the ionosphere.

It was taken into account that the minimum reflection frequency depends both on the absorption of the probe signal in the lower ionosphere, and on the noise level and technical characteristics of the ionosonde. Therefore, it seems problematic to estimate the absolute value of signal absorption in this way. However, as a qualitative characteristic, as an indicator of the absorption level of the probe signal, the parameter fmin can be used.

Arithmetic average values of median foF2 for near noon (10-14 LT), near midnight (23-01 LT) hours and average diurnal values for this period of ionosphere observation are considered as initial data. The variability of the lower ionosphere was studied using fmin data for daylight hours. The monthly average values of the solar radio emission flux F10.7 and the Ap index are considered as characteristics of solar and geomagnetic activity. The geomagnetic (Ap) indexes, the near-noon, near-midnight, diurnal averaged foF2, and near-noon fmin variations are found to be in strong dependence from the solar activity; all of them show a dominant pattern of variation with a period of ~34-36 years and linear negative trend. The correlation coefficient between the foF2 and F10.7 long-term variations is very high, up to 0.99 that permit us to believe that the solar activity can be considered to be the main driver for the long-term variations in the ionospheric F2-region.

The geomagnetic (Ap) index, the near-noon, near-midnight, daily averaged foF2, and near-noon fmin variations are found to be in strong dependence from solar activity; all of them show a dominant pattern of variation with a period of ~34-36 years and linear negative trend. The correlation coefficient between the foF2 and F10.7 long-term variations is very high, up to 0.99, that permits us to believe that the solar activity can be considered as a main driver for the long-term variations in the ionospheric F2-region.

The long-term course of fmin is similar to those found in F10.7, Ap, and foF2, i.e. the periodicity of 34-36 years is also evident in the fmin variations. However, in contrast to the F2-layer parameters, the fmin variation clearly demonstrates an upward (positive) linear trend that is opposite in sign to the trend found in F10.7. This means relatively high sensitivity of the fmin values to the solar activity changes and significant influence of other trend drivers on them, one of which is the possible impact of anthropogenic factors on the state of the lower ionosphere.

Key words: midlatitude ionosphere; upper and lower ionosphere; D-region, F2-region, long-term trends.

Introduction. In our previous work we investigated the long-term variations in the near-noon (10-14 LT) critical frequency of the ionospheric F2-layer (foF2) measured at the mid-latitude station Alma-Ata [43.25N, 76.92E] over the period from 1957 to 2012 [1]. The purpose of this paper is to provide further analyses of the F2-layer variations and study long-term trends (a long-term linear change) in the near-noon, near-midnight and daily averaged F2-layer critical frequencies measured at the Alma-Ata station in more extended time period, from 1957 to 2017 including the period of very deep minimum in solar activity observed in 2008-2009. In addition to the F2-layer parameters the minimum frequency of

reflection f_{min} is also used as a climatic characteristic of the upper atmosphere/low ionosphere (D region) to do the trend analyses in the ionospheric absorption. The f_{min} value depends, apart from the absorption of radio waves in the ionosphere, on technical characteristics of ionosondes that does not give us any opportunity to use this ionosphere parameter to determine the absolute values of the absorption of radio waves. However, as a qualitative characteristic, as an indicator of absorption of radio waves, the f_{min} value is widely used ([2-3] and references therein).

For the present trend analyses, monthly median foF2 and f_{min} values routinely measured at the Alma-Ata station have been used. The arithmetic means of the foF2 values at the near-noon (10-14 LT), near-midnight (23-01 LT) hours, and daily means foF2 are calculated for the analyses, the f_{min} values are taken only for the near-noon hours: foF2₁₀₋₁₄, foF2₂₃₋₀₁, foF2_{av}, and $f_{min_{11-13}}$ correspondingly. The monthly mean solar flux F10.7 and geomagnetic index Ap are also used as the indexes for solar and geomagnetic activities (available at <http://www.swpc.noaa.gov>) to illustrate their long-term variations.

Description of the data, and observation results *Trends in the ionospheric F-region.* As an example, median mean values foF2₁₀₋₁₄, foF2₂₃₋₀₁, foF2_{d,av} together with the corresponding data of solar and geomagnetic activity is shown in figure 1a-e. The dots represent observed data, the thick lines show the fitting curves. It should be also noted, the statistical processing of the ionospheric data series requires their continuity that is not always possible because of different reasons (ionosonde repair and other technical problems).

The regression dependences between the selected ionospheric parameters and F10.7 (figure 1f-h) have been analyzed to define the missing data. Thick lines represent the linear regression lines for these data; dashed lines correspond to the polynomial functions of the second degree that better fit the given F2-layer parameters ($R^2_{polynomial} > R^2_{linear}$) where R^2 is the coefficient of determination that provides a measure of how well the least-square curve fits the observational data, r is the correlation coefficient. Assuming the second-order polynomial dependence with F10.7 the missing foF2 values have been defined from the regression equations to fill available gaps in the data sets. Two things are evident in the figure, firstly the variations of all parameters are strictly modulated by cyclic 11-year variations of solar activity and secondly the foF2 data are much scattered relatively to the smoothed lines that can be attributed to various sources other than solar activity, including planetary waves and seasonal variations. As an example, the great part of seasonal variations is evident in Figure 1e where large deviations of the foF2₂₃₋₀₁ values from the smoothed line is caused by the fact that summer nighttime foF2 values are much higher than winter ones. Since this intraseasonal variability has to be taken into consideration in the trend analyses, annual averages of the parameters considered were calculated. Figure 2a-e demonstrates temporal variations of the calculated annual averages (denoted by symbol*) of the F10.7*, Ap* indexes, and the F2-layer parameters (foF2*₁₀₋₁₄, foF2*₂₃₋₀₁, and foF2*_{av}) for whole period of observations; variations of the ionospheric parameters with F10.7* are presented in Figure 2f-h.

The main feature of the variations is their similarity (figure 2a-e) and close connection between the ionospheric parameters and solar activity (figure 2f-h). The coefficients of determination R^2 (the coefficients R^2 and derived regression equations are shown in the figure fields) are found to be very high, from 0.97 to 0.99 it means that from 97% to 99% of the annual foF2 variations can be explained by their relationship with the 11-year cycle of solar activity.

According to our previous work [1], the 11-year (132 months) running mean values of the annual averages of the ionospheric parameters (foF2*₁₀₋₁₄(132), foF2*₂₃₋₀₁(132), and foF2*_{av}(132)) were calculated to obtain an independent picture of long-term trends in the upper ionosphere; the 11-year smoothing technique was also applied to the F10.7 and Ap data sets (F10.7*(132), Ap*(132)); Figure 3 presents these calculated values and shows that geomagnetic activity is strongly controlled by the solar activity, both show a dominant pattern of variations with the period ~34-36 years that is also reflected in all ionospheric parameters considered.




Figure 1 - The long-term variations of the near-noon(b), daily mean (c) and near-midnight (d) monthly median foF2 values in Alma-Ata [43.25°N, 76.92°E] in 1957-2017 together with the corresponding variations of solar flux F10.7 (a) and Ap index (e); solid dots – measured data, thick lines show the fitting lines, dashed lines – linear fits. Monthly median near-noon (f), daily mean (g), and near-midnight (h) foF2 values versus monthly mean F10.7; thick lines represent the linear regression lines for these data, dashed lines correspond to the polynomial functions of the second degree




Figure 2 - As the Figure 1 but the annual averages of the F10.7, Ap, and foF2 values.




Figure 3 - The 132-month smoothed values of the annual averages of the ionospheric F2-layer parameters, F10.7, and Ap.

This period is slightly differs from our earlier finding [1] where the period was estimated to be in a range of 30-32 years. The extended data sets used in the study reveal the period more clearly. It should be mentioned that similar period (31-32 years) has been found in all the solar terrestrial parameters ([4-6], and references therein) and was interpreted as the 35-year Bruter climatic periodicity. Note, the negative trend (dashed line), the lowering in the F2-layer critical frequency, is also seen in figure 3. The long-term trends found in different ionospheric characteristics (figures 1-3) are given in table.

Table						
#	Parameter	Trend, MHz/year	Parameter	Trend, MHz/year	Parameter	Trend, MHz/year
1	foF2 ₁₀₋₁₄ (med)	-0.02822	foF2* ₁₀₋₁₄	-0.02439	foF2* ₁₀₋₁₄ (132)	-0.01758
2	foF2 _{d.av} (med)	-0.01884	foF2* _{d.av}	-0.01586	foF2* _{d.av} (132)	-0.01077
3	foF2 ₂₃₋₀₁ (med)	-0.00932	foF2* ₂₃₋₀₁	-0.00756	foF2* ₂₃₋₀₁ (132)	-0.00364

The regression dependences of $foF2^*_{10-14}(132)$, $foF2^*_{23-01}(132)$, $foF2^*_{d.av}(132)$ on $F10.7^*(132)$ were studied for the period 1963-2012 (the 11-year smoothing technique that was applied to the $foF2^*$ data sets reduced the available period for study to between 1963 and 2012), the coefficients of determination (R^2) were found to be 0.96, 0.98, and 0.90 for near-noon, daily averaged and near-midnight data correspondingly figure 4a.




Figure 4 - The regression dependence between $foF2^*(132)$ and $F10.7^*(132)$ for the periods 1963-2012, (a), 1972-1984 (b) and 1984-2008 (c)

However, a combination of two groups of points was evident in figure 4 (upper panel) that assumed some different dependence between the parameters on different phases of the 34-36-years cycle (the period 1972-1984, and 1984-2008), when the $foF2^*(132)$ values increase to their maximum and then decrease to their minimum values. Examples of the regression dependences between $foF2^*(132)$ and $F10.7^*(132)$ for the two time periods are shown in figure 4a,b. It is distinctly seen in both cases that 99% of the variations in the ionospheric parameters can be explained by linear dependence between them and solar activity. So, the solar activity can be considered to be the main driver for the long-term variations in the ionospheric F2-region.




Figure 5 - The 132-month smoothed values of $fmin$ and $F10.7$ where dashed lines represent their linear trends for the whole measuring interval

Trends in the D-region. A similar analysis, as that carried out for $foF2$, has been carried out for the $fmin$ data observed at the Alma-Ata station in the years 1957-2017. As it has been already mentioned, the ionospheric parameter $fmin$ is used as a qualitative characteristic of the ionospheric absorption in the D-region, and our interest “is there any trend in the $fmin$ data, and if “yes” what is its sign?” For this purpose, we use the monthly median $fmin$ values for near-noon interval of local time averaging the data over three hours, from 11:00 LT to 13:00 LT. Then, the $fmin$ data were smoothed using the 11-year running mean smoothing to suppress effects of the 11-year solar cycle and obtain a picture of long-term variation in them. Figure 5 shows the $fmin(132)$ and $F10.7(132)$ variation together with the regression

lines (dashed lines) and demonstrates that the long-time course of $f_{min}(132)$ is similar to those found for F10.7 and foF2 (figures 1-3) in the event that the periodicity of 34-36 years is also evident in the $f_{min}(132)$ variation which is found in variations of the F2-layer parameters. However, figure 5 clearly illustrates a stable upward (positive) trend in $f_{min}(132)$ values opposite in sign to the trends observed in variations F10.7 and foF2. It means relative high sensitivity of the f_{min} values to the solar activity changes and significant influence of other trend drivers on them that is very possible the impact of anthropogenic factors on the state of the lower ionosphere.

Conclusion. The F2-layer critical frequency (foF2) and lowest frequency (f_{min}) observed at the mid-latitude ionospheric station Alma-Ata [43.25N, 76.92E] in the period 1957-2017 were used to study long-term trends in the upper (F2-layer) and lower (D-region) ionosphere.

The geomagnetic indexes (A_p), the near-noon, near-midnight, daily averaged foF2 are found to be in strong dependence from the solar activity; all of them show a dominant pattern of variation with a period of ~34-36 years and linear negative trend.

The trend magnitudes are found to be of the same order independently of the fact whether the solar activity effects in the data sets are smoothed or not; the trends are statistically significant and lie within of -0.018 to -0.028, -0.011 to -0.019, -0.0036 to -0.0093 MHz/year for near-noon, daily averaged and near-midnight values correspondingly.

The long-term course of f_{min} is similar to those found in F10.7, A_p , and foF2, i.e. the periodicity of ~34-36 years is also evident in the f_{min} variation. However, in contrast to the F2-layer parameters, the f_{min} variation clearly demonstrates an upward (positive) linear trend that is opposite in sign to the trend found in the F10.7. This means relatively high sensitivity of the f_{min} values to the solar activity changes and significant influence of other trend drivers on them, one of which is the possible impact of anthropogenic factors on the state of the lower ionosphere.

The work was carried out with accordance Grant № AP05131261 of Ministry of Education and Science RK.

А.Ф. Яковец, Г.И. Гордиенко, Ю.Г. Литвинов

ЕЖШС «Ионосфера институты» «ҰҒЗТО» АҚ, Алматы, Қазақстан

ИОНОСФЕРАНЫҢ F ЖӘНЕ D АЙМАҒЫНДАҒЫ ҰЗАҚ МЕРЗІМДІ ВАРИАЦИЯЛАРЫ

Аннотация. Алматы станциясындағы ионосфераның жер үсті тік радиозондауының деректері бойынша F2 Ионосфера қабатының (foF2) және жер маңындағы ең төменгі шағылысу жиіліктерінің (f_{min}) ұзақ мерзімді өзгерістері зерттелді [43.25 N, 76.92 E] 1957-2017 жж.

Сонымен қатар, ең аз шағылысу жиілігі төменгі ионосферада зондирлеуші сигналдың жұтылуына да, шу деңгейі мен ионозондтың техникалық сипаттамаларына да байланысты. Сондықтан сигналдың жұтылуының абсолюттік мәнін осындай жолмен бағалау проблемалы болып табылады. Алайда, қолшатыр сигналының жұту деңгейінің индикаторы ретінде сапалы сипаттама ретінде f_{min} параметрін пайдалануға болады.

Бастапқы деректер ретінде fof2 медиандық fof2 (10-14 LT), терек жанындағы (23-01 LT) сағат арифметикалық орташа мәндері және ионосфераның бақылауының қаралып отырған кезеңіндегі орташа тәуліктік мәндері қарастырылды. Төменгі ионосфераның өзгергіштігі күндізгі сағат үшін f_{min} деректерін пайдалану арқылы зерттелген. Күн радиосәулелендіру ағынының орташа айлық мәндері F10.7 және A_p индексі күн және геомагниттік белсенділіктің сипаттамалары ретінде рассматрен.

Күн және геомагниттік белсенділігі бойынша деректер ионосфераның жағдайына әсер ететін факторлар ретінде тартылды. Ионосфера аймағының F2 параметрлері, сондай-ақ A_p геомагниттік белсенділіктің индексі $T \approx 35$ жыл мерзімділігін және сызықтық теріс трендін анықтай отырып, күн белсенділігіне қатаң тәуелділікті сезінеді. FoF2 және F10.7 ұзақ мерзімді вариация арасындағы детерминация коэффициенттері 0.99 мәндеріне жетеді, бұл күн белсенділігі Ионосфера аймағының F2 ұзақ мерзімді өзгергіштігінің негізгі драйвері болып табылады деп болжайды.

Бұл нәтиже бір жағынан күн белсенділігіне жоғары тәуелділікті, екінші жағынан басқа факторлардың төменгі ионосфераның жағдайына, антропогендік факторларға әсерін болжайды.

Түйін сөздер: орташалықты ионосфера; жоғарғы және төменгі ионосфера; D-облыс; F2-облыс; ұзақ мерзімді трендтер.

А.Ф. Яковец, Г.И. Гордиенко, Ю.Г. Литвинов

ДТО «Институт ионосферы» АО «НЦКИТ», Алматы, Казахстан

ДОЛГОВРЕМЕННЫЕ ВАРИАЦИИ В ОБЛАСТИ F И D ИОНОСФЕРЫ

Аннотация. Исследованы долговременные изменения околоразуденных, среднесуточных и полуночных критических частот слоя F2 ионосферы (f_oF2) и околоразуденных минимальных частот отражения (f_{min}) по данным наземного вертикального радиозондирования ионосферы на станции Алма-Ата [43.25N, 76.92E] за период 1957-2017 гг.

Принималось во внимание, что минимальная частота отражения зависит как от поглощения зондирующего сигнала в нижней ионосфере, так и от уровня шумов и технических характеристик ионозонда. Поэтому оценить абсолютное значение поглощения сигнала таким способом представляется проблематичным. Однако, как качественная характеристика, как индикатор уровня поглощения зондирующего сигнала, параметр f_{min} можно использовать.

В качестве исходных данных рассмотрены арифметически средние значения медианных f_oF2 для околоразуденных (10-14 LT), околоразуденных (23-01 LT) часов и среднесуточные значения за рассматриваемый период наблюдений ионосферы. Изменчивость нижней ионосферы исследована с использованием данных f_{min} для дневных часов. Среднемесячные значения потока радиоизлучения Солнца F10.7 и индекс Ap рассмотрены как характеристики солнечной и геомагнитной активности.

Привлекались данные по солнечной и геомагнитной активности как факторы, оказывающие влияние на состояние ионосферы. Получено, что параметры F2 области ионосферы, а также индекс геомагнитной активности Ap испытывают строгую зависимость от солнечной активности, обнаруживая периодичность $T \approx 35$ лет и линейный отрицательный тренд. Коэффициенты детерминации взаимосвязи долгопериодных вариаций f_oF2 и F10.7 достигают значений 0.99, это предполагает, что солнечная активность является основным драйвером долговременной изменчивости F2 области ионосферы.

Значения f_{min} , обнаруживая подобно параметрам F2 области ионосферы ~35-летнюю периодичность, демонстрируют положительный тренд в наблюдаемый период 1957-2017 гг. Данный результат предполагает с одной стороны высокую зависимость от солнечной активности, с другой стороны влияние других факторов на состояние нижней ионосферы, возможно факторов антропогенных.

Ключевые слова: среднеширотная ионосфера; верхняя и нижняя ионосфера; D-область; F2-область; долговременные тренды.

Information about authors:

Gordienko Galina Ivanovna, Ph.D. mat. Sciences, ANS, Institute of the Ionosphere, JSC "NCSCIT", Almaty, ggordienko@mail.ru. 1) The author made a significant contribution to the research concept, obtaining data and interpretation, 2) the author wrote the first version of the article 3) the author approved the final version of the article before submitting it for publication. <https://orcid.org/0000-0002-9337-206X>;

Yakovets Arthur Fedorovich, Ph.D. mat. Sciences, VNS, Institute of the Ionosphere, JSC "NTSKIT", Almaty, artyak40@mail.ru. 1) The author made a significant contribution to the research concept, data analysis and interpretation, 2) the author wrote the first version of the article, 3) the author approved the final version of the article before submitting it for publication. <https://orcid.org/0000-0002-3374-9507>;

Litvinov Yuri Georgievich, Ph.D. mat. Sci., SSS, Institute of the Ionosphere, NCSCIT JSC, Almaty, yurii-litvinov@mail.ru. 1) The author made a significant contribution to obtaining the data and their analysis, 2) the author made a critical review of the article for important intellectual content, 3) the author approved the final version of the article before submitting it for publication. <https://orcid.org/0000-0002-0600-0517>

REFERENCES

- [1] Gordiyenko G. I., Vodyannikov V. V., Yakovets A. F. and Litvinov Yu. G. (2014), Earth Planets and Space, 66 (125), 1-6, <http://www.earth-planets-space.com/content/66/1/125>.
- [2] Lauter E. A., Entzian G. Winter anomaly 1980/81 as an example of stratomesospheric coupling//Phys. Solariterr. 1982. Vol. 18. P. 83-90.
- [3] Kokourov V. D. Minimum frequency of reflections as a climatic characteristic of the upper atmosphere// Geomagnetism and Aeronomy. 2003. Vol. 43(2). P.274-276 (in Russian).
- [4] Clúa de Gonzalez A. L., Gonzalez W. D., Dutra S. L. G., Tsurutani B. T. Periodic variation in the geomagnetic activity: a study based on the Ap index//J Geophys. Res. 1993. Vol. 98. P. 9215.
- [5] Echer E., Rigozo N. R., Nordemann D. J. R., and Vieira L. E. A. (2004). Prediction of solar activity on the basis of spectral characteristics of sunspot number//Annales Geophysicae. 2004. Vol.22. P. 2239-2243.
- [6] Prabhakaran Nayar S. R. Periodicities in solar activity and their signature in the terrestrial environment//IL WS WORKSHOP, GOA, February 19-24. 2006. P.1-9.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

www.nauka-nanrk.kz

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы: *М. С. Ахметова, Г. Б. Халидуллаева, Д. С. Аленов*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 08.06.2020.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
15 п.л. Тираж 300. Заказ 3.