

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
әл-Фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
Әл-Фараби атындағы
Қазақ ұлттық университетінің

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-Farabi Kazakh
National University

SERIES
PHYSICO-MATHEMATICAL

3 (331)

MAY – JUNE 2020

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р ы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі
Ғ.М. Мұтанов

Р е д а к ц и я а л қ а с ы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев У.У. проф., корр.-мүшесі (Қазақстан)
Жүсіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Д. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қырғыстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорусь)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19; 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2020

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Главный редактор
д.ф.-м.н., проф. академик НАН РК
Г.М. Мутанов

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жангаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф., чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физика-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219, 220; тел.: 272-13-19; 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2020

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

Editor in chief

doctor of physics and mathematics, professor, academician of NAS RK

G.M. Mutanov

Editorial board:

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof., corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskiy I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006.

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19; 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2020.2518-1726.32>

Volume 3, Number 331 (2020), 21 – 25

UDC 524.3-48, 524.3-59, 524.355

Y. K. Aimuratov

Fesenkov Astrophysical Institute, Almaty, Kazakhstan.

E-mail: aimuratov@aphi.kz

GAMMA-RAY BURSTS WITHOUT MEASURED REDSHIFTS I: ENERGY ESTIMATE AND LAT BORESIGHT ANGLE DISTRIBUTION

Abstract. *Context.* With this short note we start a series of communications dedicated to the study of gamma-ray bursts (GRBs) without measured redshifts.

Aims. We make an attempt to estimate energy of GRBs without measured distances based on population of bursts with measured redshift.

Methods. We assume that there is indeed universal bimodal distribution on duration which is followed by all the events. Using data on best-fit spectral parameters from GBM GRB Catalog we calculate energy for each burst given mean redshift values for short/long distribution. Additionally, we cross-correlate GBM GRB Catalog with general GBM Trigger Catalog in order to retrieve information on LAT boresight angle at the moment of event detection.

Results. We obtained estimate energy for each burst and built a distribution plot which follows the idea of bimodal distribution. A scatter plot for LAT boresight angles of short GRBs was obtained.

Conclusions. The energy-based histogram follows a general pattern for short/long distribution having distinct peaks. On the other hand, LAT boresight angles for short events shows quite uniform distribution, and being an instrumental feature, therefore cannot implicitly point to some intrinsic mechanisms responsible for high-energy photons generation.

Keywords: gamma-ray burst; redshift; LAT boresight angle; data analysis.

1. Introduction.

There is essential quantitative information one can calculate knowing the distance to the astrophysical object. Among these values isotropic-equivalent energy is the starting point for inferences in any theoretical model. An estimate of injected energy is crucial information for numerical simulations, especially for modelling the most energetic phenomenon of the universe such as gamma-ray burst (GRB).

The *Fermi Gamma-Ray Space Observatory* (previously GLAST) has detected over two thousand* unique Gamma-Ray Burst (GRB) events during the first decade of operation [1]. Only ~10% of these bursts have measured redshifts therefore making possible a reliable physical interpretation of the processes onsite. Consequently the rest of GRBs do not offer the latter possibility in full though containing spectral information.

The statistics to the date of July 20th 2018 is the following (see **Table 1**): *Fermi*-GBM telescope detected 2367 gamma-ray bursts (100%) among which there are 398 short (16.8%) and 1969 long (83.2%) duration GRBs, both are with and without measured redshift values; out of this total number of bursts the *Fermi*-LAT telescope detected 169 bursts (7.1%) among which there are 14 short and 155 long duration GRBs, both are with and without measured redshift values [2].

* <https://heasarc.gsfc.nasa.gov/W3Browse/fermi/fermigbrst.html>

Table 1 - GRB detection statistics for all-sky *Fermi*-GBM (8 keV–40 MeV) and *Fermi*-LAT (20 MeV–100 GeV) since the beginning of operational era in July 2008. The first and the last rows correspond to the first [2] and the second [4] LAT catalogs. Percentages in parentheses are related to the total number of bursts by GBM. Numbers were retrieved from online *Fermi*-GBM Burst Catalog and *Fermi*-LAT Burst list.

| End time | GBM | $T_{90} < 2$ s | $T_{90} > 2$ s | LAT | $T_{90} < 2$ s | $T_{90} > 2$ s |
|-----------|------|----------------|----------------|------------|----------------|----------------|
| July 2011 | 733 | 122 (16.6%) | 611 (83.4%) | 35 (4.8%) | 5 | 30 |
| July 2018 | 2367 | 398 (16.8%) | 1969 (83.2%) | 169 (7.1%) | 14 | 155 |

Within a decade of the *Fermi* operational era there were reported around 30 short GRBs with redshifts measured, among them there is only one GRB 090510 with both measured redshift and GeV photons detected [2]. This means a number of >350 short bursts does not have redshift and roughly same number without GeV photons were observed. As for the long bursts there are more than 200 events with known distances [3].

2. Methods.

In order to estimate energy for GRBs without redshift we take an average value of those short and long bursts with measured redshift. The *secure* median value for the redshift of short GRBs is around unity, while the median redshift value of long bursts is around two [5].

Fixing the redshift at unity (short) and two (long) values we calculate E_{iso} given a spectral model. For every GRB its `FlnC_Best_Fitting_Model` parameter from GBM catalog indicates the model best fitting the data for a spectrum accumulated over T_{90} duration of the burst [1]. The models used to fit each spectrum are: simple power law (plaw or PL), cutoff power law (comp or CPL), smoothly broken power law (SBPL) and Band function. Depending on model we will have various parameters to consider. The best-fit model is automatically selected by comparison of the values of the likelihood-based Castor C-Statistic (C-STAT). The C-STAT is a modified log-likelihood statistic based on Cash statistic, see [6], and it is used for fitting a model to data set through minimization. The isotropic energy equation has the form:

$$E_{iso} = \frac{4\pi d_L^2}{1+z} S(E_1, E_2, z), \quad (1)$$

where d_L is the luminosity distance and $S(E_1, E_2, z)$ is the fluence accumulated between minimum E_1 and maximum E_2 energy. It is defined by following expression:

$$S(E_1, E_2, z) = T_{90} \int_{E_1/(1+z)}^{E_2/(1+z)} EN(E) dE. \quad (2)$$

Here $N(E)$ defines the best-fit model of the prompt phase spectrum and the integration is done over GBM working energy band. Since GRBs are the cosmological objects then above mentioned luminosity distance will be defined, assuming a flat universe Λ CDM model with $\Omega_\Lambda=0.692$, $\Omega_M=0.308$, $H_0=67.8$ km s⁻¹ Mpc⁻¹ and $c=2.998e+10$ cm s⁻¹, by following equation [7]:

$$d_L(z, \Omega_\Lambda, \Omega_M) = (1+z) \frac{c}{H_0} \int_0^z \frac{dz'}{\sqrt{\Omega_M(1+z')^3 + \Omega_\Lambda}}. \quad (3)$$

As can be seen the redshift stays as only unknown parameter. The calculated values of $E_{iso}^{z=1}$ and $E_{iso}^{z=2}$ are shown on **Figure 1** and listed in **Table 2** of the next section.

Another issue to consider for future studies is a boresight angle of the LAT instrument at the moment of GBM trigger. This angle defines the sensitivity of LAT and simply represents itself a viewing angle outside of which no observation is possible. Given orientation coordinates two instruments can calculate mutual directions and when burst location defined by GBM the LAT normal angle is calculated with respect to that pointing and stored in GBM Trigger Catalog. We scan throughout (in order of priority) the *Fermi LAT Second GRB Catalog*,[†] the *Fermi GBM Trigger Catalog*[‡] and GCN archive[§] looking for the LAT boresight angle with regard to the best location of every burst.

[†] <https://heasarc.gsfc.nasa.gov/W3Browse/all/fermilgrb.html>

[‡] <https://heasarc.gsfc.nasa.gov/W3Browse/fermi/fermigtrig.html>

[§] <https://gcn.gsfc.nasa.gov/>

3. Results. We obtained estimate energy for each burst and built a distribution plot which follows the idea of bimodal distribution.

Table 2 lists all short GRBs observed within a decade of GBM operation and columns indicate: burst full name, duration (in seconds), rest-frame spectral peak energy calculated at $z=1.0$ (in keV), isotropic-equivalent energy calculated at $z=1.0$ (in erg, rest-frame by definition), and LAT boresight angle at the GBM trigger (in degrees). Although not listed here, a similar table is obtained for long ($T_{90} > 2$ s) duration GRBs, and both tables in full are available upon request.

Table 2 - List of 398 short ($T_{90} < 2$ s) GRBs within August 2008–July 2018 detected by *Fermi*-GBM with estimated isotropic-equivalent energy E_{iso} when given $z=1.0$ as an average redshift for short bursts. Since the table is large in size we show here only the first and the last three rows. Full table is available upon request

| GRB name | T_{90} (s) | $E_{peak}^{z=1}$ (keV) | $E_{iso}^{z=1}$ (erg) | θ (degrees) |
|-----------|--------------|------------------------|-----------------------|--------------------|
| 080723913 | 0.192 | 641.679 | 6.70615e+50 | — |
| 080725541 | 0.96 | 2965.18 | 4.87635e+51 | — |
| 080802386 | 0.576 | 1244.78 | 4.81246e+51 | — |
| ... | ... | ... | ... | ... |
| 180703949 | 1.536 | 273.749 | 4.60329e+52 | 32 |
| 180715741 | 1.664 | 1121.59 | 8.24943e+51 | 79 |
| 180715755 | 0.704 | 1804.56 | 8.58977e+51 | 75 |

We built an energy-based histogram for two populations of 1800 long and 363 short GRBs on figure 1. The median values are $2.16e+51$ erg (short) and $4.61e+52$ erg (long), and two distinct peaks are easily observed.

Regarding the LAT boresight angles there are 363 short GRBs with known θ at the trigger (for 35S-GRB there is no angle information, see table 3), of which 210 bursts were detected at $\theta < 75$ degrees, number of 6 short GRBs were triggered at $\theta = 75$ degrees and events in amount of 147 were detected at $\theta > 75$ degrees.

Table 3 - Distribution of short GRBs with regard to LAT boresight angle.

| Total S-GRBs | $\theta < 75$ degrees | $\theta = 75$ degrees | $\theta > 75$ degrees | no θ |
|--------------|-----------------------|-----------------------|-----------------------|-------------|
| 398 | 210 | 6 | 147 | 35 |

A scatter plot for LAT boresight angles of short GRBs was obtained on figure 2. Labeled red points indicate S-GRBs from the Second LAT Catalog [2]. Note that a large offset of S-GRB 170127067 still had high-energy emission at late repointing. The histogram shows the same distribution stacked in 30 bins.




Figure 1 - Histogram of isotropic-equivalent energy for 363 short and 1800 long GRBs for a decade of *Fermi* mission. The values of energy are calculated by assuming average redshifts for short ($z=1.0$, solid blue) and long ($z=2.0$, solid red) bursts. Dashed black line indicates $E_{iso}=1e52$ erg energy. Dashed blue and red lines indicate median energy values




Figure 2 - Distribution of 363 short GRBs by LAT boresight angle and isotropic-equivalent energy calculated under assumption of common $z=1$. Vertical dashed lines indicates $\theta=75$ degrees threshold angle for LAT instrument, horizontal dashed line indicates $E_{iso}=1e52$ erg energy

4. Discussion. The energy-based histogram follows a general pattern for short/long distribution having distinct peaks. These two populations when put on $E_{peak}-E_{iso}$ plane stays in accordance with spectral energy correlation for short [9] and long [10] GRBs. The prospect of further detailed study is possible due to the presence of redshift measured bursts [Aimuratov et al. (2020) in *Proceedings of the Fifteenth Marcel Grossmann Meeting*, submitted].

On the other hand, LAT boresight angles for short events shows quite uniform distribution, and being an instrumental feature, therefore, it cannot implicitly point to some intrinsic mechanisms responsible for high-energy photons generation. An outlier GRB 170127C detected ~ 300 s after GBM trigger [8] is one of the clear examples for prolonged GeV emission of still unknown nature. This implies

5. Acknowledgements. This article is dedicated to Dr. Larissa Andreevna Pavlova and Dr. Boris Ivanovich Demchenko. The work was done with partial support from research grants/programs BR05236322 from MES RK and BR05336383 from AC MDDIAI RK. Author additionally thanks research grant AP05133630 for partial support.

Е. К. Аймуратов

В. Г. Фесенков атындағы астрофизика институты, Алматы, Қазақстан

ҚЫЗЫЛ ЫҒЫСУЫ ӨЛШЕНБЕГЕН ГАММА-СӘУЛЕЛЕНУ ЖАРҚЫЛДАРЫ І: ЭНЕРГИЯНЫ БАҒАЛАУ ЖӘНЕ LAT КӨРУ БҰРЫШТАРЫНЫҢ ҮЛЕСТІРІМІ

Аннотация. Мәтін. Осы қысқаша хабарламамен біз қызыл ығысуы өлшенбеген гамма-сәулелену жарқылдарына (ГСЖ) арналған зерттеулер туралы мақалалар сериясын бастаймыз.

Мақсаты. Біз қызыл ығысуы өлшенген жарқылдар жиынтығына сүйеніп, қашықтығы өлшенбеген ГСЖ популяциясының энергия өлшемін бағалауға әрекет жасаймыз.

Әдісі. Біз барлық оқиғалар арасында әмбебап бимодалдық үлестіріміне сай болады деп негіздейміз. GBM GRB каталогындағы әр оқиғаға ең жақсы сәйкес келетін спектрлік параметрлер туралы мәліметтерді және қызыл ығысуы өлшенген қысқа/ұзақ уақытты оқиғалар үшін берілген орташа мәндерді қолдана отырып, біз энергияны есептейміз. Сонымен қатар біз оқиғаларды анықтаған сәтте LAT инструментінің көру бұрышы туралы ақпарат алу үшін GBM GRB каталогын GRB Trigger Catalog жалпы каталогымен өзара салыстырамыз.

Нәтижелері. Біз әр ГСЖ-ның энергиясын есептеп алдық және нәтижесінде бимодалды үлестірім идеясына сәйкес келетін алаңын алдық. Қысқа уақытты оқиғалардың LAT көру бұрыштары үшін шашырау сюжеті алынды.

Қорытындысы. Энергияға негізделген гистограмма нақты шыңдары барын көрсете отырып, қысқа/ұзақ уақытты оқиғалардың үлестіріміне сай жалпы үлгісін ұстанады. Екінші жағынан, қысқа оқиғаларға арналған

LAT көру бұрыштары біркелкі таралуды көрсетеді және инструменталды сипат болып табылады, сондықтан жоғары энергиялы фотондардың пайда болуына жауап беретін кейбір ішкі механизмдерді нақты көрсете алмайды.

Түйін сөздер: гамма-сәулелену жарқылы; қызыл ығысуы; LAT көру бұрышы; деректерді талдау.

Е. К. Аймуратов

Астрофизический институт имени В. Г. Фесенкова, Алматы, Казахстан

ГАММА-ВСПЛЕСКИ БЕЗ ИЗМЕРЕННЫХ КРАСНЫХ СМЕЩЕНИЙ I: ОЦЕНКА ЭНЕРГИЙ И РАСПРЕДЕЛЕНИЕ УГЛОВ ПРИЦЕЛИВАНИЯ LAT

Аннотация. *Контекст.* С этой короткой заметки мы начинаем серию сообщений, посвященных изучению гамма-всплесков (ГВ) без измеренных красных смещений.

Цели. Мы предпринимаем попытку оценить энергию ГВ без измеренных расстояний на основе совокупности всплесков с измеренным красным смещением.

Методы. Мы предполагаем, что, действительно, существует универсальное бимодальное распределение по продолжительности, которому следуют все события. Используя данные о спектральных параметрах из каталога GBM GRB для наиболее подходящих моделей, мы вычисляем изотропную энергию для каждого всплеска, учитывая средние значения красного смещения для распределения коротких/длинных событий. Кроме того, мы производим взаимную корреляцию каталога GBM GRB с общим каталогом GBM Trigger Catalog для получения информации об угле прицеливания LAT в момент обнаружения события.

Результаты. Мы получили оценку энергии для каждого всплеска и построили график распределения, который следует идее бимодального распределения. Получен график распределения углов прицеливания LAT для коротких GRB.

Заключение. Гистограмма по энергиям ГВ следует общей схеме распределения по коротким/длинным событиям, показывая два отчетливых пика. С другой стороны, углы прицеливания LAT для коротких ГВ показывают довольно равномерное распределение, и, являясь инструментальной особенностью, не могут неявно указывать на какие-либо внутренние механизмы, ответственные за образование фотонов высоких энергий.

Ключевые слова: гамма-всплеск; красное смещение; угол прицеливания LAT; анализ данных.

Information about author:

Yerlan Kairatovich Aimuratov, a PhD in Relativistic Astrophysics, Junior Research Associate at the Laboratory of Stars and Nebulae Physics, Department of Observational Astronomy, Fesenkov Astrophysical Institute. Address: Observatory 23, Kamenskoye Plateau, Medeu District, 050020 Almaty, Kazakhstan. aimuratov@aphi.kz, <https://orcid.org/0000-0001-5717-6523>

REFERENCES

- [1] von Kienlin A. et al. (2020) The Fourth Fermi-GBM Gamma-Ray Burst Catalog: A Decade of Data // *ASTROPHYS J*, 893:46. <https://doi.org/10.3847/1538-4357/ab7a18>
- [2] Ajello M. et al. (2019) A Decade of Gamma-Ray Bursts Observed by Fermi-LAT: The Second GRB Catalog // *ASTROPHYS J*, 878:52. <https://doi.org/10.3847/1538-4357/ab1d4e>
- [3] Zhang Z. B. et al. (2018) Spectrum-Energy Correlations in GRBs: Update, Reliability, and the Long/Short Dichotomy // *PUBLICATIONS OF THE ASTRONOMICAL SOCIETY OF THE PACIFIC*, **130**:054202. <https://doi.org/10.1088/1538-3873/aaa6af>
- [4] Ackermann M. et al. (2013) The First Fermi-LAT Gamma-Ray Burst Catalog // *ASTROPHYS J SUPPL S*, 209:11. <https://doi.org/10.1088/0067-0049/209/1/11>
- [5] Berger E. (2014) Short-Duration Gamma-Ray Bursts // *ANNUAL REVIEW OF ASTRONOMY AND ASTROPHYSICS*, **52**:43. <https://doi.org/10.1146/annurev-astro-081913-035926>
- [6] Cash W. (1979) Parameter Estimation in Astronomy through Application of the Likelihood Ratio // *ASTROPHYS J*, 228:939.
- [7] Weinberg S. (1972) *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*. Weinheim: Wiley-VCH, 688. ISBN: 978-0-471-92567-5
- [8] Moretti E. et al. (2017) GRB 170127C: Fermi-LAT detection // *GCN Circular Archive #20544*.
- [9] Zhang Z. B. et al. (2018) Spectrum-Energy Correlations in GRBs: Update, Reliability, and the Long/Short Dichotomy // *PUBLICATIONS OF THE ASTRONOMICAL SOCIETY OF THE PACIFIC*, **130**:054202. <https://doi.org/10.1088/1538-3873/aaa6af>
- [10] Amati L. et al. (2019) Addressing the Circularity Problem in the Ep-Eiso Correlation of Gamma-Ray Bursts // *MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY LETTERS*, **486**:L46. <https://doi.org/10.1093/mnras/slz056>

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

www.nauka-nanrk.kz

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы: *М. С. Ахметова, Г. Б. Халидуллаева, Д. С. Аленов*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 08.06.2020.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
15 п.л. Тираж 300. Заказ 3.