

ISSN 2518-1726 (Online),  
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ  
әл-Фараби атындағы Қазақ ұлттық университетінің

# Х А Б А Р Л А Р Ы

---

---

## ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН  
Қазақстан Республикасының  
Ғылым Академиясының  
Әл-Фараби атындағы  
Қазақ ұлттық университетінің

## NEWS

OF THE NATIONAL ACADEMY OF SCIENCES  
OF THE REPUBLIC OF KAZAKHSTAN  
Al-Farabi  
Kazakh National University

**SERIES  
PHYSICO-MATHEMATICAL**

**5 (333)**

**SEPTEMBER – OCTOBER 2020**

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р ы  
ф.-м.ғ.д., проф., ҚР ҰҒА академигі  
**Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

**Асанова А.Т.** проф. (Қазақстан)  
**Бошкаев К.А.** PhD докторы (Қазақстан)  
**Байгунчеков Ж.Ж.** проф., академик (Қазақстан)  
**Вишневский И.Н.** проф., академик (Украина)  
**Quevedo Hernando** проф. (Мексика),  
**Жүсіпов М.А.** проф. (Қазақстан)  
**Ковалев А.М.** проф., академик (Украина)  
**Калимолдаев М.Н.** проф., академик (Қазақстан)  
**Михалевич А.А.** проф., академик (Белорусь)  
**Молдабеков М. М.** проф., академик (Қазақстан)  
**Мырзакулов Р.** проф., академик (Қазақстан)  
**Өмірбаев У.У.** проф., академик (Қазақстан)  
**Пашаев А.** проф., академик (Әзірбайжан)  
**Рамазанов Т.С.** проф., академик (Қазақстан)  
**Такибаев Н.Ж.** проф., академик (Қазақстан), бас ред. орынбасары  
**Тигиняну И.** проф., академик (Молдова)  
**Тулешов А.К.** проф., чл.-корр. (Қазақстан)  
**Уалиев З.Г.** проф., чл.-корр. (Қазақстан)

**«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».**

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан Республикасының Ақпарат және коммуникациялар министрлігінің Ақпарат комитетінде 14.02.2018 ж. берілген № 16906-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Тақырыптық бағыты: *физика-математика ғылымдары және ақпараттық технологиялар саласындағы басым ғылыми зерттеулерді жариялау.*

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19; 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2020

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Главный редактор  
д.ф.-м.н., проф. академик НАН РК  
**Г.М. Мутанов**

Редакционная коллегия:

**Асанова А.Т.** проф. (Казахстан)  
**Бошкаев К.А.** доктор PhD (Казахстан)  
**Байгунчеков Ж.Ж.** проф., академик (Казахстан)  
**Вишневский И.Н.** проф., академик (Украина)  
**Quevedo Hernando** проф. (Мексика),  
**Жусупов М.А.** проф. (Казахстан)  
**Ковалев А.М.** проф., академик (Украина)  
**Калимолдаев М.Н.** проф., академик (Казахстан)  
**Михалевич А.А.** проф., академик (Беларусь)  
**Молдабеков М. М.** проф., академик (Казахстан)  
**Мырзакулов Р.** проф., академик (Казахстан)  
**Пашаев А.** проф., академик (Азербайджан)  
**Рамазанов Т.С.** проф., академик (Казахстан)  
**Такибаев Н.Ж.** проф., академик (Казахстан), зам. гл. ред.  
**Тигиняну И.** проф., академик (Молдова)  
**Тулешов А.К.** проф., чл.-корр. (Казахстан)  
**Уалиев З.Г.** проф., чл.-корр. (Казахстан)  
**Умирбаев У.У.** проф., академик (Казахстан)

**«Известия НАН РК. Серия физика-математическая».**

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и коммуникаций Республики Казахстан № 16906-Ж, выданное 14.02.2018 г.

Тематическая направленность: *публикация приоритетных научных исследований в области физико-математических наук и информационных технологий.*

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219, 220; тел.: 272-13-19; 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© Национальная академия наук Республики Казахстан, 2020

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

Editor in chief  
doctor of physics and mathematics, professor, academician of NAS RK  
**G.M. Mutanov**

Editorial board:

**Asanova A.T.** prof. (Kazakhstan)  
**Boshkayev K.A.** PhD (Kazakhstan)  
**Baigunchekov Zh.Zh.** prof., akademik (Kazakhstan)  
**Vishnevskiy I.N.** prof., academician (Ukraine)  
**Quevedo Hernando** prof. (Mexico),  
**Zhusupov M.A.** prof. (Kazakhstan)  
**Kovalev A.M.** prof., academician (Ukraine)  
**Kalimoldaev M.N.** prof., akademik (Kazakhstan)  
**Mikhalevich A.A.** prof., academician (Belarus)  
**Moldabekov M. M.** prof., akademik (Kazakhstan)  
**Myrzakulov R.** prof., akademik (Kazakhstan)  
**Pashayev A.** prof., academician (Azerbaijan)  
**Ramazanov T.S.** prof., akademik (Kazakhstan)  
**Takibayev N.Zh.** prof., academician (Kazakhstan), deputy editor in chief.  
**Tiginyanu I.** prof., academician (Moldova)  
**Tuleshov A.K.** prof., chl.-korr. (Kazakhstan)  
**Ualiev Z.G.** prof., chl.-korr. (Kazakhstan)  
**Umirbayev U.U.** prof., academician (Kazakhstan)

**News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.**  
ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Communications of the Republic of Kazakhstan **No. 16906-Ж**, issued on 14.02.2018.

Thematic scope: *publication of priority research in the field of physical and mathematical sciences and information technology.*

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19; 272-13-18,  
<http://physics-mathematics.kz/index.php/en/archive>

---

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN  
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2020.2518-1726.89>

Volume 5, Number 333 (2020), 106 – 112

UDK 524.4; 524.82; 524.83; 524.85

IRSTI 41.29.15; 41.29.17; 41.29.25; 41.29.33

**S.R. Myrzakul<sup>1,2</sup>, Y.M. Myrzakulov<sup>1,2</sup>, M. Arzymbetova<sup>3</sup>**

<sup>1</sup>L.N. Gumilyov Eurasian National University, Nur-Sultan, Kazakhstan;

<sup>2</sup>Ratbay Myrzakulov Eurasian International Centre for Theoretical Physics LLP, Nur-Sultan, Kazakhstan;

<sup>3</sup>Al-Farabi Kazakh National University, Almaty, Kazakhstan.

E-mail: [smyrzakul@gmail.com](mailto:smyrzakul@gmail.com), [ymyrzakulov@gmail.com](mailto:ymyrzakulov@gmail.com), [arzymbetova.moldir@gmail.com](mailto:arzymbetova.moldir@gmail.com)

## INFLATION IN $F(R, T)$ GRAVITATION WITH $f$ -ESSENCE

**Abstract.** Modified theories of gravity have become a kind of paradigm in modern physics because they seem to solve several shortcomings of the standard General Theory of Relativity (GTR) related to cosmology, astrophysics and quantum field theory. The most famous modified theories of gravity are  $F(R)$  and  $F(T)$  theories of gravity. A generalization of these two modified theories and gravitations, which was first proposed by Myrzakulov Ratbay. In this paper, we study an inhomogeneous isotropic cosmological model with a fermion field  $f$ -essence whose action has the form  $S_{RT} = \int d^4x \sqrt{-g} [F(R, T) + L_m]$ , where  $R$  is the scalar of curvature, and  $T$  is the torsion scalar, and  $L_m$  is the Lagrangian  $f$ -essence. A particular case  $F(R, T) = \alpha R + \beta T$  is studied in detail when parameters are obtained that describe the current accelerated expansion of the Universe. The type of Lagrangian  $f$ -essence of this model is determined. The presented results show that gravity with  $f$ -essence can describe inflation in the early evolution of the Universe. A modified  $F(R, T)$  gravity with  $f$ -essence is considered. Equations of motion were obtained and the inflationary period of the early Universe was considered. To describe the inflationary period, the form of the Hubble parameter and the slow-roll parameter were determined.

**Key words:** Universe, accelerated expansion, inflation, cosmology,  $f$ -essence, slow-roll parameter, Friedmann-Robertson-Walker metric,  $F(R, T)$ -gravity.

**Introduction.** The idea of expanding Einstein's theory of gravity is productive and economical in terms of several attempts that try to solve problems by adding new and, in most cases, unjustified new components to give a self-consistent picture of dynamics. The accelerated expansion of the Hubble flow observed today and the missing material of astrophysical large-scale structures are primarily included in these considerations. Both problems can be solved by changing the gravity sector, i.e. field equations. Philosophy is an alternative for adding new cosmic fluids (new components in the field equations), which should lead to cluster structures (dark matter) or to accelerated dynamics (dark energy) due to exotic equations of state. In particular, weakening the hypothesis that the gravitational Lagrangian should be a linear function of the scalar of Ricci curvature  $R$ , as in the Hilbert-Einstein formulation, one can take into account the effective action as a minimum extension when the gravitational Lagrangian is a general function.

The main and well-known examples of such modified theories of gravity are  $F(R)$  theory of gravity and  $F(T)$  theory of gravity. In this direction, the generalization of the theories of gravity  $F(R)$  and  $F(T)$  promises to be a promising and interesting version of modified theories of gravity. One of these generalizations of the  $F(R)$  and  $F(T)$  theories was first presented in [1] (see also [2-7]). There was shown that the torsion scalar, as well as the curvature scalar without a source of matter, leads to the formation of this union, the expansion of the Universe, inflation, and so on. This generalization includes all the properties of the  $F(R)$  and  $F(T)$  theory of gravity.

Astronomical observations suggest the universe can be viewed as homogeneous and isotropic for cosmological purposes that at very large scales, and so this is well described by the Friedmann-Robertson-Walker (FRW) metric. In this paper, we investigate the dynamics of the fermionic field of the  $f$ -essence in the framework of  $F(R, T)$  gravity described by the FRW metric. The special case of  $F(R, T) = \alpha R + \beta T$ , where  $\alpha$  and  $\beta$  are real constants, is investigated in detail. The necessary cosmological parameters have been obtained.

The expansion in the early time of space-time in the evolution of the Universe, namely inflation, is introduced to solve some cosmological problems. The discovery of a gravitational wave from a heavy celestial body and the observation of a black hole (BH) is a great success of Einstein's general relativity. However, from a cosmological point of view, some problems remain, such as flatness, horizon, monopole problems.

The theory of inflation is one of the elegant and simple solutions for these problems. An alternative approach is to modify Einstein's gravity.

The standard method for determining observable inflation is to perform detailed disturbance analysis [8]. However, the approach can be simplified by introducing slow-roll parameters [9], either when inflation occurs due to modification of standard gravitational actions, or due to scalar fields.

### $F(R, T)$ gravity

The action has the following form [1] (see also [2-17]):

$$S_{RT} = \int d^4x \sqrt{-g} [F(R, T) + L_m], \quad (1)$$

where  $L_m$  is the matter Lagrangian,  $R$  is the curvature scalar,  $T$  is the torsion scalar

$$R = g^{\mu\nu} R_{\mu\nu}, \quad T = S_{\rho}^{\mu\nu} T^{\rho}_{\mu\nu}. \quad (2)$$

This generalization was first proposed in [1], and these properties were investigated in [2-17]. In general,  $G^{\lambda}_{\mu\nu}$  has the form

$$G^{\lambda}_{\mu\nu} = e_i^{\lambda} \partial_{\mu} e^i_{\nu} + e_j^{\lambda} e^i_{\nu} \omega^j_{i\mu} = \Gamma^{\lambda}_{\mu\nu} + K^{\lambda}_{\mu\nu}, \quad (3)$$

where  $\Gamma^{\lambda}_{\mu\nu}$  - Levi-Civita connection, and  $K^{\lambda}_{\mu\nu}$  - contorsion tensor. We write the metric in the standard form as

$$ds^2 = g_{ij} dx^i dx^j. \quad (4)$$

Orthonormal tetrad components  $e_i(x^{\mu})$  are related to the metric tensor as follows

$$g_{\mu\nu} = \eta_{ij} e^i_{\mu} e^j_{\nu}. \quad (5)$$

The orthonormalized condition is given as

$$\eta_{ij} = g_{\mu\nu} e^{\mu}_i e^{\nu}_j. \quad (6)$$

Here's  $\eta_{ij} = \text{diag}(-1, 1, 1, 1)$  where we used the relation

$$e^i_{\mu} e^{\mu}_j = \delta^i_j. \quad (7)$$

### Basics of $f$ -essence

Briefly describe some of the basics fermion model  $f$ -essences indicated in [7]. The action of the model has the form [7]

$$S_f = \int d^4x \sqrt{-g} [R + 2K(Y, \psi, \bar{\psi})], \quad (8)$$

where  $\psi$  and  $\bar{\psi} = \psi^{\dagger} \gamma^0$  denote the fermionic field and its conjugation, respectively, the cross means complex conjugation and  $R$  is the Ricci scalar.  $K$  is the density of the Lagrangian of the fermionic field, the canonical kinetic term of which has the form

$$Y = \frac{1}{2}i[\bar{\psi}\Gamma^\mu D_\mu\psi - (D_\mu\bar{\psi})\Gamma^\mu\psi]. \quad (9)$$

Here  $\Gamma^\mu = e_a^\mu \gamma^a$  is the generalized Dirac-Pauli matrices satisfying the Clifford algebra

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}, \quad (10)$$

where the brackets denote the anti-commutation relation.  $e_a^\mu$  denotes a tetrad, while the covariant derivative is given by

$$D_\mu\psi = \partial_\mu\psi - \Omega_\mu\psi, \quad D_\mu\bar{\psi} = \partial_\mu\bar{\psi} + \bar{\psi}\Omega_\mu. \quad (11)$$

The fermionic coupling  $\Omega_\mu$  is determined by the formula

$$\Omega_\mu = -\frac{1}{4}g_{\rho\sigma}[\Gamma_{\mu\delta}^\rho - e_b^\rho\partial_\mu e_\delta^b]\Gamma^\sigma\Gamma^\delta, \quad (12)$$

where  $\Gamma_{\mu\delta}^\rho$  Christoffel symbols.

Fermionic fields are considered here as classically commuting fields. By the classical fermionic field, we mean a set of four complex space-time functions that transform in accordance with the spinor representation of the Lorentz group. The existence of such fields have of decisive importance in our work, because, despite the fact that fermions are described by quantized fermion fields which that do not have a classical limit, we assume that such classical fields exist and use them as matter fields associated with gravity.

### **$F(R,T)$ gravity with $f$ -essence**

Now let's consider  $F(R,T)$  gravity with  $f$ -essence. Its action is given as

$$S = \int d^4x \sqrt{-g}[F(R,T) + 2K(Y, \psi, \bar{\psi})], \quad (13)$$

where  $R$  and  $T$  are curvature and torsion scalars, respectively, expressed as

$$R = g^{\mu\nu}R_{\mu\nu}, \quad T = S_\rho^{\mu\nu}T^\rho_{\mu\nu}. \quad (14)$$

### **FRW metric case**

To simplify, this section will make the next two to simplify the problem, namely, we assume that:

1)  $F$  is of the form

$$F = \alpha R + \beta T, \quad (15)$$

where  $\alpha$  and  $\beta$  are some positive constants.

2) consider only the FRW space-time

$$ds^2 = -dt^2 + a(t)^2(dx^2 + dy^2 + dz^2). \quad (16)$$

Then action (13) takes the form

$$S = \int dx^4 \sqrt{-g}[\alpha R + \beta T + 2K(Y, \psi, \bar{\psi})]. \quad (17)$$

Next important suggestion is that for the FRW metric,  $R$  and  $T$  take the forms

$$R = u + 6\left(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2}\right) = u + 6(\dot{H} + 2H^2) = u + R_S, \quad (18)$$

$$T = v - 6\frac{\dot{a}^2}{a^2} = v - 6H^2 = v - T_S,$$

where  $u$  and  $v$  are some functions from  $a$ ,  $a$ ,  $\frac{d^j a}{dt^j}$ ,  $R, T$  and so on. Here

$$R_S = 6\left(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2}\right) = 6(\dot{H} + 2H^2), \quad (19)$$

$$T_S = -6\frac{\dot{a}^2}{a^2} = -6H^2.$$

Next step for our model can be rewritten as

$$S = \int dx^4 \sqrt{-g} [\alpha R_r + \beta T_w + \bar{L}_m], \tag{20}$$

where

$$\bar{L}_m = \alpha u + \beta v + L_m = \alpha u + \beta v + 2K(Y, \psi, \bar{\psi}) = 2\bar{K}(Y, \psi, \bar{\psi}). \tag{21}$$

For the FRW metric, the Dirac matrices of the curved space-time  $\Gamma^\mu$  have the form

$$\Gamma^0 = \gamma^0, \Gamma^j = a^{-1}\gamma^j, \Gamma^5 = -i\sqrt{-g}\Gamma^0\Gamma^1\Gamma^2\Gamma^3 = \gamma^5, \Gamma_0 = \gamma^0, \Gamma_j = a\gamma^j (i=1,2,3). \tag{22}$$

Hence we get

$$\Omega_0 = 0, \Omega_j = \frac{1}{2}\dot{a}\gamma^j\gamma^0 \tag{23}$$

and

$$Y = \frac{1}{2}i(\bar{\psi}\gamma^0\dot{\psi} - \dot{\bar{\psi}}\gamma^0\psi) \tag{24}$$

Finally, note that the gamma matrices that we write in the Dirac basis are of the form

$$\gamma^0 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \gamma^k = \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix}, \gamma^5 = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \tag{25}$$

where  $I = diag(1,1)$  and  $\sigma^k$  are Pauli matrices having the following forms

$$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \tag{26}$$

We rewrite action (20) in the following form

$$S = \int dt \sqrt{-g} L, \tag{27}$$

where  $L$  is the point-like Lagrangian. To find the equations of motion of the model, we use the Euler-Lagrange equations. In our case, the Euler-Lagrange equations are defined as follows

$$\frac{\partial L}{\partial a} - \frac{d}{dt} \frac{\partial L}{\partial \dot{a}} = 0, \frac{\partial L}{\partial R} - \frac{d}{dt} \frac{\partial L}{\partial \dot{R}} = 0, \frac{\partial L}{\partial T} - \frac{d}{dt} \frac{\partial L}{\partial \dot{T}} = 0, \frac{\partial L}{\partial \psi} - \frac{d}{dt} \frac{\partial L}{\partial \dot{\psi}} = 0, \frac{\partial L}{\partial \bar{\psi}} - \frac{d}{dt} \frac{\partial L}{\partial \dot{\bar{\psi}}} = 0. \tag{28}$$

Accordingly, the condition is given as

$$E_L = \frac{\partial L}{\partial \dot{a}} \dot{a} + \frac{\partial L}{\partial \dot{R}} \dot{R} + \frac{\partial L}{\partial \dot{T}} \dot{T} + \frac{\partial L}{\partial \dot{\psi}} \dot{\psi} + \frac{\partial L}{\partial \dot{\bar{\psi}}} \dot{\bar{\psi}} - L = 0. \tag{29}$$

Finally, the equations of motion for our model take the form

$$\begin{aligned} 3H^2 - \bar{\rho} &= 0, \\ 2\dot{H} + 3H^2 + \bar{p} &= 0, \\ \bar{K}_Y \dot{\psi} + 0.5(3H\bar{K}_Y + \dot{\bar{K}}_Y)\psi - i\gamma^0 \bar{K}_{\bar{\psi}} &= 0, \\ \bar{K}_Y \dot{\bar{\psi}} + 0.5(3H\bar{K}_Y + \dot{\bar{K}}_Y)\bar{\psi} + i\bar{K}_{\psi} \gamma^0 &= 0, \end{aligned} \tag{30}$$

where

$$\bar{\rho} = Y\bar{K}_Y - \bar{K}, \quad \bar{p} = \bar{K}. \tag{31}$$

where  $\dot{H} = \dot{a}/a$  is the Hubble parameter,  $a(t)$  is the scale factor, and  $\dot{a}(t)$  is cosmic time derivative  $t$ .

From the second equation of system (30) can be seen that, according to the rules of differentiation, if expressions from the spirit of the sides of equality depend on different arguments, then this equality will be constant

$$2\dot{H}(a) + 3H(a)^2 = -\bar{K}(Y, \psi, \bar{\psi}) = const.$$



To study inflation is enough to determine the form of the Hubble parameter, which has the form:

$$H = \frac{1}{6} \frac{\left( \left( 12\sqrt{3} \sqrt{\frac{4C_1^3 + 27e^{2t}C_2}{C_2}} + 108e^t \right) C_2^2 \right)^{1/3}}{C_2} \frac{2C_1}{\left( \left( 12\sqrt{3} \sqrt{\frac{4C_1^3 + 27e^{2t}C_2}{C_2}} + 108e^t \right) C_2^2 \right)^{1/3}} \quad (32)$$

**Inflation.** Let us consider an inflationary model with a minimum kinetic term, where the behavior of the system is described by the FRW and Dirac equations (30). To describe the evolution of the background is convenient to introduce the functions of the Hubble flow  $\varepsilon_n$ , which are defined by the formula

$$\varepsilon_{n+1} \equiv -\frac{d \ln|\varepsilon_n|}{dN}, \quad n \geq 0, \quad (33)$$

where  $\varepsilon_0 \equiv H_{ini}/H$  and  $N \equiv \ln(a_{ini}/a)$  are the amount of e-folding. By definition, inflation is a phase of accelerated expansion  $\ddot{a}/a > 0$  or, equivalently  $\varepsilon_1 < 1$ . As a consequence, the end of inflation is determined by the condition  $\varepsilon_1 = 1$ . On the other hand, the conditions of slow-rolling down (or the slow-rolling approximation) refer to the situation when all  $\varepsilon_n$  satisfy  $\varepsilon_n \ll 1$ . For our case, the inflation solution evolves with the (positive) Hubble flow functions

$$\varepsilon_1 = -\frac{\dot{H}}{H^2}, \quad (34)$$

$$\varepsilon_2 = -\frac{2\dot{H}}{H^2} + \frac{\ddot{H}}{H\dot{H}}, \quad (35)$$

where  $C_1$  and  $C_2$  are integration constants. Figure 1 shows the typical behavior of the slow-roll parameter for different  $C_1$  and  $C_2$  values.




Figure 1 - Graphical analysis of  $\varepsilon_1$  (black lines) and  $\varepsilon_2$  (red lines). Dotted line:  $C_1 > 0$  and  $C_2 > 0$ ; dotted line:  $C_1 < 0$  and  $C_2 > 0$ ; long dotted line:  $C_1 < 0$  and  $C_2 < 0$ ; dot-dotted line:  $C_1 > 0$  and  $C_2 < 0$

**Conclusion.** In this work we have considered a modified  $F(R, T)$  gravity with  $f$ -essence. Equations of motion were obtained and the inflationary period of the early Universe was considered. To describe the inflationary period, we determined the form of the Hubble parameter and the slow-roll parameter. As can be seen from the graph for our model, the slow-roll parameter does not satisfy the inflation conditions given in [9]. Thus, the given  $F(R, T)$  gravity model with  $f$ -essence cannot describe viable inflation.

**Acknowledgement.** This work was supported by the Ministry of Education and Science of RK Grant No. AP08052197.

Ш.Р. Мырзақұл<sup>1,2</sup>, Е.М. Мырзақулов<sup>1,2</sup>, М. Арзимбетова<sup>3</sup>

<sup>1</sup>Л.Н. Гумилев атындағы Еуразия ұлттық университеті, Нұр-Сұлтан, Қазақстан;

<sup>2</sup>Ratbay Myrzakulov Eurasian International Centre for Theoretical Physics, Нұр-Сұлтан, Қазақстан;

<sup>3</sup>Әл-Фараби атындағы Қазақ ұлттық университеті, Алматы қ., Қазақстан

### *f*-ЭССЕНЦИЯСЫ БАР $F(R,T)$ ГРАВИТАЦИЯСЫНДАҒЫ ИНФЛЯЦИЯ

**Аннотация.** Жаңартылған гравитациялық теориялары қазіргі физикада өзгеше парадигманың түріне айналды, өйткені олар космология, астрофизика және кванттық өріс теориясымен байланысты стандартты жалпы салыстырмалылық теориясының бірнеше кемшіліктерін шешкен болуы мүмкін. Гравитациялықтың ең танымал модификацияланған теориялары –  $F(R)$  және  $F(T)$ . Осы екі теорияның жалпыланған түрін алғаш рет Мырзақулов Ратбай ұсынған. Бұл жұмыста біз *f*-эссенция фермиондық өрісі бар біртекті емес изотропты космологиялық модельді зерттейміз, әсерін былай көрсетеміз:  $S_{RT} = \int d^4x \sqrt{-g} [F(R,T) + L_m]$ , мұнда  $R$  – қисық скаляры және  $T$  – ширату скаляры,  $L_m$  – *f*-эссенция лагранжианы. Әлемнің үдетілген кеңею жағдайын сипаттайтын параметрлер алынған. Нақты жағдай  $F(R,T) = \alpha R + \beta T$  қарастырылып, толықтай зерттеледі. Бұл модельдің *f*-эссенция лагранжианың түрі анықталған. Ұсынылған нәтижелер *f*-эссенциясы бар  $F(R,T)$  гравитациясы әлемнің ерте инфляциялық эволюциясын сипаттайтындығын көрсетеді.

Жұмыста фермиондық өрісі бар және канондық емес кинетикалық мүшесі (*f*-эссенция) бар космологиялық модель қарастырылған. Модельдің кейбір нақты шешімдері табылған және біз осындай гравитациялық-фермиондық өзара әрекеттесудің әлемнің бақыланатын үдемелі ұлғаю әсерін тексереміз. Әрі қарай, осы модельдердің космологиялық қосымшалары талқыланады. Сондай-ақ, *f*-эссенция моделінде Эйнштейннің статикалық әлемі секілді шешім жоқ. *f*-эссенциясы бар модификацияланған  $F(R,T)$  гравитациясы қарастырылған. Қозғалыс теңдеулері алынды және алғашқы Әлемнің инфляциялық кезеңі қарастырылған. Инфляциялық кезеңді сипаттау үшін Хаббл параметрінің формасы және баяу оралу параметрі анықталған.

**Түйін сөздер:** әлем, үдемелі ұлғаю, инфляция, космология, *f*-эссенция, баяу сырғу параметрі, Фридман-Робертсон-Уокер метрикасы,  $F(R,T)$ -гравитациясы.

Ш.Р. Мырзақұл<sup>1,2</sup>, Е.М. Мырзақулов<sup>1,2</sup>, М. Арзимбетова<sup>3</sup>

<sup>1</sup>Евразийский национальный университет им. Л.Н.Гумилева, г. Нур-Султан, Казахстан

<sup>2</sup>Ratbay Myrzakulov Eurasian International Centre for Theoretical Physics, г. Нур-Султан, Казахстан

<sup>3</sup>Казахский национальный университет имени аль-Фараби, г.Алматы, Казахстан

### ИНФЛЯЦИЯ В $F(R,T)$ ГРАВИТАЦИИ С *f*-ЭССЕНЦИЕЙ

**Аннотация.** Модифицированные теории гравитации стали своего рода парадигмой в современной физике, поскольку они, кажется, решают несколько недостатков стандартной Общей теории относительности (ОТО), связанной с космологией, астрофизикой и квантовой теорией поля. Наиболее известные модифицированные теории гравитации являются  $F(R)$  и  $F(T)$  теории гравитации. Обобщение этих двух модифицированных теорий  $F(R)$  и  $F(T)$  гравитации впервые было предложено Мырзақуловым Ратбаем. В данной работе исследована неоднородная изотропная космологическая модель с фермионным полем *f*-эссенцией, действие которой имеет вид  $S_{RT} = \int d^4x \sqrt{-g} [F(R,T) + L_m]$ , где  $R$  – скаляр кривизны, а  $T$  – скаляр кручения,  $L_m$  – лагранжиан *f*-эссенции. Детально изучается частный случай при  $F(R,T) = \alpha R + \beta T$ , получены параметры, описывающие текущее ускоренное расширение Вселенной. Определён вид лагранжиана *f*-эссенции данной модели. Представленные результаты показывают, что  $F(R,T)$  гравитация с *f*-эссенцией может описывать инфляцию в ранней эволюции Вселенной.

В этой работе рассмотрена космологическая модель с фермионным полем и с неканоническим кинетическим членом (*f*-эссенция). Найдены некоторые точные решения модели и проверяем влияние такого гравитационно-фермионного взаимодействия на наблюдаемое ускоренное расширение Вселенной. Далее, обсуждаются космологические применения точных моделей. Также модель *f*-эссенции не имеет решения типа статической Вселенной Эйнштейна. Рассмотрено модифицированная  $F(R,T)$  гравитация с *f*-эссенцией. Было получены уравнения движения и рассмотрен инфляционный период ранней Вселенной. Для описания инфляционного периода, были определены вид параметра Хаббла и параметр медленного скатывания.

**Ключевые слова:** Вселенная, ускоренное расширение, инфляция, космология, *f*-эссенция, параметр медленного скатывания, метрика Фридмана-Робертсона-Уокера,  $F(R,T)$  - гравитация.

**Information about authors:**

Myrzakul Shynaray Ratbaykyzy, Candidate of Physical and Mathematical Sciences, PhD in Physics, L.N. Gumilyov Eurasian National University, «Ratbay Myrzakulov Eurasian International Centre for Theoretical Physics» LLP, Nur-Sultan, Kazakhstan, e-mail: srmyrzakul@gmail.com, ORCID: <https://orcid.org/0000-0003-0044-7323>;

Myrzakulov Yerlan Muratbayevich, PhD in Physics, L.N. Gumilyov Eurasian National University, «Ratbay Myrzakulov Eurasian International Centre for Theoretical Physics» LLP, Nur-Sultan, Kazakhstan, e-mail: srmyrzakul@gmail.com, ORCID: <https://orcid.org/0000-0003-0160-0422>;

Arzymbetova Moldir, doctoral student in physics, Al-Farabi Kazakh National University, e-mail: arzymbetova.moldir@gmail.com, ORCID: <https://orcid.org/0000-0002-4589-2941>

**REFERENCES**

- [1] Myrzakulov R. (2012) F(T) gravity and k-essence. *General Relativity and Gravitation*, 44, 3059-3080. DOI: [10.1007/s10714-012-1439-z](https://doi.org/10.1007/s10714-012-1439-z)
- [2] Myrzakulov R. (2012) Dark Energy in F(R,T) Gravity. <https://arxiv.org/abs/1205.5266>
- [3] Myrzakulov R. (2012) FRW Cosmology in F(R,T) gravity. *The European Physical Journal C*, 72:11, 2203. DOI: [10.1140/epjc/s10052-012-2203-y](https://doi.org/10.1140/epjc/s10052-012-2203-y)
- [4] Sharif M., Rani S., Myrzakulov R. (2013) Analysis of F(R,T) Gravity Models Through Energy Conditions, *The European Physical Journal Plus*, 128, 123. DOI: [10.1140/epjp/i2013-13123-0](https://doi.org/10.1140/epjp/i2013-13123-0)
- [5] Yi-Fu Cai, Capozziello S., De Laurentis M., Saridakis E.N. (2016) f(T) teleparallel gravity and cosmology, *Reports on Progress in Physics*, 79:4, 106901. DOI: [10.1088/0034-4885/79/10/106901](https://doi.org/10.1088/0034-4885/79/10/106901)
- [6] Capozziello S., De Laurentis M., Myrzakulov R. (2015) Noether Symmetry Approach for teleparallel-curvature cosmology, *International Journal of Geometric Methods in Modern Physics*, 12:9, 1550095. DOI: [10.1142/S0219887815500954](https://doi.org/10.1142/S0219887815500954)
- [7] Myrzakulov R. (2010) Cosmological models with non-canonical scalar and fermion fields: k-essence, f-essence and g-essence. [arXiv:1011.4337]
- [8] Benisty D., Guendelman E.I., Saridakis E.N. (2020) The scale factor potential approach to inflation, *The European Physical Journal C*, 80, 480. DOI: [10.1140/epjc/s10052-020-8054-z](https://doi.org/10.1140/epjc/s10052-020-8054-z)
- [9] Martin J., Ringeval C., Vennin V. (2014) *Encyclopædia Inflationaris, Physics of the Dark Universe*, 5-6, 75-235. DOI: [10.1016/j.dark.2014.01.003](https://doi.org/10.1016/j.dark.2014.01.003)
- [10] Amani Ali R., Samiee-Nouri A. (2015) Logarithmic entropy corrected holographic dark energy with F(R,T) gravity, *Communications in Theoretical Physics*, 64:4, 485-490. DOI: [10.1088/0253-6102/64/4/485](https://doi.org/10.1088/0253-6102/64/4/485)
- [11] Amani Ali R., Dehneshin S.L. (2015) Interacting F(R,T) gravity with modified Chaplygin gas, *Canadian Journal of Physics*, 93:12, 1453-1459. DOI: [10.1139/cjp-2015-0024](https://doi.org/10.1139/cjp-2015-0024)
- [12] Salti M., Korunur M., Acikgoz I., Pirinccioglu N., Binbay F. (2018) f(T,R) theory of gravity, *International Journal of Modern Physics D*, 27:05, 1850062. DOI: [10.1142/S0218271818500621](https://doi.org/10.1142/S0218271818500621)
- [13] Chattopadhyay S. (2013) A Study on the Interacting Ricci Dark Energy in f(R,T) Gravity, *Proceedings of the National Academy of Sciences, India Section A: Physical Sciences*, 84:1, 87-93. DOI [10.1007/s40010-013-0090-8](https://doi.org/10.1007/s40010-013-0090-8)
- [14] Meirbekov B., Bauyrzhan G., Yerzhanov K. (2018) FRW cosmology of Myrzakulov gravity with k-essence. DOI: [10.13140/RG.2.2.33421.64481](https://doi.org/10.13140/RG.2.2.33421.64481)
- [15] Meirbekov B., Razina O., Tsyba P. (2018) Cosmological solutions of Myrzakulov gravity with fermionic fields. [https://www.researchgate.net/publication/326200230\\_Cosmological\\_solutions\\_in\\_Myrzakulov\\_gravity\\_model\\_with\\_a\\_fermionic\\_field](https://www.researchgate.net/publication/326200230_Cosmological_solutions_in_Myrzakulov_gravity_model_with_a_fermionic_field)
- [16] Meirbekov B., Myrzakulov Y., Yerzhanov K. (2018) Myrzakulov gravity with f-essence and its FRW cosmology. [https://www.researchgate.net/publication/326200767\\_Myrzakulov\\_gravity\\_with\\_f-essence\\_and\\_its\\_FRW\\_cosmology](https://www.researchgate.net/publication/326200767_Myrzakulov_gravity_with_f-essence_and_its_FRW_cosmology)

## **Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www.nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

**ISSN 2518-1726 (Online), ISSN 1991-346X (Print)**

Редакторы: *М. С. Ахметова, Д. С. Аленов, А. Ахметова*  
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 22.09.2020.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.  
7,75 п.л. Тираж 300. Заказ 5.